

Interactive comment on “Efficient retention of mud drives land building on the Mississippi Delta plain” by Christopher R. Esposito et al.

Anonymous Referee #3

Received and published: 14 April 2017

This manuscript concerns the important issue of sediment retention, particular that of the finer mud fraction, on floodplains. As set out in the manuscript, the retention of sediment on floodplains is vital for the management of deltaic and wetland environments. Furthermore, previous work on this subject has focussed intently on the sand, or coarser, fractions of the sediment load. This work adds a novel and important re-focus on the finer fraction which is shown here to represent ~95% of the sediment volume in the studied crevasse-splay system. As such this manuscript is an important addition to the literature. It is well written and clearly presented. However, I think the framing of the work, and some of the comparisons to other splay systems needs revisiting, or more justification.

The splay system under consideration in the manuscript is one that developed in a swamp environment (line 27) and which is characteristic of several such features found

[Printer-friendly version](#)

[Discussion paper](#)



along the Mississippi Delta which are key to the maintenance of the landward portions of the MD (line 28 - 30). These splay systems are not developing in open water basins at the marine edges of the delta where they are being effected by tides and waves in the Gulf of Mexico, rather they likely develop in empty basins where the main control is topography and fluvial inputs and where mud is potentially rapidly deposited (rather than resuspended) as sediment enters the basin. It is therefore interesting that the main area of comparison is with the Wax Lake Delta, which is a coastal deltaic deposit developing at the marine interface and is influenced by tides and waves. It seems to me that these two systems are not directly comparable as the marine influences are likely to effect the processes of deposition and resuspension which occur at the splay edges, thus impacting the proportions and locations of the sediment fractions under study. It would be more useful to compare the findings of this work to other terrestrial/fluvial crevasse-splay systems found along the main channels of the "inland" MD rather than deltaic deposits at the marine interface.

It would also be interesting to have a discussion around the errors in the authors estimates of ratings curves used to estimate sediment load and fractionations, and also the water levels in the assumed trunk channels from which the splay emanates. Is there any hysteresis displayed in the sediment ratings curves that could impact upon the functioning of the splays? How good is the fit of the ratings curves and what the propagated errors through the estimates of sediment concentration and discharge? The estimates of SRE the authors report are likely to vary with these and it would be useful to have an idea of the sensitivity of the metrics used by the authors to characterise the ACS to these input parameters.

Interactive comment on Earth Surf. Dynam. Discuss., doi:10.5194/esurf-2017-5, 2017.

Printer-friendly version

Discussion paper

