

Interactive comment on “Stable water isotopes in the MITgcm (checkpoint 64w)” by Rike Völpel et al.

Anonymous Referee #1

Received and published: 9 March 2017

General comments Völpel et al., present the first results of the implementation of oxygen stable isotopes in the ocean general circulation model MITgcm. They compare the results of an equilibrium simulation under pre-industrial conditions to observational oxygen stable isotopes data and late Holocene data from planktonic foraminifera. They discuss the accordance or discrepancy of this data-model comparison. I find always very interesting the implementation of water stable isotopes in a climate model as it will offer wide prospects for simulations of past climate. The manuscript reads well overall, some figure captions or figures could be improved. I have several concerns that I would like to see addressed before the publication of the manuscript. I therefore recommend major revision of the present manuscript. Below are my specific comments.

Specific comments “Part2.1 Ocean model in lines 28-30” Can the authors explained in more details the rescaling of the vertical coordinate?

“Part 2.2 in lines 27, Note that the prescribed atmospheric forcing fields obtained from

[Printer-friendly version](#)

[Discussion paper](#)



the PI ocean state estimate by Kurahashi-Nakamura et al., (submitted) and the corresponding isotopic fluxes are not entirely consistent and might introduce an error in our model simulation". The authors refer to unpublished results here. They should show some results that indicate what could be the error and if the use of ratio of the isotopic content indeed minimize the uncertainty.

"Part 2.3.1 in line 30". The authors compare long-term mean monthly value with GISS sample. This is indeed better than to compare with the annual mean isotope values. However, in the rest of the text it is difficult to know when the comparison is based on monthly value or annual. This is also not clear in the different figures and captions on the manuscript.

"part 3.2 line21" The number of measurements for dD is rather small. According to the GISS database there is more than 1000 data points in dD. This is indeed more reduce than for the d18Osw but enough to realize a data-model comparison. Rather, the authors can mentioned that they choose to focus on the d18O and will work on the dD in the future.

"Figure 3" Is it the annual or monthly value that are plotted for the model? Is it the surface data that are compared to the average 50m of the model or the data between 0 and 50 m? What could be the error associated if this is the surface data versus the average 50m?

"part 3.2 lines 3-4, the subtropical gyres are less enriched. ..." There is also a discrepancy for the Mediterranean Sea. What is the reason for such discrepancy in the subtropical gyres and Mediterranean region?

"Figures 7 and 8" What is the depth used in the model (50 m?), is it annual or monthly?

"Part 4.1 lines 20 to 30 and Figure 9" A zoom on the artic region would be very helpful here. The isotopic values for rivers discussed in the text could eventually be added to this figure of the artic region.

[Printer-friendly version](#)[Discussion paper](#)

“Part 4.3: Planktonic foraminiferal d18Oc” When reading part 4.3 it seems that the main discrepancy between data and model results is because of the gametogenic calcification of foraminifera and so that paleotemperature equations derived from plankton-tow data are more appropriate to reconstruct surface water conditions than the commonly used paleotemperature equations like Shackleton (1974) or Kim and O’Neil (1997). This discussion is extremely interesting for paleoceanographic studies. Nonetheless I find that all the potential factors that can affect the d18Oc and so the data-model comparison and mismatch are not developed enough. Indeed, the temperature bias in the model (2°C or more in some regions, see figure 1) can affect significantly the d18Oc reconstruction with the model. Similarly, the bias in d18Osw could contribute significantly to this “biased towards lower values”. For example, the d18Osw is 0.4‰ too depleted in the model in comparison to data in the tropics (see part 3.2) and 0.9‰ too enriched in the Arctic Ocean (see part 3.2). These biases can affect the d18Ocalcite reconstruction and comparison. Also, it seems that the shift on figure 8a is more important for tropical species than for polar species. The data-model agreement or disagreement seems different depending on the oceanic region (or species considered). So I recommend to the authors to realize a data-model comparison for the d18Oc for the different species of foraminifera separately. This analysis is important not only to try to discuss the oceanic region separately but also because other factors can affect each species of foraminifera in a different way. The seasonality is one of these important factors. Although there is one sentence in part 4.3 that mentions that “seasonality could be a problem and is not considered” it would be interesting to estimate how much bias could be introduced by such inconsideration. One way to do that could be to calculate the simulated seasonal amplitude for ocean calcite d18O in the model. It could be that the “biased towards lower values” is partly or totally explained by a distortion of the foraminifera flux towards a specific season or period than the annual mean. Similarly, the effect of vertical migration is not completely developed. The author discusses the gametogenic calcification that is indeed related to this effect of vertical migration but the different species that are grouped on Figure 8 have different depth habitats and

[Printer-friendly version](#)[Discussion paper](#)

this affect their $\delta^{18}\text{O}_c$. They can also change their depth habitat (for example during upwelling conditions). Again a data-model comparison for each species separately and with a different mean depth habitat of calcification would be interesting. The data on figure 8 are only presented for the first 50 m (although not clearly indicated in the text or on the Figure 8 caption). Although it will be difficult to examine the results for the very surface only (because of the grid of the model), the authors can investigate how the integration of the results for deeper water depth affect the data-model comparison.

The authors also suggest that the more enriched $\delta^{18}\text{O}_c$ values obtained with the equation of Shackleton (1974) is because this equation is based on *Uvigerina* spp shells that are relatively enriched in ^{18}O . In fact, Shackleton (1974) proposed that *Uvigerina peregrina* is in isotopic equilibrium with seawater contrary to *Cibicides*. On the contrary, Bemis et al. (1998) (not cited in the discussion) suggested that *Cibicides* might also calcify in isotopic equilibrium and that the heavier $\delta^{18}\text{O}$ values of *Uvigerina* are due to calcification at lower porewater pH. More recently, Marchitto et al., 2014 (also not cited in the discussion) investigated this difference in more details. Their results agree with Bemis et al. (1998) that *Cibicidoides* and *Planulina* appear to be closer to isotopic equilibrium (as represented by the Kim and O'Neil (1997) inorganic precipitates, which is also a matter of debate) than *Uvigerina*, although scatter in the measurements limits their confidence in this statement. They also recommend that *Uvigerina* $\delta^{18}\text{O}$ be adjusted to the *Cibicidoides* scale by subtracting 0.47‰ and not 0.64‰. They were also unable to discern an impact of bottom water pH on benthic foraminiferal $\delta^{18}\text{O}$, but they speculate that *Uvigerina*'s deviation from equilibrium could be explained by admixture of rapidly-precipitated non-equilibrium CaCO_3 that would be subject to a pH influence. So, to my knowledge, the question as to why the $\delta^{18}\text{O}$ of *Uvigerina* and *Cibicides* are different remains. The question of the pH influence is also not discussed for planktonic foraminifera whereas it could also have a significant effect on the oxygen isotopic composition (Bijma et al., 1999; Zeebe 1999). This pH effect could be a primary mechanism to explain the differences between the equations (Mulitza et al., 2004). Again, the pH effect will be different with the latitudes and so it is important to discuss the species

[Printer-friendly version](#)[Discussion paper](#)

(that are associated to different oceanic regions) separately.

To resume, I like the discussion in part 4.3, this is of strong interest for paleoceanographic studies and the gametogenic calcification is a factor that certainly need to be considered. Nonetheless, the authors do not discuss in details all the factors and biases that can affect the $d_{18}O_c$ of their data-model comparison. For each foraminifera specie, how the bias in $d_{18}O_{sw}$ in the model, the depth use in the model to generate the $d_{18}O_c$ signal, the seasonality and vertical migration and the pH can affect the $d_{18}O_c$ signal modelled and the comparison with data? At the end, if we consider all these factors and potential biases for $d_{18}O_c$ and the data-model comparison, can the authors really conclude that the differences between data and model is mainly linked to gametogenic calcification? If the authors cannot confirm their hypothesis in a revised version, they should also reformulate this conclusion from the abstract and conclusion part.

References: Bemis, B. E., Spero, H. J., Bijma, J., & Lea, D. W. (1998). Reevaluation of the oxygen isotopic composition of planktonic foraminifera: Experimental results and revised paleotemperature equations. *Paleoceanography*, 13(2), 150-160. Bijma, J., Spero, H. J., & Lea, D. W. (1999). Reassessing foraminiferal stable isotope geochemistry: Impact of the oceanic carbonate system (experimental results). In *Use of proxies in paleoceanography* (pp. 489-512). Springer Berlin Heidelberg.

Kim, S. T., & O'Neil, J. R. (1997). Equilibrium and nonequilibrium oxygen isotope effects in synthetic carbonates. *Geochimica et Cosmochimica Acta*, 61(16), 3461-3475.

Marchitto, T. M., Curry, W. B., Lynch-Stieglitz, J., Bryan, S. P., Cobb, K. M., & Lund, D. C. (2014). Improved oxygen isotope temperature calibrations for cosmopolitan benthic foraminifera. *Geochimica et Cosmochimica Acta*, 130, 1-11.

Mulitza, S., Donner, B., Fischer, G., Paul, A., Pätzold, J., Rühlemann, C., and Segl, M.: The South Atlantic oxygen-isotope record of planktic foraminifera, in: *The South*

[Printer-friendly version](#)

[Discussion paper](#)



Atlantic in the Late Quaternary: Reconstruction of Mass Budget and Current Systems edited by Fischer, G. and Wefer, G., 121-142, Springer, New York, 2004.

Shackleton, N. J.: Attainment of isotopic equilibrium between ocean water and the benthonic foraminifera genus *Uvigerina*: isotopic changes in the ocean during the last glacial, Gif-sur-Yvette, Colloque international du CNRS, 219, 203-210, 1974.

Zeebe, R. E. (1999). An explanation of the effect of seawater carbonate concentration on foraminiferal oxygen isotopes. *Geochimica et Cosmochimica Acta*, 63(13), 2001-2007.

[Interactive comment on Geosci. Model Dev. Discuss., doi:10.5194/gmd-2017-7, 2017.](#)

GMDD

[Interactive
comment](#)

[Printer-friendly version](#)

[Discussion paper](#)

