

Interactive comment on “Statistical characteristics of convective wind gusts in Germany” by Susanna Mohr et al.

G. R. Solari (Referee)

giovanni.solari@unige.it

Received and published: 20 February 2017

Separating the occurrences and measurements of different Aeolian phenomena such as synoptic cyclones, thunderstorms, tornadoes and so on is a key topic of modern wind engineering in order to perform distinct statistical analysis, to extract the main statistical parameters related to each phenomenon, and to build wind field models suitable to represent the wind loading and response of structures. Merging these separate evaluations in a unitary formulation is a further aim still in the embryonic stage.

This paper provides very interesting and new information on several aspects in the above framework, thus represents a useful and pertinent contribution to the advance of the knowledge in this field. In its whole I appreciate it and support its publication.

This paper contains a broad literature both in the fields of atmospheric sciences and

[Printer-friendly version](#)

[Discussion paper](#)



wind engineering, perhaps a little biased towards the first field. Despite this I believe that some relevant contributions to this topic are not considered and some choices inherent the methods herein applied seem to be based on a limited view of some previous contributions. Under this point of view, without changing anything in the substance of this paper, I believe that a wider critical discussion on the advantages and shortcomings related to such choices may improve the quality of this paper and inspire future step forwards.

More in detail, I recommend Authors to take into account the following remarks and observations:

- Section 1: at least two additional references should be considered. The first [1] is the fundamental paper that in 1977 introduced the concept of mixed wind climate and the idea of processing separately the statistical analysis of different wind phenomena. The second [2], published in 2002, deals with the same topic of the present paper just with reference to Germany. A comparison with previous methods and results is recommended.
- Section 2: I am quite doubtful on the decision of restricting analyses to the summer half-year. In my experience thunderstorm events are concentrated in this part of the year but are present also, in minor proportion, all over the year. Restricting analyses to a period is even more dangerous considering the aim of performing a statistical analysis of the extreme wind speed. Unavoidably this produces underestimated results. I suggest to revise this choice in next contributions.
- Section 2.1: Authors base their analyses on the daily peak and subsequent mean wind speeds on 10-min and 1-h periods. They also use pressure measurements. A very similar approach is used in Uruguay and described in [3]. I suggest to examine this contribution.
- Section 2.1: Also in the light of the occurrence of gust factors in the order of 6-10, I suggest Authors to consider the possibility that some peak values in the database may be wrong [4]. The potential presence of some mistakes and the difficulty of recognizing them is a major shortcoming of this kind of analyses, where the control is very good in terms on mean values but almost impossible with reference to single peaks.
- Section 2.3: I understand that Authors have probably no other opportunity than this use of lightning data. In my experience the presence of cloud-to-

[Printer-friendly version](#)[Discussion paper](#)

cloud lightning not detected by measurements may provide some relevant drawback. I verified this by comparing similar lightning data with high-sampling velocity records.

â€” Section 2.4: The problem of the separation of different wind events is a key topic because any mistake in this stage may compromise the quality of further evaluations. I suggest Authors to dedicate a few more words to this problem for instance using a citation to Lombardo et al. (2009) (included in references but not cited here) and to [5].

â€” Section 2.5: Authors speak of GEV and POT/GPD and make the choice of using POT/GPD. This is fine but again, without changing the substance of this paper, this topic is a "world" that may necessitate a some more "delicate" approach. First of all the use of POT/GPD is widely supported by some Authors but drastically opposed by others. Ref. [5], for instance, is fully devoted to demonstrate that this method is wrong or at least unreliable. Our research group recently published a series of papers based on long-term Monte Carlo simulations [6,7] that confirms the limited reliability of the POT/GPD technique and arrives to the conclusion that the Process Analysis [8] (probably not easy to apply to thunderstorms) and the Penultimate distribution [9,10] are the best methods.

â€” Section 2.5: At the end of this section Authors write "that the differences between the return values estimated by both methods are considerably smaller than the uncertainties of the method itself". This is absolutely correct with reference to return periods in the order of the number of years of available data, for instance 20-50 years. Structural safety, however, needs evaluations extrapolated to return periods in the order of 500-1000 years. Here, different methods lead to divergent results [9,10,11].

â€” Section 3.1: Authors write: "we considered every single measurement at each station, which means that one event can be recorded on two or more stations". I think that this sentence may result misleading. Downburst are phenomenon with a radius of a few km. It is almost impossible that the same downburst may be detected by two stations of this network. The situation is different if Authors refer to the large scale wind event that generates downbursts. This point should be clarified.

â€” Section 3.3: The last sentence deserves a citation to Authors that first expressed this concept [1].

â€” Section 3.4: Line 11. The dependence

[Printer-friendly version](#)[Discussion paper](#)

of the gust factor on the averaging period is discussed also by Solari et al. (2015).
â€” Section 3.4: Line 20. I do not agree on the sentence according to which “turbulent factors (to replace with gust factors ?) fluctuate usually between 1.2 and 2.3”. Such a large variability necessarily depends not only on the roughness length but even more on the stability conditions. If wind is intense and of synoptic type, then atmosphere is neutrally stratified and the gust factor may vary between 1.25 and 1.75 with an average value around 1.5. In my opinion gust factors in the order of 1.75-2.3 may be ascribed to unstable conditions and intermediate events between large scale depressions and mesoscale downbursts.

Independently of the above remarks, that I wrote in a fully constructive spirit, I confirm my appreciation towards this contribution and that I consider it appropriate for publication. I hope that Authors may consider or discuss my remarks.

I suggest that this paper is accepted subject to minor revisions but I believe that a quick re-review may be useful.

Giovanni Solari Department of Civil, Chemical and Environmental Engineering Polytechnic School, University of Genoa, Italy

References
Gomes L, Vickery BJ (1977/1978). Extreme wind speeds in mixed climates. *J Ind Aerod*, 2, 331-344.
Kasperski M (2002). A new wind zone map of Germany. *J Wind Eng Ind Aerod*, 90, 1271-1287.
Duranona V (2015). The significance of non-synoptic winds in the extreme wind climate of Uruguay. *Proc 14th Int Conf on Wind Engineering*, Porto Alegre, Brasil.
Cook NJ (2014) Review of errors in archived wind data. *Weather* 69: 72-78.
De Gaetano P, Repetto MP, Repetto T, Solari G (2013). Separation and classification of extreme wind events from anemometric data. *J Wind Eng Ind Aerod*, 126, 132-143.
Harris, R.I., 2005. Generalized Pareto methods for wind extremes. Useful tool or mathematical mirage? *JWEIA* 93, 897-918.
Torrielli A, Repetto MP, Solari G (2013). Extreme wind speeds from long-term synthetic records, *J Wind Eng Ind Aerod*, 115, 22-38.

Torrielli A, Repetto MP, Solari G (2014). A refined analysis and simulation of the wind speed macro-meteorological components. *J Wind Eng Ind Aerod*, 132, 54-65.

Gomes, L., Vickery, B.J., 1977. On the prediction of extreme wind speeds from the parent distribution. *J. Ind. Aerodyn.* 2, 21-36.

Cook, J., Harris, I., 2004. Exact and general FT1 penultimate distributions of extreme wind speeds drawn from tail-equivalent Weibull parents. *Struct. Saf.* 26, 391-420.

Cook, J., Harris, I., 2008. Postscript to "Exact and general FT1 penultimate distributions of extreme wind speeds drawn from tail-equivalent Weibull parents". *Struct. Saf* 30, 1-10.

Lagomarsino, S., Piccardo, G., Solari G., 1992. Statistical analysis of high return period wind speeds, *JWEIA* 41, 485-496.

[Interactive comment on Nat. Hazards Earth Syst. Sci. Discuss., doi:10.5194/nhess-2016-402, 2017.](#)

[Printer-friendly version](#)

[Discussion paper](#)

