

Atmos. Chem. Phys. Discuss., author comment AC2  
<https://doi.org/10.5194/acp-2020-981-AC2>, 2021  
© Author(s) 2021. This work is distributed under  
the Creative Commons Attribution 4.0 License.

## Reply on RC2

Xuewu Fu et al.

---

Author comment on "Isotopic compositions of atmospheric total gaseous mercury in 10 Chinese cities and implications for land surface emissions" by Xuewu Fu et al., Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2020-981-AC2>, 2021

---

## Response to Anonymous Referee #2

RC- Reviewer's Comments; AC – Authors' Response Comments

**RC1:** The manuscript presents data of TGM concentrations and their isotopic compositions in 10 large cities, many of them considered mega cities, in China. Documentation of such data in open literature is valuable, the data quality is good and the scientific presentation is sound. The primary weakness of the manuscript is a deterministic scientific argument based on the relatively scattered data, which is difficult for a study of this nature since the sampling was performed at different time (and perhaps by different sets of researchers) at different locations. Given the level of data scattering, it is somewhat uncertain to provide a clear scientific finding, which seemed to be the main criticisms of Reviewer #3 in the previous round of peer review.

Other than a lack of deterministic scientific conclusion, the reasoning and interpretation of data appear to be sound. One interesting feature of the conclusive remarks made by the author group is the attribution of mercury source and the TGM concentration variation primarily to soil evasion, which is somewhat counter-intuitive to the intensive human activities in large cities. The measured TGM concentrations in those cities are highly elevated (Figure 4). Previous work has attributed the elevated concentrations to human activities and the seasonal variation to meteorological factors, which seems reasonable and intuitive. Although this work presents additional mercury isotope data, the level of data scattering does not seem justified to rule out the past attribution to human activities and meteorology. Perhaps the authors should at least make an attempt to strengthen the arguments described in their conclusion.

**AC1:** We greatly appreciate the reviewer for recognizing the value of our study and for providing constructive comments, which would help us improve the manuscript.

We understand the concerns of this reviewer regarding the interpretation of the factors influencing the TGM concentrations and isotopic compositions in our investigated sites. Actually, we have not ruled out the effect of anthropogenic emissions and meteorology on our measurements. For example, we roughly estimated that primary anthropogenic emission is of similar importance as land surface emission in atmospheric Hg emission

budget in our investigated cities (line 371 in the revised manuscript). In this study, we are trying to use multiple evidences (including the unique Hg isotope data) to show that the land surface emission likely played a dominant role in regulating the seasonal variations in TGM isotopic compositions and concentrations. We have strengthened this arguments by a comparison of GEM emission fluxes between land surface and primary anthropogenic sources in line 374-379 in the revised manuscript, which reads: "However, our estimate is overall consistent with pervious studies on GEM emission fluxes from land surfaces and anthropogenic sources in Chinese urban areas. For example, Previous studies on GEM emission fluxes from urban surfaces in China showed a mean value of  $83.2 \pm 170 \text{ ng m}^{-2} \text{ h}^{-1}$  ( $1\sigma$ ,  $n = 39$ ) (Fu et al., 2012;Feng et al., 2005;Wang et al., 2006;Fang et al., 2004), which was relatively higher than the mean anthropogenic GEM flux ( $48.4 \pm 48.1 \text{ ng m}^{-2} \text{ h}^{-1}$ ,  $1\sigma$ ,  $n = 10$ ) in the ten investigated cities (Table S5) (AMAP/UNEP, 2013)".

There are also several other areas that can use additional clarity:

**RC2:** The selection of the ten city sites needs to be justified and the characteristics can be more detailed. Was the selection by design or by incident? If it is by design, discussion should be provided for the intended scientific goals. If it is by incident, discussion should be provided to argue why the data collected from the 10 sites can sufficiently support the conclusion.

**AC2:** These cities were selected by design, and the intended scientific goals are added in line 132-137 in the revised manuscript, which reads: "These cities are located in different geographical regions of China, which were potentially characterized by specific source emission patterns, climate, and atmospheric chemistry. The designated investigations in these cities may therefore provide a comprehensive information on the variations of TGM concentrations and isotopic compositions in mega cities of China, and help to explore the major factors influencing the atmospheric Hg in Chinese cities".

**RC3:** Since the samples were collected at different times and locations where the chemistry of various urban airshed could be substantially different such that the samplers may behave inconsistently. Based on the description in the method section, it seems that the sampling was not duplicated but the analysis was repeated. Some discussion in regard to the consistency of the carbon trap samplers will ensure the confidence on the data quality.

**AC3:** We are confidential that the collections of TGM using the carbon trap would not be biased by various atmospheric chemistry. This method has been used successfully in free troposphere, marine boundary layer, arctic, and forests by many previous studies (Fu et al., 2019;Fu et al., 2016;Obrist et al., 2017;Yu et al., 2020), and the data quality have been proven by these studies. In the present study, we investigated the blanks ( $n = 27$ ), breakthrough ( $n = 10$ ), and standard addition of Hg(0) vapor to carbon trap ( $n = 11$ ), which showed good results (more details in Section 2.3 and 2.4) and suggest our carbon trap method could measure the TGM concentrations and isotopic composition precisely.

**RC4:** The 10 city sites have drastically different meteorological patterns other than the generic seasonal patterns described in the manuscript. It is possible that there are local processes forcing the observed isotopic characteristics? This is not clear in the manuscript.

**AC4:** We agree that the local meteorological patterns would be a potential factor in some

cities, but we suppose they are not a dominant cause to drive the consistent seasonal variations in the ten investigated cities. The reason of this argument is added in line 421-429 in the revised manuscript, which reads: "Prevailing wind directions during the wintertime and summertime sampling campaigns were similar Jinan, Lanzhou, Zhengzhou, and Shanghai, but were different in other remaining cities (Figure S6). Variations in predominant wind directions would change the relationships between receptor and regional anthropogenic emissions, which could further influence the TGM levels and isotopic compositions in these cities. Given the similarity in wintertime and summertime prevailing wind directions in some cities and consistent summertime lower CO concentrations in most cities, it is postulated that the variations in local anthropogenic emissions and transport of regional anthropogenic emissions were not likely the main cause for the seasonal variations in TGM concentrations and isotopic compositions".

**RC5:** It will be useful if the authors can specify what statistical criteria is considered significant for using the relatively scattered data to draw the conclusion.

**AC5:** The statistical method used in this study is specified in line 266-267 in the revised manuscript, which reads: "Linear regression analysis was performed with IBM SPSS Statistics using the forced entry method".

**RC6:** Overall, the manuscript is considered acceptable after revisions on the scientific arguments and editorial improvements.

**AC6:** The reviewers' and editorial comments have been addressed and revisions have been made accordingly.

## References:

Fang, F. M., Wang, Q. C., and Li, J. F.: Urban environmental mercury in Changchun, a metropolitan city in Northeastern China: source, cycle, and fate, *Sci Total Environ*, 330, 159-170, DOI 10.1016/j.scitotenv.2004.04.006, 2004.

Feng, X. B., Wang, S. F., Qiu, G. A., Hou, Y. M., and Tang, S. L.: Total gaseous mercury emissions from soil in Guiyang, Guizhou, China, *J Geophys Res-Atmos*, 110, Artn D14306, Doi 10.1029/2004jd005643, 2005.

Fu, X., Zhang, H., Liu, C., Zhang, H., Lin, C.-J., and Feng, X.: Significant Seasonal Variations in Isotopic Composition of Atmospheric Total Gaseous Mercury at Forest Sites in China Caused by Vegetation and Mercury Sources, *Environmental Science & Technology*, 53, 13748-13756, 10.1021/acs.est.9b05016, 2019.

Fu, X. W., Feng, X. B., Zhang, H., Yu, B., and Chen, L. G.: Mercury emissions from natural surfaces highly impacted by human activities in Guangzhou province, South China, *Atmos Environ*, 54, 185-193, DOI 10.1016/j.atmosenv.2012.02.008, 2012.

Fu, X. W., Maruszczak, N., Wang, X., Gheusi, F., and Sonke, J. E.: Isotopic Composition of Gaseous Elemental Mercury in the Free Troposphere of the Pic du Midi Observatory, France, *Environmental Science & Technology*, 50, 5641-5650, 10.1021/acs.est.6b00033, 2016.

Obrist, D., Agnan, Y., Jiskra, M., Olson, C. L., Colegrove, D. P., Hueber, J., Moore, C. W., Sonke, J. E., and Helmig, D.: Tundra uptake of atmospheric elemental mercury drives Arctic mercury pollution, *Nature*, 547, 201-204, 10.1038/nature22997, 2017.

Wang, D. Y., He, L., Shi, X. J., Wei, S. Q., and Feng, X. B.: Release flux of mercury from different environmental surfaces in Chongqing, China, *Chemosphere*, 64, 1845-1854, DOI 10.1016/j.chemosphere.2006.01.054, 2006.

Yu, B., Yang, L., Wang, L. L., Liu, H. W., Xiao, C. L., Liang, Y., Liu, Q., Yin, Y. G., Hu, L. G., Shi, J. B., and Jiang, G. B.: New evidence for atmospheric mercury transformations in the marine boundary layer from stable mercury isotopes, *Atmos Chem Phys*, 20, 9713-9723, 10.5194/acp-20-9713-2020, 2020.