Atmospheric Chemistry and Physics Discussions

Atmos. Chem. Phys. Discuss., community comment CC1
https://doi.org/10.5194/acp-2021-123-CC1, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Reply on AC2

Jacob Carlin

Community comment on "Comment on "Review of experimental studies of secondary ice production" by Korolev and Leisner (2020)" by Vaughan T. J. Phillips et al., Atmos. Chem. Phys. Discuss., https://doi.org/10.5194/acp-2021-123-CC1, 2021

In support of Vaughan's hypothesis, I wanted to chime in with some evidence we have recently come across of potential sublimational SIP. Specific differential phase (K_{DP}) has recently been proposed as a potential marker of SIP zones as it is sensitive to large concentrations of anisotropic fragments, but this has so far been limited to SIP during riming (e.g., Grazioli et al. 2015; Sinclair et al. 2016; Kumjian and Lombardo 2017). However, looking at quasi-vertical profiles (QVPs) we have recently found bands of enhanced K_{DP} situated squarely within the sublimation zone in 9 of 12 cases of strong sublimation, coincident with gradients of Z and RH w.r.t. ice (taken from coincident RAP analyses). We are currently following up and exploring these signatures more quantitatively. However, given the observational laboratory evidence for sublimational SIP (e.g., Oraltay and Hallet 1989; Dong et al. 1994; Bacon et al. 1998), I contextualized these signatures with the existing literature by emulating Fig. 13 of Oraltay and Hallett (1989; reproduced below) and plotting the median K_{DP} as a function of RHi and ice bulb temperature. Note that Oraltay and Hallett (1989) only included down to 50% RHi and -6C for ice bulb temperatures, so my figure is an extrapolation and it is unclear whether their 70% RHi threshold is maintained at much colder ice bulb temperatures. Regardless, it is apparent that these regions of enhanced K_{DP} indicative of high concentrations of highly anisotropic particles are concentrated primarily in the region identified by Oraltay and Hallett (1989) as the favorable zone for sublimational breakup. The plot using mean K_{DP} was similar. These sorts of values are not likely to be due to aggregates which we believe characterize the majority of parent particles in the cases examined, and riming is precluded due to the degree of dry air present. Thus, although the use of K_{DP} is indirect evidence, under proper conditions (e.g., a continual and sufficiently high flux of parent particles into a sufficiently dry layer and favorable particle habits), I believe Vaughan's hypothesis of an equilibrium being established between fragment generation and fragment consumption to be plausible.

Jacob Carlin

Please also note the supplement to this comment:
https://acp.copernicus.org/preprints/acp-2021-123/acp-2021-123-CC1-supplement.pdf