

Atmos. Chem. Phys. Discuss., referee comment RC3
<https://doi.org/10.5194/acp-2021-133-RC3>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Comment on acp-2021-133

Anonymous Referee #3

Referee comment on "Identifying the sources of uncertainty in climate model simulations of solar radiation modification with the G6sulfur and G6solar Geoengineering Model Intercomparison Project (GeoMIP) simulations" by Daniele Visoni et al., Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2021-133-RC3>, 2021

This manuscript investigates the difference between simulations where solar radiation management is implemented as a reduction of incoming solar radiation and as an injection of stratospheric aerosols. The authors find that the two methods lead to a different response in the surface climate, and therefore are not equivalent in simulating geoengineering (as suggested in a few previous papers).

I have found this paper interesting, clear, and well written. I only have very minor comments.

L45: I supposed with "baseline simulation" the author means a simulation of a desirable climate reached via emission mitigation. My first understanding, though, was that a baseline simulation was one without geoengineering, which would not be correct because I believe G4 also required a simulation without geoengineering.

L49: The discriminant is not the presence of microphysics, but rather the presence of a sulfur cycle, or a way to represent the formation of new particles from sulfur dioxide. A microphysics representation allows for the evolution of particle size, which is one additional process beyond the formation of new particles.

L130: Shouldn't the reference be to Fig. 2f?

L132: what does it mean that the solar reduction and the AOD seem unrelated? In Fig. 2e you show a fit with R^2 of 0.72. The word "unrelated" should be replaced with a quantitative metric/

L142: eliminate "much"

L147: can you explain better how this is a feature of the current feedback controller? I understand that you refer to Tilmes et al. (2018) but I think it would be more complete if there was a not too specific explanation here without having to look for another paper

L152: I don't understand the connection between the sentences before and the conclusion that they explain why the AOD in 2050 is the same among all models.

L160: Does Solar produce any difference in the stratosphere? I imagine that reducing the solar flux would impact UV absorption by ozone (hence the temperature profile) as well as ozone concentrations

L189: I'm really not sure how meaningful is the comparison between Pinatubo and geoengineering simulations, and I would not include it at all. As you write here, this is a sustained injection, and you already mentioned several times that Visoni et al. showed the importance of the injection seasonality on the transport. Jones et al. (2016, doi:10.1002/2016JD025001) showed the dependence of the Pinatubo dispersal from the initial conditions. What I find incorrect is that this is the only "real" point of comparison. The only real point of comparison is a simulation of Pinatubo (which all models could do - and probably have done already) vs observations of Pinatubo. That would be informative with respect to the ability of each models to simulate the transport of stratospheric aerosols.

L200: Is this true also if you calculate the spread as % of multimodel mean? I am not sure that the better agreement isn't simply a result of the AOD values being smaller in 4c than 4a

L210: It would be interesting to show the optical properties (the extinction efficiency) used by each model and/or the simulated particle size distribution

L214: Canty et al. (2013, <https://doi.org/10.5194/acp-13-3997-2013>) finds a decrease of 0.14 K globally.

Fig. 7: I am a bit confused by the statistics. The ensembles contain two, three at the most, ensemble members. Did you perform a T test with only 2 realizations per experiment?