

Atmos. Chem. Phys. Discuss., referee comment RC1
<https://doi.org/10.5194/acp-2021-577-RC1>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Comment on acp-2021-577

Anonymous Referee #1

Referee comment on "On the evolution of sub- and super-saturated water uptake of secondary organic aerosol in chamber experiments from mixed precursors" by Yu Wang et al., Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2021-577-RC1>, 2021

General comments:

In this paper, authors conducted a series of laboratory studies on the evolution of sub- and super-saturated water uptake of secondary organic aerosol in chamber experiments from mixed precursors. The comparison of aerosol hygroscopicity under sub- and super-saturated conditions involves complex SOA characteristics are important for CCN predictions and understanding the RH effects (from sub-saturation to super saturation) on aerosol hygroscopicity and further improve aerosol hygroscopicity parameterizations. Therefore, the theme of this paper is scientifically meaningful, and the designed laboratory studies are also valid. However, more insightful analysis is required to better interpret the laboratory results.

Two main conclusions are drawn in this study.

The first one is that SOA composition played a second role in the kappa variations. I agree with the authors, and this is obvious considering that the well-known fact that inorganic aerosol is much more hygroscopic than SOA, thus, of course the SOA composition should play the second role. However, at the same MR_SOA/PM, considerable variations can be found, for example, at MR_SOA/PM ~0.4, the k_{HTDMA} varies from less than 0.2 to about 0.4, which demonstrates that the SOA compositions played significant roles in determining the apparent kappa. The authors should quantitatively derive the apparent/effective hygroscopicity parameter K_{OA} using the volume mixing rule under sub- and super-saturated conditions and discuss their differences with previous results (Kuang et al., 2020b), quantitatively present the relationships between K_{OA} and SOA oxidation levels under sub- and super-saturated conditions and discuss more about the controlling factors of K_{OA} variations based on AMS signals and VOC precursors (Wang et al., 2019). Otherwise, the authors are just talking about the sub- and super-saturated water uptake of mixed aerosol system includes organic and inorganic aerosols but not that of secondary organic aerosol, and is not consistent with the theme of the title.

The second one is " $K_{\text{HTDMA}}/k_{\text{CCN}}$ increased as a function of SOA mass fraction, independent of initial VOC concentrations and sources, the mean k_{HTDMA} can be 60% higher than k_{CCN} on average when aerosol fraction approached 0.8". This finding is quite interesting but the explanation is not very convincing. The authors conclude that this finding is possibly attributable to the non-ideality of solutes at different RH or different co-condensation of condensable organic vapors within the two instruments. At L397, the authors claim that water increase in CCN set-up is not favorable for co-condensation of semi-volatile vapors. However, the condensation of these vapors such as HNO_3 is influenced by both water vapor content and temperature. Although the temperature in CCN is higher than the temperature in HTDMA which seems not favorable for co-

condensation, however, the aerosol water content in the CCN is much higher than in the HTDMA due to the super-saturated conditions. Thus, I cannot agree upon this argument. As to the possible role of non-ideality, the authors cited two papers published by Brechtel and Kreidenweis (2000a) and Brechtel and Kreidenweis (2000b) and said "the interactions of inorganic ions and organic molecules can exert both positive and negative effects in the water uptake, depending on the organic fraction and inorganic species". The organic fraction and inorganic species are clear in this study, can authors perform some quantitative analysis? or at least deliver a clear qualitative result that the interactions of inorganic ions and organic molecules exerted a positive or a negative effect on the water uptake.

Specific comments:

L45-L50, just a suggestion: the first two sentences of the introduction lack continuity in logic, the authors jumped from aerosol-cloud interaction to the reliability of CCN prediction, I suggest a sentence that claim the accurate CCN prediction is essential for investigating aerosol-cloud interactions in climate models might be needed between the original two sentences.

L53, the SOA can be formed not just through gas-phase partitioning, but also aqueous-phase reactions (Ervens et al., 2011; Kuang et al., 2020a).

L58, about the role of organic aerosol hygroscopicity in climate and aerosol cloud-interactions, references such as (Liu and Wang, 2010; Rastak et al., 2017) might be better choices.

L164 more details about the DMA-CCN set-up should be given, for example, the detailed supersaturation points and time schedule

L175 Please report how the AMS vacuum aerodynamic diameter is converted to the mobility diameter, and estimate potential MR_SOA/PM uncertainties associated with this respect due to that obvious size-dependent chemical composition is observed.

L226, a smaller particle size is not specific (compare to which size range), please presents diameter range directly

Brechtel, F. J., and Kreidenweis, S. M.: Predicting Particle Critical Supersaturation from Hygroscopic Growth Measurements in the Humidified TDMA. Part I: Theory and Sensitivity

Studies, *Journal of the Atmospheric Sciences*, 57, 1854-1871, 10.1175/1520-0469(2000)057<1854:PPCSFH>2.0.CO;2, 2000a.

Brechtel, F. J., and Kreidenweis, S. M.: Predicting Particle Critical Supersaturation from Hygroscopic Growth Measurements in the Humidified TDMA. Part II: Laboratory and Ambient Studies, *Journal of the Atmospheric Sciences*, 57, 1872-1887, 10.1175/1520-0469(2000)057<1872:PPCSFH>2.0.CO;2, 2000b.

Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmos. Chem. Phys.*, 11, 11069-11102, 10.5194/acp-11-11069-2011, 2011.

Kuang, Y., He, Y., Xu, W., Yuan, B., Zhang, G., Ma, Z., Wu, C., Wang, C., Wang, S., Zhang, S., Tao, J., Ma, N., Su, H., Cheng, Y., Shao, M., and Sun, Y.: Photochemical Aqueous-Phase Reactions Induce Rapid Daytime Formation of Oxygenated Organic Aerosol on the North China Plain, *Environmental science & technology*, 10.1021/acs.est.9b06836, 2020a.

Kuang, Y., Xu, W., Tao, J., Ma, N., Zhao, C., and Shao, M.: A Review on Laboratory Studies and Field Measurements of Atmospheric Organic Aerosol Hygroscopicity and Its Parameterization Based on Oxidation Levels, *Current Pollution Reports*, 10.1007/s40726-020-00164-2, 2020b.

Liu, X., and Wang, J.: How important is organic aerosol hygroscopicity to aerosol indirect forcing?, *Environmental Research Letters*, 5, 044010, 10.1088/1748-9326/5/4/044010, 2010.

Rastak, N., Pajunoja, A., Acosta Navarro, J. C., Ma, J., Song, M., Partridge, D. G., Kirkevåg, A., Leong, Y., Hu, W. W., Taylor, N. F., Lambe, A., Cerully, K., Bougiatioti, A., Liu, P., Krejci, R., Petäjä, T., Percival, C., Davidovits, P., Worsnop, D. R., Ekman, A. M. L., Nenes, A., Martin, S., Jimenez, J. L., Collins, D. R., Topping, D. O., Bertram, A. K., Zuend, A., Virtanen, A., and Riipinen, I.: Microphysical explanation of the RH-dependent water affinity of biogenic organic aerosol and its importance for climate, *Geophysical Research Letters*, 44, 5167-5177, 10.1002/2017gl073056, 2017.

Wang, J., Shilling, J. E., Liu, J., Zelenyuk, A., Bell, D. M., Petters, M. D., Thalman, R., Mei, F., Zaveri, R. A., and Zheng, G.: Cloud droplet activation of secondary organic aerosol is mainly controlled by molecular weight, not water solubility, *Atmos. Chem. Phys.*, 19, 941-954, 10.5194/acp-19-941-2019, 2019.