

Atmos. Chem. Phys. Discuss., referee comment RC1
<https://doi.org/10.5194/acp-2021-728-RC1>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Comment on acp-2021-728

Anonymous Referee #2

Referee comment on "Formation of ice particles through nucleation in the mesosphere" by
Kyoko K. Tanaka et al., Atmos. Chem. Phys. Discuss.,
<https://doi.org/10.5194/acp-2021-728-RC1>, 2021

Formal review of the manuscript

Formation of ice particles through nucleation in the mesosphere

by K. Tanaka, I. Mann and Y. Kimura

The nucleation mechanism of polar mesospheric clouds has been a longstanding problem. At least since the late 1960s it has been debated whether homogeneous or heterogeneous nucleation is the dominant nucleation mechanism leading to the formation of ice particles under the extreme conditions of the polar summer mesopause (e.g., Witt 1969). While homogeneous nucleation had been deemed very unlikely given the contemporary knowledge on temperatures, water vapor mixing ratios and the (at that time only conceived) occurrence of meteoric smoke particles, observations of extreme gravity wave-induced temperature perturbations by Lübken et al. (2009) triggered Murray and Jensen (2010) to reinvestigate the problem. Based on (slightly modified) classical nucleation theory they concluded that homogeneous nucleation could indeed lead to the formation of amorphous solid water particles in the mesopause region if such extremely strong gravity wave-induced temperature perturbations (and hence cooling rates) occurred. However, they also found that if homogeneous nucleation had to compete with heterogeneous nucleation on meteoric smoke particles, the latter was more efficient and homogeneous nucleation became negligible.

In their current manuscript Tanaka and coauthors reconsider this problem based on the fact that the classical nucleation theory used in the work of Murray and Jensen is known to strongly disagree with laboratory observations for the case of water. Hence, Tanaka et al. apply a semi-phenomenological model which is known to be in much better agreement

with observations. This model shows a much higher free energy barrier for nucleation such that homogeneous nucleation of ice particles in the mesopause region would require unrealistically low temperatures, i.e., well below 100K. Hence, this nucleation pathway can be ruled out (because it contradicts observed temperatures) while heterogeneous nucleation is found to be feasible (in agreement with the recent groundbreaking laboratory measurements by Duft et al. 2019).

In all this is a sound study that contributes to the important fundamental problem of ice nucleation in the mesopause region. While the study of Murray and Jensen predicted homogeneous nucleation to possibly occur under extreme, but still conceivable conditions (extreme cooling rates, no competing meteoric smoke) the work by Tanaka et al. now clarifies that even under such extreme conditions homogeneous nucleation cannot be expected. This result certainly warrants publication.

My recommendation is hence to publish this work provided that the following mostly minor issues are properly addressed before publication:

The referencing in the introduction could be improved by referring to the original papers for the statements made. Here are my suggestions:

- line 16/17: original reference for noctilucent clouds: Jesse 1885; maybe also Vestine 1934

- line 20: the original reference for satellite-based PMC observations is Donahue et al 1972; a very good review until 2006 is DeLand et al., 2006.

- line 21: reference for particle sizes: Thomas and MacKay, 1985, von Cossart et al., 1999;

- line 23/24: well, this statement is not correct as it stands here: the ground based visual sightings of NLC actually do not show a unique trend as shown in Kirkwood and Stebel (2003); however, a trend is observed in the brightness of satellite-based PMC observations as presented in Thomas et al (2003) and updated in DeLand and Thomas (2015).

- line 25: while the reference to Lübken et al. (2018) is good, the original paper posing this hypothesis should also be mentioned, i.e., Thomas et al., Nature 1989.

- line 27: original reference on gravity wave-NLC-interaction: Witt, 1962.

- line 28: to my knowledge temperatures as low as 100K (and even lower) have only been reported in Lübken et al. (2009). Lübken 1999 is a climatology for mean temperatures at 69°N (from falling sphere measurements); Rapp et al. (2002) do show gravity wave perturbed temperature measurements in NLC but with minimum temperatures of 110K.

- line 33: homogeneous nucleation has only been considered feasible again (after many years during which it was regarded extremely unlikely) after Lübken et al. (2009) reported enormous temperature variability due to gravity waves (see their figures 9, 10 and 11). Until then the consensus in the community was that it was rather heterogeneous nucleation on meteoric smoke (see e.g., Rapp and Thomas 2006 for a discussion).

- line 39 and 40: The authors are mixing two things here: as reviewed in Rapp and Thomas, the stated species have been suggested in the literature as potential nuclei for mesospheric ice particle formation. However, not all of the stated species are candidates for the composition of meteoric smoke (e.g., proton hydrates, soot are independent of meteoric origin). Meteoric smoke composition is indeed discussed in Plane (2015). I recommend to have a look at this paper and change the sentence accordingly.

- Section 3: in order to put the results in perspective, it would be useful if the authors included a short section describing typical ranges of mesospheric variables like observed temperatures, water vapor mixing ratios or partial pressures, concentrations of meteoric smoke particles (e.g. from rocket borne observations), and cooling rates due to tides and gravity waves. This will help assessing the assumptions made and results achieved in the paper. In this context, the authors should clearly state if derived or used values are way outside of observed ranges.

- line 238/239: the authors should point out that cooling rates as low as $1\text{e-}6\text{ K s}^{-1}$ at initial temperature of 135K also corresponds to a completely unrealistic time that the nucleation would take. However, observations do show that PMSE (which are also evidence for ice particles, but already at times when they have not yet grown large enough to be optically detectable) form rapidly for example in updrafts of gravity waves (i.e., within minutes).

- line 235/236: These formulations are misleading. "the amount of water vapor present was 20 times higher at 145K than at 135K" – this certainly doesn't have anything to do with the atmosphere. In the atmosphere, the water vapor mixing ratio in the mesosphere is determined by transport across the tropical tropopause and oxidation of methane in the stratosphere (roughly at a ratio 50:50) and does not depend on the local temperature. Please clarify what you mean.

- Figures 6 and 7: please give “dust density” in number densities and not mass densities for easier interpretation in terms of known values from previous models and observations.

- Section 3.3: these are important results. The authors should maybe also state that measurements with SOFIE on AIM can only then be properly explained if the refractive index for crystalline ice is used, but not for amorphous ice. I remember that this was presented by Mark Hervig at several meetings. The authors might like to check back with him where this is published.

References

von Cossart, G., J. Fiedler, and U. von Zahn (1999), Size distributions of NLC particles as determined from 3-colour observations of NLC by ground-based lidar, *Geophys. Res. Lett.*, 26(11), 1513 – 1516, doi:10.1029/1997GL900226.

DeLand, M. T., and G. E. Thomas (2015), Updated PMC trends derived from SBUV data, *J. Geophys. Res. Atmos.*, 120, 2140–2166, doi:10.1002/2014JD022253.

Matthew T. DeLand, Eric P. Shettle, Gary E. Thomas, John J. Olivero, A quarter-century of satellite polar mesospheric cloud observations, *Journal of Atmospheric and Solar-Terrestrial Physics*, Volume 68, Issue 1, 2006, Pages 9-29, <https://doi.org/10.1016/j.jastp.2005.08.003>.

Donahue, T.M., Guenther, B., Blamont, J.E., 1972. Noctilucent clouds in daytime: circumpolar particulate layers near the summer mesopause. *Journal of the Atmospheric Sciences* 30, 515–517.

Duft, D., Nachbar, M., and Leisner, T.: Unravelling the microphysics of polar mesospheric cloud formation, *Atmos. Chem. Phys.*, 19, 2871–2879, <https://doi.org/10.5194/acp-19-2871-2019>, 2019.

Jesse, O., Auffallende Erscheinungen am Abendhimmel, *Meteorol. Z.*, 2, 311 –312, 1885.

Kirkwood, S., and K. Stebel, Influence of planetary waves on noctilucent cloud occurrence

over NW Europe, *J. Geophys. Res.*, 108(D8), 8440, doi:10.1029/2002JD002356, 2003.

Lübken, F.J., Lautenbach, J., Höffner, J., Rapp, M., Zecha, M., 2009. First continuous temperature measurements within polar mesosphere summer echoes. *Journal of Atmospheric and Solar-Terrestrial Physics* 71, 453–463.

Benjamin J. Murray, Eric J. Jensen, Homogeneous nucleation of amorphous solid water particles in the upper mesosphere, *Journal of Atmospheric and Solar-Terrestrial Physics*, Volume 72, Issue 1, 2010, Pages 51-61, <https://doi.org/10.1016/j.jastp.2009.10.007>.

Rapp, M., Thomas, G.E., 2006. Modeling the microphysics of mesospheric ice particles: assessment of current capabilities and basic sensitivities. *Journal of Atmospheric and Solar-Terrestrial Physics* 68, 715–744.

Thomas, G. E., Olivero, J. J., Jensen, E. J., Schröder, W., & Toon, O. B. (1989). Relation between increasing methane and the presence of ice clouds at the mesopause. *Nature*, 338, 490–492.

Thomas, G. E., and C. P. McKay (1985), On the mean particle size and water content of polar mesospheric clouds, *Planet. Space Sci.*, 33, 1209– 1224.

Thomas, G.E., Olivero, J.J., DeLand, M., Shettle, E.P., 2003. A response to the article by U. von Zahn. "Are noctilucent clouds truly a miner's canary of global change?". *EOS, Transactions—American Geophysical Union* 84 (36), 352–353.

Vestine, E., Noctilucent clouds, *Journal of the Royal Astronomical Society of Canada*, 28, 249-272, 1934

Witt, G., Height, structure and displacements of noctilucent clouds, *Tellus*, 14, 1 – 18, 1962.

Witt, G., The nature of noctilucent clouds, *Space Res.*, 9, 157– 169, 1969