

Atmos. Chem. Phys. Discuss., referee comment RC1  
<https://doi.org/10.5194/acp-2021-866-RC1>, 2022  
© Author(s) 2022. This work is distributed under  
the Creative Commons Attribution 4.0 License.

## Comment on acp-2021-866

Anonymous Referee #1

---

Referee comment on "Measurement report: Variations in surface SO<sub>2</sub> and NO<sub>x</sub> mixing ratios from 2004 to 2016 at a background site in the North China Plain" by Xueli Liu et al., Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2021-866-RC1>, 2022

---

### General Comments

This paper reports on NO<sub>x</sub> and SO<sub>2</sub> measurements at a background site in the North China Plain. The site, the instrumental setup, quality control and the data processing procedures have been described in detail. Data are compared to other data from other measurement sites. The long-term trend of both SO<sub>2</sub> and NO<sub>x</sub>, their diurnal and seasonal behavior are discussed and compared to emission data. As publications of long time series of NO<sub>x</sub> and SO<sub>2</sub> are rare, the manuscript should be published after these questions have been answered:

### Specific Comments

Line 66 and following

Here the authors describe the setup of the instrument. For NO<sub>x</sub> detection, a chemiluminescence analyzer has been used. Unfortunately, it is not stated how NO<sub>2</sub> is converted to NO for detection.

Has a thermal or a photolytical converter been used? If a thermal converter has been used this would mean that a large fraction of the NO<sub>x</sub> would be in fact NO<sub>y</sub>, as nitric acid and nitrates would cause significant interferences in the NO<sub>2</sub> channel (Jung et al., 2017, Steinbacher et al., 2007). This interference should be discussed.

How frequent was the conversion efficiency determined? Was an NO<sub>2</sub> gas standard used or was it done by gas phase titration?

How was mixing depth determined?

Line 162 and following

The authors state that there is a lag between SO<sub>2</sub> mixing ratios and emissions. What can cause such a lag? Change in meteorology would only lead to a decoupling of emissions and measured SO<sub>2</sub> and NO<sub>2</sub> values but not to a time shift.

Line 166 and following

The authors write that the 2008 Olympic games should affect the emissions. However, in figure 5 the measured SO<sub>2</sub> values are highest in 2008 and low in 2009. Is it possible the effect of the worldwide economic crisis in 2009 is larger than the effect of the Olympic games and not well represented in the emission data?

Line 246 and following

In chapter 4.3. the authors explain the different diurnal features of SO<sub>2</sub> and NO<sub>x</sub>. The profile of NO<sub>x</sub> coincides with the change in mixing depth while the diurnal profile of SO<sub>2</sub> is opposite to it. The profile of SO<sub>2</sub> is explained by transport from SO<sub>2</sub>-rich air from above which originate from remote, not necessarily elevated, sources, which will increase SO<sub>2</sub> concentration during daytime. However, during nighttime an efficient loss process must reduce the SO<sub>2</sub> concentration again. What is the loss process that reduces the concentration of SO<sub>2</sub> reduced during nighttime?

Line 246 and following

With respect to the daily profile of NO<sub>x</sub> it was argued that it is the result of transport processes during noontime. But isn't it more likely that NO<sub>x</sub> is emitted from local sources close to the ground, as motor vehicles and small burners This was observed at the background site at Linan (Yin et al., 2022), which showed similar diurnal cycles and similar mean NO<sub>x</sub> concentrations.

Jung, J., Lee, J., Kim, B., & Oh, S. (2017). Seasonal variations in the NO<sub>2</sub> artifact from chemiluminescence measurements with a molybdenum converter at a suburban site in Korea (downwind of the Asian continental outflow) during 2015–2016. *Atmospheric Environment*, 165, 290-300.  
<https://doi.org/https://doi.org/10.1016/j.atmosenv.2017.07.010>

Steinbacher, M., Zellweger, C., Schwarzenbach, B., Bugmann, S., Buchmann, B., Ordóñez, C., Prevot, A. S. H., & Hueglin, C. (2007). Nitrogen oxide measurements at rural sites in Switzerland: Bias of conventional measurement techniques [<https://doi.org/10.1029/2006JD007971>]. *Journal of Geophysical Research: Atmospheres*, 112(D11). <https://doi.org/https://doi.org/10.1029/2006JD007971>

Yin, Q., Ma, Q., Lin, W., Xu, X., & Yao, J. (2022). Long-term variations in surface NO<sub>x</sub> and SO<sub>2</sub> mixing ratios from 2006 to 2016 at a background site in the Yangtze River Delta region, China. *Atmos. Chem. Phys.*, 22(2), 1015-1033.  
<https://doi.org/10.5194/acp-22-1015-2022>

## Technical corrections

Line 91 High values in winter and low values in summer

Line 102 lower mixing depth heights

Line 112 6.27 ppb (ppb is missing)

Line 112 I think a one-digit precision of the values is sufficient here

Line 238 More difficulties

Line 247 high values in winter and low values

Line 249 greatly different from each other

Line 284 low values in

Line 285 exhibited great differences