

Atmos. Chem. Phys. Discuss., author comment AC2  
<https://doi.org/10.5194/acp-2021-957-AC2>, 2022  
© Author(s) 2022. This work is distributed under  
the Creative Commons Attribution 4.0 License.

## Reply on RC1

Baoshuang Liu et al.

---

Author comment on "Dramatic changes in atmospheric pollution source contributions for a coastal megacity in northern China from 2011 to 2020" by Baoshuang Liu et al., Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2021-957-AC2>, 2022

---

The manuscript applied a machine learning-based meteorological normalization approach to decouple the meteorological effects from air quality trend in a coastal city in northern China, and further assessed the changes in the contributions of pollution sources in Qingdao in the past ten years. There are some minor issues need to be solved.

- In the last paragraph of the introduction, if any, it is suggested to add some evaluation articles on the changes in air quality and the effectiveness of control measures before and after major events held in Qingdao (e.g., the 2018 the Shanghai Cooperation Organization summit in Qingdao). Clarify the differences with this paper to support the particularity of this study.

**Response:** Thanks for your advice. We added the related contents in the revised version (on the lines 102-106). The more details as following:

Liu et al. (2020a) assessed the changes in O<sub>3</sub> concentrations during the Shanghai Cooperation Organization (SCO) Summit in Qingdao and analyzed the impact of control measures on the emissions reduction of its precursors, and Liu et al. (2020b) also analyzed the reasons for the increase of O<sub>3</sub> concentration at nighttime during the SCO Summit.

- In the sampling and analysis section (L144-145), what is the basis for collecting 22 hours a day? Why not 24 hours?

**Response:** Thanks for your advice. The sampling period was from 11:00 to 09:00 the next day, and the sampling time was 22 hours. The sampling time covers the peak time of morning and evening traffic and the period of strong photochemical reaction at noon, and includes the peaks and troughs of air pollutant concentration. The sampling period of this study is representative. Meanwhile, because the sampling instruments in this study cannot continuously replace the filters, therefore, it needs to take a certain time to replace the sampling filters for subsequent sampling. Furthermore, similar sampling times (22-23 hours) occurred in the studies of Liu et al., (2021), Fang et al. (2017), Turap et al. (2019), and Wang et al. (2016).

Liu, B.S., Wu, J.H., Wang, J., Shi, L.Y., Meng, H., Dai, Q.L., Wang, J., Song, C.B., Zhang, Y.F., Feng, Y.C., Hopke, P.K., 2021. Chemical characteristics and sources of ambient PM<sub>2.5</sub> in a harbor area: Quantification of health risks to workers from source-specific selected toxic elements. *Environ. Pollut.*, 2021, 268, 115926.

Fang, X.Z., Bi, X.H., Xu, H., Wu, J.H., Zhang, Y.F., Feng, Y.C., 2017. Source apportionment of ambient PM<sub>10</sub> and PM<sub>2.5</sub> in Haikou, China. *Atmos. Res.* 190, 1–9.

Turap, Y., Talifu, D., Wang, X.M., Abulizi, A., Maihemuti, M., Tursun, Y., Ding, X., Aierken, T., Rekefu, S., 2019. Temporal distribution and source apportionment of PM<sub>2.5</sub> chemical composition in Xinjiang, NW-China. *Atmos. Res.* 218, 257–268.

Wang, Y.N., Jia, C.H., Tao, J., Zhang, L.M., Liang, X.X., Ma, J.M., Gao, H., Huang, T., Zhang, K., 2016. Chemical characterization and source apportionment of PM<sub>2.5</sub> in a semi-arid and petrochemical-industrialized city, Northwest China. *Sci. Total Environ.* 573, 1031–1040.

- Have the sample data of 2011-2012, 2016, and 2019 been carried out for source analysis, respectively? Please clarify. In addition, the data of all sites were chronologically ordered end to end for each PMF analysis? PMF is usually used for source analysis based on the long-time data of one site. Please explain the rationality.

**Response:** (1) Yes. The related contents were described in the lines 477-479. (2) Yes. The data of all sites were chronologically ordered end to end for each PMF analysis, and the related contents have shown in text S2 in the supplementary material. (3) In this study, the data of all sampling sites were included in PMF for source analysis, mainly because all sampling sites were located in Qingdao urban area, and the impact source-categories at these sites were the same. The stability of the model running can be improved due to the increase of the number of samples by incorporating multiple sites data into PMF. Escrig et al. (2009) suggested that because the three sampling sites were affected by the same source-categories, therefore, they can be treated as one site and then included in PMF for analysis. Mooibroek et al. (2011) incorporated the receptor data of five sampling sites into PMF for source apportionment, which considered that combining the receptor data of five sites can increase the amount of input data for the model, and then more stable analysis results can be obtained. Furthermore, many related studies have adopted similar methods (Lu et al., 2018; Ezeh et al., 2019; Zhao et al., 2019).

Escrig, A., Monfort, E., Celades, I., Querol, X., Amato, F., Minguillón, M.C., and Hopke, P.K.: Application of optimally scaled target factor analysis for assessing source contribution of ambient PM<sub>10</sub>, *J. Air & Waste Manage.*, 59, 1296–1307, <https://doi.org/10.3155/1047-3289.59.11.1296>, 2009.

Mooibroek, D., Schaap, M., Weijers, E.P., and Hoogerbrugge, R.: Source apportionment and spatial variability of PM<sub>2.5</sub> using measurements at five sites in the Netherlands, *Atmos. Environ.*, 45, 4180-4191, <https://doi.org/10.1016/j.atmosenv.2011.05.017>, 2011.

Lu, Z.J., Liu, Q. Y., Xiong, Y., Huang, F., Zhou, J. B., and Schauer, J.J.: A hybrid source apportionment strategy using positive matrix factorization (PMF)

and molecular marker chemical mass balance (MM-CMB) models, *Environ Pollut.*, 238, <https://doi.org/10.1016/j.envpol.2018.02.091>, 2018.

Ezeh, G, C., Obioh, I, B., Asubiojo, O., Abiye, O, E., and Onyeuwaoma, N, D.: A study of PM<sub>2.5-10</sub> pollution at three functional receptor sites in a sub-Saharan African megacity, *Aerosol Science and Engineering.*, 3, <https://doi.org/10.1007/s41810-019-00044-3>, 2019.

Zhao, Z.P., Lv, S., Zhang, Y.H., Zhao, Q.B., Shen, L., Xu, S., Yu, J.Q., Hou, J.W., and Jin, C.Y.: Characteristics and source apportionment of PM<sub>2.5</sub> in Jiaxing, China, *Environ. Sci. Pollut. R.*, 26, <https://doi.org/10.1007/s11356-019-04205-2>, 2019.

- In the process of PMF calculation, the author analyzed DISP and BS (Fig. S9-11, Table S14), but the author did not mention it in the introduction of PMF method. Please add.

**Response:** Thanks for your advice. The related contents have been added in the revised manuscript (on the lines 238-239). The more details as following:

Bootstrap (BS) and displacement (DISP) analyses were used to investigate the effects of measurement error and rotation ambiguity on the resulting solutions.

- In recent years, scholars have studied the change of air quality in many cities around the world (such as Beijing), many of which use random forest and other methods. In section 3.11, it is suggested to add the content of comparative analysis with other cities.

**Response:** Thanks for your advice. We added the contents of comparative analysis with other cities in the revised manuscript (on the lines 288-299). The more details as following:

Similar to this study, Vu et al. (2019) found that primary emission controls required by the CAAP in Beijing have led to substantial reductions in PM<sub>2.5</sub>, PM<sub>10</sub>, NO<sub>2</sub>, SO<sub>2</sub>, and CO from 2013 to 2017 of approximately 34%, 24%, 17%, 68%, and 33%, respectively, after meteorological normalization. Zhai et al. (2019) suggested that the mean PM<sub>2.5</sub> decreased across China was 4.6  $\mu\text{g m}^{-3} \text{ yr}^{-1}$  in the meteorology-corrected data from 2013 to 2018, and the Beijing–Tianjin–Hebei, the Yangtze River Delta, the Pearl River Delta, the Sichuan Basin, and the Fenwei Plain decreased 8.0, 6.3, 2.2, 4.9, and 5.0  $\mu\text{g m}^{-3} \text{ yr}^{-1}$ , respectively. Overall, the concentrations of most air pollutants (i.e., PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub>, NO<sub>2</sub>, and CO) in China have showed a decreasing trend in recent years (Zhao et al., 2021; Fan et al., 2020, while that of O<sub>3</sub> has performed an increasing trend (Li et al., 2020; Ma et al., 2021), which further facilitated secondary particles formation (Wang et al., 2016; Nøjgaard et al., 2012).

- L401-404 "The observed and normalized concentrations of PM<sub>2.5</sub> during the whole study period were 93 and 83  $\mu\text{g m}^{-3}$ , suggesting that unfavorable meteorological conditions

generated approximately  $10 \mu\text{g m}^{-3}$  of growth of  $\text{PM}_{2.5}$ ", can the difference of simple subtraction represent the influence value of meteorology? Is there any basis?

**Response:** Thanks for your advice. In this study, we used the ventilation coefficient to normalize the impact on  $\text{PM}_{2.5}$  concentrations from the meteorology, and the observed and normalized concentrations of  $\text{PM}_{2.5}$  during the whole study period were  $93$  and  $83 \mu\text{g m}^{-3}$ , respectively, suggesting that the meteorological conditions might explain approximately  $10 \mu\text{g m}^{-3}$  of  $\text{PM}_{2.5}$  variation. The inference here was mainly based on the studies from Ding et al. (2021) and Zhai et al. (2019). Ding et al. (2021) found that the mean measured  $\text{PM}_{2.5}$  mass concentration and dispersion coefficient normalized  $\text{PM}_{2.5}$  mass concentration were  $67.5$  and  $45.6 \mu\text{g m}^{-3}$  during the COVID-19 lockdown period, indicating that unfavorable meteorological condition generated approximately  $22 \mu\text{g m}^{-3}$  growth of  $\text{PM}_{2.5}$ . Zhai et al. (2019) found that the mean  $\text{PM}_{2.5}$  decrease across China was  $4.6 \mu\text{g m}^{-3} \text{yr}^{-1}$  in the meteorology-corrected data from 2013 to 2018, 12% lower than in the original data, meaning that 12% of the  $\text{PM}_{2.5}$  decrease in the original data was attributable to the meteorology. Furthermore, we also added the related contents from other studies in the revised manuscript (on the lines 422-426).

Ding, J., Dai, Q. L., Li, Y. F., Han, S. Q., Zhang, Y. F., and Feng, Y. C.: Impact of meteorological condition changes on air quality and particulate chemical composition during the COVID-19 lockdown, *J. Environ. Sci.*, 109, 45-56, <https://doi.org/10.1016/j.jes.2021.02.022>, 2021.

Zhai, S., Jacob, D.J., Wang, X., Shen, L., Li, K., Zhang, Y., 2019. Fine particulate matter ( $\text{PM}_{2.5}$ ) trends in China, 2013–2018: separating contributions from anthropogenic emissions and meteorology. *Atmos. Chem. Phys.* 19, 11031-11041.

- L542-581, this paper analyzed the changes in source contributions in the winter heating periods in Qingdao from 2011-2012, 2016, and 2019. The author should clarify the reasons and necessity of this analysis.

**Response:** Thanks for your advice. The related reasons and necessity have been added in the revised manuscript (on the lines 563-569). The more details as following:

Furthermore, with the beginning of heating season in northern cities in China (Liu et al., 2016; Li et al., 2019a), the atmospheric pollutant emissions increased substantially (Chen et al., 2020a). Coupled with the adverse meteorological conditions (Li et al., 2019a), haze episodes occurred frequently during this period (Liu et al., 2018a; Yang et al., 2020). Therefore, the control effects of pollution sources and key control sources in the specific period can be better highlighted through analyzing the changes in the contributions of emission sources during heating seasons over the years.

Chen, J.Y., Shan, M., Xia, J.J., and Jiang, Y.: Effects of space heating on the pollutant emission intensities in "2+26" cities, *Building Environ.*, 175, 106817, <https://doi.org/10.1016/j.buildenv.2020.106817>, 2020a.

Li, H., You, S.J., Zhang, H., Zheng, W.D., and Zou, L.J.: Analysis of the impacts of heating emissions on the environment and human health in North China, *J. Clean Prod.*, 207,

728-742, <https://doi.org/10.1016/j.jclepro.2018.10.013>, 2019a.

Liu, B.S., Song, N., Dai, Q.L., Mei, R.b., Sui, B.H., Bi, X.H., and Feng, Y.C.: Chemical composition and source apportionment of ambient PM<sub>2.5</sub> during the non-heating period in Taian, China, *Atmos. Res.*, 170, 23-33, <https://doi.org/10.1016/j.atmosres.2015.11.002>, 2016.

Liu, B.S., Cheng, Y., Zhou, M., Liang, D.N., Dai, Q.L., Wang, L., Jin, W., Zhang, L.Z., Ren, Y.B., Zhou, J.B., Dai, C.L., Xu, J., Wang, J., Feng, Y.C., and Zhang, Y.F.: Effectiveness evaluation of temporary emission control action in 2016 in winter in Shijiazhuang, China, *Atmos. Chem. Phys.*, 18, 7019-7039, <https://doi.org/10.5194/acp-18-7019-2018>, 2018a.

Yang, S., Duan, F., Ma, Y., Li, H., Ma, T., Zhu, L., Huang, T., Kimoto, T., and He, K.: Mixed and intensive haze pollution during the transition period between autumn and winter in Beijing, China, *Sci. Total Environ.*, 711, 134745, <https://doi.org/10.1016/j.scitotenv.2019.134745>, 2020.