

# ***Interactive comment on “Comparison of the GRUAN data products for Meisei RS-11G and Vaisala RS92-SGP radiosondes at Tateno (36.06° N, 140.13° E), Japan” by E. Kobayashi et al.***

**Fasso**

alessandro.fasso@unibg.it

Received and published: 3 February 2019

Regarding comments of RC2, please note that in my comment SC1, I suggest that "the histogram of differences (d) in each layer ... may help".

In fact,  $P(|d| > 2 \cdot \sigma) = 0.95$  is true if the differences are zero-mean Gaussian coherently with authors' lines 32-33: "Assume that  $m_1 = m_2$  is true and that uncertainty follows normal distribution.". If the differences are zero mean, but non-Gaussian, for example, zero mean Student's t with 3 degrees of freedom, then the constant 2 is wrong and the corresponding Student's t percentile should be used.

Printer-friendly version

Discussion paper



So if, after subtracting the mean of  $d$ , the distribution of  $d$  at a certain pressure level is not approximately Gaussian, then the constant 2 is not the appropriate one.

Which would be the correct constant depends on the distribution of  $d$ . So my general consideration does not give the final answer but may help in understanding a step more on this issue.

---

Interactive comment on Atmos. Meas. Tech. Discuss., doi:10.5194/amt-2018-416, 2019.

Printer-friendly version

Discussion paper

