

Atmos. Meas. Tech. Discuss., author comment AC3
<https://doi.org/10.5194/amt-2021-113-AC3>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Reply on RC1

Longjiang Li et al.

Author comment on "A new zenith hydrostatic delay model for real-time retrievals of GNSS-PWV" by Longjiang Li et al., Atmos. Meas. Tech. Discuss.,
<https://doi.org/10.5194/amt-2021-113-AC3>, 2021

My response:

Thanks for the reviewer's suggestions and we have made changes for all the suggestions in the new manuscript accordingly. The detailed modifications are as follows.

1) The motivation of the study includes:

- a) Improving the blind models used in real-time retrieval of GNSS-PWV under the condition that meteorological sensor are not equipped at the GNSS stations.
- b) Providing an alternative method in consideration of the inconvenience in acquiring forecast pressure from numerical weather models. The forecast data need to be downloaded in advance, which increases the complexity of data processing, not to mention the fact that the forecast data may not be obtained due to various reasons, e.g. problems of some servers or agencies.
- c) Coming up with a new method for the determination of the ZHD in the retrieval of GNSS-PWV.

2) Line 54 has been corrected.

3) There are no topography and interpolation used in Figure 8. When ZHD-ERA5 was used as the reference, the ZHD-ERA5 at a grid point was calculated by the integral from the lowest level (1000 hPa) to the highest level (1 hPa), and the ZHD-GPT3 at the same grid point was calculated using the Saastamonien model with the input pressure derived from GPT3 at the lowest level. Note that the geopotential height in ERA5 was converted to ellipsoidal height using the method described in the following paper:

Wang, X., Zhang, K., Wu, S., Fan, S., and Cheng, Y.: Water vapor-weighted mean temperature and its impact on the determination of precipitable water vapor and its linear trend: Water Vapor-Weighted Mean Temperature, J. Geophys. Res. Atmos., 121, 833–852, <https://doi.org/10.1002/2015JD024181>, 2016.

The reason for the large biases in Figure 8 may be that the GPT3 model performed poorly in the regions where the height is significantly different from the earth surface. The height of the grid points from ERA5 was above the earth surface in most regions but below the earth surface in mountainous regions, since the pressure at the lowest level of the grid points from ERA5 is almost constant (1000 hPa) but the heights at the lowest level in

different grid points are different.

4) Figure 3 caption has been corrected, as advised.

5) Lines 110, 277 and 305 have been corrected, as advised.

6) In addition to the above suggestions, we also found a new problem: when the integral method was used to calculate the ZHD, geoidal height or ellipsoidal height should be used, so the geopotential height contained in ERA5 and sounding data at each pressure level needs to be converted to geoidal or ellipsoidal height. However, in the previous manuscript, the height was not converted. Although this fault might cause large biases in the ZHD, it is not necessarily to largely affect the RMSE of the ZHD. This is due to that the effect of this fault changes biases from positive to negative, but the values are about equal. In the new manuscript, all the problems have been corrected.

Please also note the supplement to this comment:

<https://amt.copernicus.org/preprints/amt-2021-113/amt-2021-113-AC3-supplement.pdf>