

Atmos. Meas. Tech. Discuss., referee comment RC2  
<https://doi.org/10.5194/amt-2021-178-RC2>, 2021  
© Author(s) 2021. This work is distributed under  
the Creative Commons Attribution 4.0 License.



## Comment on amt-2021-178

Anonymous Referee #1

---

Referee comment on "Retrieval algorithm for OCIO from TROPOMI by Differential Optical Absorption Spectroscopy" by Jānis Puķīte et al., Atmos. Meas. Tech. Discuss.,  
<https://doi.org/10.5194/amt-2021-178-RC2>, 2021

---

### General Comments:

In this work, a new retrieval algorithm of the slant column densities (SCDs) of OCIO is proposed. This algorithm, aimed to be applied to TROPOMI DOAS measurements, takes into account different spectral effects not considered in previous retrievals. A corresponding error analysis of some retrieval settings has been performed. The authors also present a comparison between OCIO SCDs, obtained by TROPOMI through the new algorithm, and from ground-based zenith DOAS measurements at Kiruna and Neumayer stations. The results show a very good agreement with these instruments (especially at Kiruna). The SCDs of OCIO obtained in this work have been also compared to preliminary S5p+I OCIO products during different periods of the year, showing similar SCD evolution but presenting an offset between both datasets.

All the manuscript (text and figures) is clearly presented. The new concepts and settings introduced in this new algorithm are exhaustively explained (appendixes), as well as the corresponding error analysis and sensitivity studies. I think that the results exposed in this work will be useful for the treatment and analysis of the OCIO SCDs obtained by TROPOMI. Thus, I think the paper should be published in AMT. However, I think that some questions should be better clarified.

My main concern is that the authors of this work claim that the new algorithm improves the retrieval results, but looking at the comparison between SP5+I and the results of this work, can that be really stated? The new algorithm takes into account several fine effects that, in principle, should improve the OCIO SCDs retrieval and decrease the corresponding errors. But, in practice, how can we say that the results of these works are better than those of SP5+I? For high SCDs, results are very similar, and for low SCDs the offset between both datasets cannot be explained. It is true that, as the authors explained, OCIO observation is not expected when the temperatures are still warm, as it is observed in the results of this study (Figure 10). Contrarily, SP5+I results show a background level of OCIO. But, it can be affirmed that the results of this work are better than those provided by the SP5+I? Did the authors compare the results of both algorithms with independent measurements (as those of Kiruna or Neumayer)? Please, explain better.

