

Atmos. Meas. Tech. Discuss., author comment AC2
<https://doi.org/10.5194/amt-2022-133-AC2>, 2022
© Author(s) 2022. This work is distributed under
the Creative Commons Attribution 4.0 License.

Reply on RC2

Kimberlee Dubé et al.

Author comment on "An improved OSIRIS NO₂ profile retrieval in the upper troposphere–lower stratosphere and intercomparison with ACE-FTS and SAGE III/ISS" by Kimberlee Dubé et al., Atmos. Meas. Tech. Discuss.,
<https://doi.org/10.5194/amt-2022-133-AC2>, 2022

Thank you for taking the time to review our manuscript and for providing helpful suggestions. Our responses to each comment are included here in bold text.

This is a solid paper on a topic that fits in AMT. It describes a new OSIRIS NO₂ data product (version 7.2) that improves upon previous versions, addressing deficiencies in the retrieval methodology such as how the reference radiance is calculated and changes in spectral resolution with instrument temperature. Comparisons are made with co-incident observations from occultation instruments. It is well written and the subject is worthy of publication in AMT.

My only criticism of substance is that considering "... version 7.2, was designed to fix a low bias and to improve performance in the UTLS ..." it really does not do an adequate job evaluating the performance of the new product in this region. This paper should attempt to better answer questions such as "precisely how good is v7.2 in the UTLS" or "is v7.2 actually better than the old version", and if this cannot be answered, then why not. This is the region where diurnal effect errors can be quite substantial. Of the three datasets used, it was ignored in OSIRIS and ACE-FTS, and corrected for in ISS/SAGEIII, which muddies the waters. For example, the OSIRIS-ACE and OSIRIS-SAGE results seem to conflict. Does that mean there are biases between ACE and SAGEIII? Is it related to the diurnal effect? I made some suggestions below as to how to investigate the diurnal effect as, especially for OSIRIS, it is not that easy. That said, the lead author is well equipped to examine this further. It is also difficult to know what to make of the comparison that go all the way down an altitude of 10 km. Is there sufficient data for these to be meaningful? And does the variability swamp the signal? All-in-all this part of the paper needs additional analysis such that they would support some clear summary findings.

Once this, along with several smaller clarifications outlined below, is completed I would support publication.

Thank you for the thoughtful response. We have performed further analysis to determine the effect of diurnal variations on the intercomparisons presented in our paper. Several paragraphs discussing these results, along with a discussion of why we think v7.2 is an improvement over v6, have been added to the conclusion of our manuscript:

NO₂ from the SAGE III/ISS DV retrieval shows improved agreement with OSIRIS compared to NO₂ from standard SAGE retrieval. The diurnal effect produces a high bias in NO₂ retrieved from occultation instruments below 25 km. There is no version of the ACE-FTS NO₂ retrieval that accounts for diurnal variations along the line of sight, which could be increasing the difference with OSIRIS below about 25 km. It is not expected that the diurnal effect would be greater in ACE-FTS than in SAGE III/ISS, so this will add at most a 5% to 40% bias, with the largest bias at the lowest altitudes (Dube et al 2021). If we assume an approximate high bias of 25% below 20 km in the ACE-FTS NO₂, the difference between ACE-FTS and OSIRIS will become greater near the tropopause, and especially in the NH. The bias in this region will be >50%, as it is for SAGE III/ISS and OSIRIS. In the SH the bias between ACE-FTS and OSIRIS is ~ -15% below 20 km. In this region including the diurnal correction will result in a bias of ~10%, corresponding to improved agreement between OSIRIS and ACE-FTS. This is opposite in sign and smaller in absolute value than the bias between SAGE III/ISS and OSIRIS in this region. There are significantly more coincidences between ACE-FTS and OSIRIS than there are between SAGE III/ISS and OSIRIS at SH mid-high latitudes, which could explain why we see a smaller bias between ACE-FTS and OSIRIS.

We also considered the effect of diurnal variations along the OSIRIS LOS by repeating the analysis using only OSIRIS scans with a SZA <85, corresponding to scans shown by McLinden et al (2006) to have a significant bias. Removing these scans had a minimal effect on the comparisons between OSIRIS and ACE-FTS in the stratosphere, changing the bias by at most 3%. It had a greater effect on the comparison with SAGE III/ISS, where the agreement improved by up to 10% in some bins in the lower stratosphere.

Overall, we conclude based on comparison with NO from ACE-FTS and SAGE III/ISS that the OSIRIS v7.2 NO₂ product is an improvement over the previous v6.0 OSIRIS NO₂, particularly below 30 km. The bias between OSIRIS v7.2 and

both ACE-FTS and SAGE III/ISS has decreased from v6.0, except for in a few bins near the tropopause where including the diurnal effect in SAGE III/ISS (or estimating the magnitude of the effect for ACE-FTS) makes the bias greater than it was in v6.0

Line 20: you spelled out the acronyms in the abstract, but do so when first used in the main text as well.

The AMT style guide requires defining acronyms in both the abstract and the main text (<https://www.atmospheric-measurement-techniques.net/submission.html#english>)

Line 27-28: use "... PRATMO stratospheric photochemical ..." and "... McLinden et al. (2000) and later Adams et al. (2017)." (<https://doi.org/10.5194/acp-17-8063-2017>)

This reference has been added

Figure 2: what time period is this for?

It is from November 2001 to May 2020. This is now mentioned in the figure caption.

Line 95: "worse agreement between the OSIRIS measurements and the forward model"
Worse how? Shouldn't this be one of the parameters one can optimize by simulating OSIRIS radiances accounting for precision and sampling/field of view considerations. The 2-3 km over most of the stratosphere argument would seem to apply for any of the values of alpha used. Would a smaller alpha help in the UTLS?

We have added ", particularly above 30 km." to be clear what this specific statement was referring to.

Generally ideal retrieval theory would say that reducing alpha increases sensitivity, at the cost of poorer precision. This is something that could be simulated as the reviewer suggests. However we have noticed that the retrieval tends to behave poorly at low altitudes for smaller values of alpha, and in particular fails to converge often. This is likely due to an effect where small biases in the forward model (or assumptions made) are amplified as the amount of regularization decreases. The biases are things that are challenging to simulate such as patchy clouds, or even identify in some cases. Essentially what we try to do is pick the lowest value of alpha where the retrieval converges most of the time and still behaves reasonably.

Line 110: Be more specific about where v6 and v7.2 begin to differ. E.g. do they both solve for the minimum in equation

A table has been added which summarizes the main features of each retrieval version.

Line 125: "The log-normal distributions are less physically realistic" ... why is this? This could be true but maybe it is not that obvious. Please give some rationale for this. If this is in the upper troposphere one might expect something non-Gaussian for if there is occasional lightning-NO_x.

This section was a little unclear. At the lowest altitudes these distributions are

actually dominated by the precision of the measurements rather than geophysical NO2 variations which is why we expect them to be normally distributed. We have added a statement to that effect.

Line 139 or thereabouts: what is the DOFS, or range in DOFS, for this data product?

The value depends on the profile, but it is typically around 13. However, we prefer to use vertical resolution as the diagnostic because the DOFS depends on the lowerbound altitude.

Line 144: please motivate the use of 1.5 km as the threshold a little more.

The larger the value the less data is removed. We tested several values for the threshold and 1.5 km was a good compromise between removing what is likely good data and not removing what is likely bad data. A larger difference means we are including information that is quite far from the tangent point. This is now mentioned in the manuscript.

Line 155: "... kernel filter, as used in version 6.2"?

The kernel filter is applied to the version 7.2 data. This is now mentioned in the manuscript.

Line 170: "change in the processor"... what does this mean? Same for "global environment settings" later on.

This means that the processing was moved to a different computer.

Line 181: would the lower altitude of the ISS, relative to SCISAT, mean there is ~1 more orbit per day... I think there are generally 16 orbits per day for the ISS

You are correct, this has been fixed.

Line 192: does OSIRIS not need to worry about O4 absorption in their NO2 retrieval?

The inclusion of O2-O2 absorption typically makes very little difference in the NO2 spectral region since it is largely uncorrelated with the NO2 absorption. For SAGE-III it likely makes a larger difference for the O3 retrieval.

Line 203: So this is an off-line (unofficial) retrieval, correct? This should be clarified. Related: later on you use "SAGE III/ISS DV v5.2"... what does this mean? It implies, I think, that the product is v5.2 but with a diurnal correction. If you did your own retrieval then I would not use the 'v5.2' label which implies to me that it is an official product.

The correction is applied to the official v5.2 data product. This is what we mean to suggest by the label DV v5.2. We have clarified this in the manuscript.

Line 221-224: this is good information to have in section 3.2.1

This information has been moved to section 3.2.1

Line 225: "The effect of changes in the input parameters on the PRATMO NO₂ was estimated by perturbing them in the model." What was this used for?

This information was not used directly, it is just included to provide an idea of how sensitive the modelled NO₂ is to the input parameters.

Line 228-229: "... it is not always enough to account for the difference between sunrise and sunset occultations from a single instrument." Is the purpose of this statement to point out that systematic errors may still remain suggesting that the scaling is not perfect? My guess is that there could be sampling issues that would also cause the SR and SS to not be the same anyway.

Yes, we are just suggesting that the scaling is not perfect. You are correct, the sampling differences likely also contribute.

Figure 7: "... coincident NO₂, when shifted from their individual local times to 12:00 pm, for several altitudes..." is this what you mean? If not, please clarify.

Yes, this is what we mean. The wording has been adjusted.

Line 245: Shouldn't v7 have more data lower down based on Figure 5?

Figure 5 is comparing the amount of v7.2 data retrieved before and after the application of the averaging kernel filter. The combination of this filter, plus the

improved cloud filtering in v7.2, means that we are removing more (unrealistic) data from v7.2 than from v6 at the low altitudes.

Line 245: showing the standard deviations of the average differences, or the average \pm s.d., would suggest if this is due to fewer coincidences or not.

It does not really make sense to compare the standard deviations at these lower altitudes because the data is normally distributed in v7.2, but has closer to a lognormal distribution in v6.0.

Section 4.1: any discussion of the differences between ACE and OSIRIS below 20 km needs to factor in diurnal effect errors. Neither ACE nor OSIRIS corrects for them, but you should be able to deduce the impact of these based on your work on ISS/SAGE III and other publications on the subject. Previous comparisons, e.g., see Brohede et al. (doi:10.1029/2006JD007586) Figure 9, show the expected high bias in ACE-FTS. The effect in OSIRIS is harder to generalize due to the varying geometry, but was estimated in McLinden et al. (doi:10.1029/2005JD006628) Figures 4 and 5. You might consider limiting OSIRIS to $SZA < 85$ OR [$SZA < 85$ or ($SZA < 88$ and $75 < dAZ < 105$)]

Thank you for the suggestion. We have further investigated the diurnal effect in ACE and OSIRIS NO₂. Requiring the OSIRIS scans to have $SZA < 85$ reduced the number of coincidences with ACE by $\sim 14\%$. Similarly, there are $\sim 12\%$ fewer coincidences between SAGE III/ISS and OSIRIS. Most of the lost coincidences are in the SH. The change in the number of coincidences in each latitude bin is shown in Figure 1 of the attached file. It is worth noting that we only retrieve data for scans with $SZA < 88$ to begin with.

We redid the comparisons after removing these coincidences and found that the effect is generally negligible. Removing these coincidences changes the bias between OSIRIS and ACE in the stratosphere by at most $\sim +/ - 3\%$. For SAGE III/ISS the effect is larger in a few bins, particularly near 50 S for the SS occultations (which is the bin that loses the most profiles after filtering).

Figure 2 in the attached file compares the difference between the mean bias in each bin with and without including scans where OSIRIS has an SZA>85. The top row is for sunrise occultations, and the bottom row is at sunset. A negative value shows that the bias with OSIRIS was reduced after removing scans with a significant diurnal effect.

Based on this analysis we conclude that the diurnal effect in OSIRIS data has a negligible effect on the comparison with ACE-FTS in the stratosphere, and a significant effect on the comparison with SAGE, but only in a few bins.

The diurnal effect will have a greater impact on ACE NO₂ than it does on OSIRIS NO₂. From Dube et al 2021 and Brohede et al 2007 we expect there to be a high bias of greater than 10% in the ACE NO₂ below ~30 km. Accounting for this effect would improve the agreement with OSIRIS in the SH, but decrease agreement in the NH where ACE is already low compared to OSIRIS.

A section discussing these findings has been added to the manuscript (section 4.3).

Section 4.2: What is the impact of the diurnal correction on the SAGEIII profiles? It is a little hard to tease in the figure. What do you make of the OS-S3 comparisons vs the OS-ACE below 20 km?

The impact of the diurnal correction on SAGE III is described in detail in Dube et al 2021(<https://amt.copernicus.org/articles/14/557/2021/>). Including this correction reduces the SAGE III NO₂ by about 10% at 25 km and more than 10% at lower altitudes (see figure 7 of Dube et al 2021). This is now mentioned in the manuscript.

Below 20 km we expect the ACE NO₂ to be biased high in the same way that the SAGE III NO₂ was before being corrected for diurnal effects. If we assume an approximate high bias of 25% in the ACE NO₂, the difference between ACE and

OSIRIS will become greater near the tropopause, and especially in the NH. The bias in this region will be >50%, as it is for SAGE III and OSIRIS. In the SH the bias between ACE and OSIRIS is ~-15% below 20 km. In this region including the diurnal correction will result in a bias of ~10%, corresponding to improved agreement between OSIRIS and ACE. This is opposite in sign and smaller in absolute value than the bias between SAGE and OSIRIS in this region. There are significantly more coincidences between ACE and OSIRIS than there are between SAGE and OSIRIS at SH mid-high latitudes, which could explain why we see a smaller bias between ACE and OSIRIS. We have added this discussion to the conclusion of the manuscript.

Line 274: "...from a data set..." what does this mean?

We intended to say "The relative anomaly is calculated by subtracting the overall mean value for a given month from all values for that month to remove the seasonal cycle (e.g. the mean June NO₂ concentration is subtracted from each individual June NO₂ concentration)". This has been corrected in the manuscript.

Line 284-285: "At sunrise ACE-FTS samples a region of the atmosphere that has not been illuminated long enough for the NO₂ to reach equilibrium, however this is not an issue at sunset." Are you agreeing this is true? I am not sure it is. Or are you merely restated the argument from Sheese et al.? Neither would be in a pseudo-steady-state at the terminator.

We are restating the theory of Sheese et al (2016)

Line 291: was the trend removed in these timeseries?

No, the trend was not removed. The only thing we removed was the seasonal cycle.

Line 316: "ACE-FTS retrieval it would likely improve agreement with OSIRIS" ... would it not make ACE-FTS values larger, thereby increasing the bias? I agree the diurnal effect will be similar for ACE and SAGEIII, and thus, either way you could be more quantitative here.

The ACE values would become smaller if the diurnal effect was included. We have elaborated on the discussion of the diurnal effect in the conclusion of the manuscript.

Please also note the supplement to this comment:

<https://amt.copernicus.org/preprints/amt-2022-133/amt-2022-133-AC2-supplement.pdf>