

Atmos. Meas. Tech. Discuss., author comment AC2 https://doi.org/10.5194/amt-2022-261-AC2, 2022 © Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.

Reply on RC2

Javier Gorroño et al.

Author comment on "Understanding the potential of Sentinel-2 for monitoring methane point emissions" by Javier Gorroño et al., Atmos. Meas. Tech. Discuss., https://doi.org/10.5194/amt-2022-261-AC2, 2022

Dear reviewer,

We have carefully considered your comments in the reviewed version of the manuscript. These comments have been very helpful to improve the overall quality of the manuscript. Here below we individually respond to each one of the comments:

 While the conclusions are indubitably important, one can regret the low amount of testing. Indeed, only 5 plume shapes and three locations (at a specific date) were used. I think it is too little to derive an accurate conclusion, especially given the variance of the results.

This is a very important comment and we thank the reviewer for bringing it up. In this reviewed version we have added a new subsection named "Extending the validation to a large plume dataset and season changes". This section includes 221 methane plumes and the winter vs. solstice acquisitions. These three sites are representative of a typical scenario for S2 methane detection. However, we agree that more sites could be studied. As the reviewer points out, follow-up work is expected which focuses on a large number of sites and conditions with an already consolidated methodology.

How stable is the simulation process regarding to the spectral characterization? The ESA regularly updates the spectral characterization of both S2A and S2B so I assume that each time the analysis needs to done again but one can wonder how much would be the error is this wasn't done (and as such what is the error that is done when using an imprecise spectral characterization).

We use version 3.0 of the spectral response. The spectral characterization changes mostly affect B1, B2 and B8. For methane detection and quantification, we select B12 and B11. The changes are not significant in these bands. Issue #39 in ESA: Sentinel-2 MSI Level-1C data quality report.

The authors mentioned the spectral sensitivity difference between S2A and S2B. It is not entirely clear the satellites is used for the different experiences (it seems that the observation is from S2A and the reference from S2B based on 1.106-112, and thus the synthetic experiment is using purely S2B). I think it could be interesting to study in that case the difference in detection limit between the two satellites (I assume that S2A limit is smaller than S2B since it is more sensitive to methane) and if there is a gain in

using the reference image from the same satellite as well.

We have studied a couple of examples over Hassi Messaoud which suggest a small increase in the detection limit (somewhere between 0-500kg/h). It is an interesting point that we just mentioned at the end of the manuscript. However, we believe that this is an issue to incorporate with more detailed analysis in a follow-up study. Temporal normalisation is a complex multi-criteria decision. We explicitly mention in the manuscript that we start from the idea that a 5 or 10-day separated acquisition (same viewing) is a priori the best observation available. However, on a case-by-case basis this varies depending on spectral homogeneity, geolocation, or simply cloud coverage.

The difference with the approach from Cusworth et al. needs to be more explicit. From the text, I notice at least two differences (simulated vs real images and the proposed correction term), are they other differences? I also think that it comparing the proposed retrieval approach to the one described in Cusworth et al. can be a valuable additional experiment. As it stands it is difficult to figure out the added value of the proposed quantification methods compared to the one previously presented since no explicit comparison is done.

No comparison with Cusworth et al. was included since the missions and scope of the work are different from this study. The proposed correction term here results in a correction of 15-20% as shown in Fig5. This correction results in a reliable benchmark product which we believe is an important added value.

• Some of the analysis is missing (or should be cited appropriately if it comes from another source), such as 1.170 "it was found that ..." or 1.178 "". I think that the experiments that lead to these assumptions should be presented as well to prove that these are indeed valid assumption.

We have made an extra effort to provide details of the methodology and results. It is possible that minor details that do not affect the content reproducibility are not included in order to reduce the manuscript length and improve readability.

 In section 3, the authors go from a 3D plume simulation to a 2D mass field thus neglecting the impact of the altitude of the plume. This should be clarified in the text (i.e clearly stating the assumptions + justification). Clearly specifying all hypothesis for the study is very important.

We agree with the reviewer that altitude might introduce small differences in the column of dry air. This was mentioned as upcoming improvements in the conclusion section.

 Given that the correction term is scene dependent, It would have been interesting to have Fig.5 for the other sites as well to see if there are noticeable differences between sites.

The correction is scene dependent but variations are expected to be small. The major component of the correction is the downwelling irradiance which will be highly similar between sites in the SWIR due to the high atmospheric transmittance. The surface effect is considered a second order error. Further adaptations of the study to this or other missions, this scene dependence (like the surface) could be simplified with a minor error on the simulated products. In the conclusions section we now completed the paragraph with these comments.

• It is not entirely clear when the temporal normalization is the same as the observation image. I think this should be clarified for the different experiments. Another interesting experiment is the measure of the variability of the estimation as a function of the

reference date. The method recommend using the closest reference but this allows to answer important questions such as "What happens when the closest is a poor reference?" or "Is the closest always the best choice or could another criterium be better?"

The new subsection, "Extending the validation to a large plume dataset and season changes" introduces a discussion about normalization. It brings, for example, the importance of phenological considerations. Different periods of the year might show different vegetation covers (for example, grass in the summer). Cloud and cirrus would definitely be another important criteria. Temporal normalization is a topic that requires understanding of multiple conditions (orbit, phenology, cloud, viewing...). We covered different cases and highlighted some of these issues. However, this will require a dedicated study in upcoming studies. The conclusions section has included a better explanation of the temporal normalization at this point.

• I think that a metric such as the SNR between the plume (delimited by the mask) and the rest of the scene used in Ehret et al. (2022) could be useful to describe the difference behavior in this section.

This is an important suggestion that could be added as a potential metric in future reviews. However, we feel at this point such a metric could be counter-intuitive. The error in the enhancement image is not spatially uncorrelated but contains features (blob-like shapes are mentioned here).

• I suggest that the authors cite the 2021 report of the Intergovernmental Panel on Climate Change that contains a very detailed analysis of the impact of methane emissions on the planet.

Done

 "high-quality calibration and high temporal revisit": I think that this should be clarified. The calibration and temporal revisit is poor compared to other satellites such as Sentinel-5P so I think it would be better to explicit the characteristic of Sentinel-2 instead of just qualifying as "high".

The sentence has been changed to "detailed instrument characterization, and a 5-day temporal revisit"

For me some information are difficult to find in the text. For example, 1.198 the authors state that "methane plume quantification is obtained from isolation of the term ...". It is only two sentence later that it is explained how (I think since no clear link is made). These links must be clearly explicit so that these sections are easier to understand and follow.

This sentence has been rewritten to improve readability.

• I suggest adding an equation that clearly explicit that $L_{B12} = \int_{B12} L_{TOA}(\lambda) d\lambda$ (I.~150)

Done

- The paper is filled with small typos. I will list the one that stood out to me but I'm sure there are many others:
 - 96: missing ref
 - "normalisation" (I.108) vs "normalization" (I.204)
 - 235: "acquitions" -> "acquisitions"
 - 390: 5Ìf0 -> ~50

These changes have been correctly implemented.