

Ann. Geophys. Discuss., author comment AC2
<https://doi.org/10.5194/angeo-2021-68-AC2>, 2022
© Author(s) 2022. This work is distributed under
the Creative Commons Attribution 4.0 License.

Reply on RC2

Chrystal Moser et al.

Author comment on "High bandwidth measurements of auroral Langmuir waves with multiple antennas" by Chrystal Moser et al., Ann. Geophys. Discuss.,
<https://doi.org/10.5194/angeo-2021-68-AC2>, 2022

Response to Reviewer #2

"High bandwidth measurements of auroral Langmuir waves with multiple antennas" by C. Moser, J. LaBelle, and I. H. Cairns

Comment:

1. The subject of Langmuir turbulence in the auroral ionosphere has been reviewed recently by Akbari, LaBelle and Newman (Front. Astron. Space Sci. 7, 617792, 2021) which should be inserted in the reference. In addition, a discussion should be added to explain the source of field-aligned electron beams that excite the auroral Langmuir waves, and if these substorm events are the ionospheric signatures of magnetic reconnection in the tail regions of the magnetosphere.

Response:

The Akbari et al. reference was already in the paper, though mis-cited as 2020 rather than 2021. In addition to correcting the citation, we elaborate further on this reference at lines 22-24. We add information about what types of electron distributions are associated with auroral Langmuir waves at lines 36-39. We further elaborate on the geomagnetic conditions at time of launch at lines 75-77.

Comment:

2. In addition to wave-wave processes involving Langmuir and Lower Hybrid waves, the authors should mention other nonlinear wave interaction studies in the auroral ionosphere. For example, Bohm et al. (JGR 95, 12157, 1990) showed that the most intense Langmuir and whistler waves measured by auroral rocket flights occur in association with Alfvén waves. The theory of auroral Langmuir-Alfvén-whistler events were studied by Chian et al. (A&A 290, L13, 1994) and Lopes and Chian (A&A 305, 669, 1996). In particular, these theoretical papers showed the feasibility of a 3-wave process involving the parametric decay of a Langmuir wave into a whistler wave and an Alfvén electromagnetic ion-cyclotron wave which may apply to the observation of electromagnetic ion-cyclotron waves in a flickering aurora by Lund and LaBelle (JGR 17, 241, 1996).

Response:

These are excellent references and show the common occurrences of non-linear processes in the auroral ionosphere. We've supplemented an existing introductory paragraph on nonlinear Langmuir wave phenomena with these references (see lines 46-54).

Comment:

3. Although the approach of this paper is based on the concepts of linear wave dispersion and weakly nonlinear wave-wave interactions, for the sake of completeness a discussion should be inserted to relate these linear and weakly nonlinear wave studies to the description of Langmuir intermittent turbulence consisted of coherent structures such as cavitons discussed by Akbari et al. (JGR Space Phys. 118, 3576, 2013) and phase-space vortices such as electron and ion holes discussed by Ergun et al. (Phys. Rev. Lett. 81, 826, 1998) and Schamel et al. (Phys. Plasmas 27, 062302, 2020).

Response:

The study by Akbari et al. is very interesting and presents observations somewhat similar to those observed in this paper. For completeness, we have added mention of strong turbulence phenomena in the introduction at lines 36-39. We feel it's unlikely that these strong turbulence phenomena relate to our observations, due to the low energy densities of the waves we observe (wave energy density to thermal energy ratio of order 10^{-12}). For completeness, however, we add a paragraph discussing this topic at lines 317-319.