

Interactive comment on “New insights into mechanisms of sunlight-mediated high-temperature accelerated diurnal production-degradation of fluorescent DOM in lake waters” by Yijun Liu et al.

Yijun Liu et al.

mostofa@tju.edu.cn

Received and published: 3 July 2020

Interactive comment on “New insights into mechanisms of sunlight-mediated high-temperature accelerated diurnal production-degradation of fluorescent DOM in lake waters” by Yijun Liu et al. Dear Prof. Dr. Koji Suzuki, Associate Editor, Biogeosciences: Anonymous Referee 1 Received and published: 2 June 2020. We are very grateful to Review1 for the valuable and constructive comments on our manuscript. We are submitting the manuscript and Figures revised according to the Reviewer comments. We have considered duly all Reviewer comments, providing more examples, which could

[Printer-friendly version](#)

[Discussion paper](#)



contribute to better understanding our FDOM research. Thank you.

Itemized responses (R) to Reviewer comments below: Note: all line numbers refer to the revised manuscript. Major comments The authors measured diurnal changes in CDOM components, DOC concentrations, and nutrients concentrations in two small lakes in Tianjin University, China to identify the biogeochemical processes controlling the diurnal DOM variation which is feasibly related to global warming. I think the research topics described in the manuscript would be of great interests to readers in Biogeosciences. However, I also think that the manuscript is not clearly written and difficult to follow, not technically sound and not appropriately discussed in the context of previous literature. Some of the figures are not clear. Please see comments listed below (1) The authors ran PARAFAC modeling to determine fluorescent components for each time period and compare the components among time periods. First of all, since there is no description regarding how the authors determined the number of components and validated PARAFAC models, I cannot evaluate whether the conclusion derived from PARAFAC are scientifically/technically sound or not. My opinion from our experiences of PARAFAC modeling is that it is not reasonable to apply PARAFAC for a dataset comprising the small number of samples ($n < 20$). It seems that the authors used small data set for PARAFAC modeling (lines 197-200; I cannot understand the sentence though...). While, since the validity depends on the dataset for PARAFAC modeling, the authors should describe the validation method to identify the number of PARAFAC components and show the results of the validation. R-1. "Number of Samples". Our research group has provided ample previous evidence that only two EEM spectra from two samples are enough to identify the authentic PARAFAC components for freshwater and seawater samples (Mostofa et al., 2019; Mostofa et al., 2018-conference paper). In the 2019-EST paper and earlier studies, i.e. Mostofa et al. 2010; Mostofa et al. 2013, we have clearly assessed that "two freshwater samples, six inshore seawater samples and 12 offshore seawater samples are sufficient to run the PARAFAC model". My research group is doing FDOM research since 1999 and I have presented EEM data as a Keynote speaker in the Conference on Organic Matter

Spectroscopy 2018 (WOMS18), held on 23-27 October 2018, in Carqueiranne City, France. Website: <http://woms18.univ-tln.fr/moftofa-abstract/>. In that presentation the number of samples issue, along with many other issues have been discussed providing substantial evidence. Three examples are provided below for further supporting the number of samples used in our PARAFAC model. (i) The PARAFAC model applied to upstream water (Namtaki, site-KR 1) of Kurose River was fitted by only one component (terrestrial humic-like substances) in three cases, i.e. all ten individual EEM data sets on samples from May 2002 to Feb 2003 (Fig. 1a), two EEM data sets on summer (May and July) samples (Fig. 1b) and two EEM data sets on winter (January and February) samples (Fig. 1c). Details about sampling can be found in Mostofa KMG et al. (2005) *Geochemical J* 39: 257-271.

(ii) The PARAFAC model applied to sewerage-impacted downstream water (Izumi, site-KR 5) of Kurose River was fitted two components in three cases, i.e. all 12 individual EEM data sets on samples from May 2002 to Feb 2003 (Fig. 2a-b), two EEM data sets on summer (May and July) samples (Fig. 2c-d), and two EEM data sets on winter (January and February) samples (Fig. 2e-f). Details about sampling can be found in Mostofa KMG et al. (2005) *Geochemical J* 39: 257-271. (iii) The PARAFAC model applied to inshore seawater of Seto Inland Sea was fitted by three components in two cases, i.e. all 12 individual EEM data sets (Fig. 3a-b-c) and six EEM data sets (Fig. 3d-e-f). Details about sampling can be found in Mostofa KMG et al. (2019) *Environ Sci and Technol* 53: 561-563.

R-1. 'Components and Model Validation'. First of all, we wish to describe our knowledge of DOM and FDOM compositions in diverse surface waters. Without understanding of DOM or FDOM composition, without which we cannot discuss about our views on "components and model validation". FDOM are primarily originated from three key sources. First, the terrestrial source, i.e. FDOM derived from soil/land, which is predominantly detected in streams and rivers, and is largely decomposed in lakes/reservoirs/ponds/oceans (Coble, 1996; Mostofa et al. 2013; Mostofa et al.

[Printer-friendly version](#)[Discussion paper](#)

2019). It is well known that extracted soil humic substances are composed of three key components that include humic acids (HA), fulvic acids (FA) and protein-like fluorophores (PLF), with soil PLF entirely different from those detected in surface waters (Fig. 4: below- soil HA (a), FA (b) and PLF (c) of forest soil origin in Mohinuzzaman et al., 2020).

Second, the autochthonous source, i.e. FDOM originated from phytoplankton, which is mostly detected in stagnant waters such as lakes/reservoirs/ponds (Zhang et al., 2009; Parlanti et al. 2000; Mostofa et al. 2013; Yijun et al.- this study). However, some rivers/streams in winter dry season can present stagnant water areas where phytoplankton can grow and produce autochthonous FDOM. As phytoplankton cannot grow in streams or upstream rivers where water is only released from groundwater/terrestrial land, these waters do not contain autochthonous source-FDOM. It is well known that the production of microorganisms/phytoplankton needs time (several days) and stagnant waters for their growth under sunlight conditions. Therefore, stream/upstream river FDOM can only be of terrestrial source and not autochthonous source. Third, an anthropogenic source or sewerage-derived FDOM (e.g. detergent-like components), is detected only in untreated sewerage-impacted rivers in specific site (Baker et al. 2001; Mostofa et al. 2005; Mostofa et al. 2010). Fourth, another important issue is that seawater FDOM is strongly affected by salinity, pH and photochemical degradation, which are completely different from those of freshwater FDOM. For example, salinity can cause the peak C of terrestrial humic-like substances to shift at longer wavelength region (red-shifted phenomena), together with other changes of other FDOM components with respect to river freshwaters (Coble 1996; Yamashita and Zaffé, 2008; Yamashita et al. 2010-Deep-Sea Res-II, 57, 1478-1485; Mostofa et al. 2010; Mostofa et al. 2013; Mostofa et al. 2019). Fifth, in this study (Yijun et al. 2020; see Figure 5) and other studies (Zhang et al., 2009; Mostofa et al. 2013; Parlanti et al. 2000). Terrestrial humic acid and terrestrial humic-like substances (C type) show fluorescence properties (peak positions and EEM images) similar to those of autochthonous humic-like substances (C type) of phytoplankton origin The two autochthonous and terrestrial

[Printer-friendly version](#)[Discussion paper](#)

sources of FDOM components cannot be distinguished by PARAFAC analysis when all diverse samples are mixed together. This is the reason why one cannot distinguish between these two autochthonous sources.

Sixth, in this and other studies (Yijun et al. 2020; see Figure 6; Zhang et al., 2009; Mostofa et al. 2013; Shammi et al. 2017) terrestrial fulvic acid and terrestrial humic-like substances (M type) (Mostofa et al. 2019; Mostofa et al. 2013; Mohinuzzaman et al. 2020) show fluorescence properties (peak positions and EEM images) similar to autochthonous humic-like substances (M type) of phytoplankton origin). The two autochthonous (Fig. 6a-d) and terrestrial (Fig. 6e-g) sources of FDOM components cannot be distinguished by PARAFAC analysis when all diverse samples are mixed together. This is the reason why one cannot distinguish between these two autochthonous sources.

Seventh, untreated sewerage-affected river could transport detergent-like substances and standard household detergents into river waters (see Fig. 7a-c; Mostofa et al. 2013; Mostofa et al. 2010; Mostofa et al. 2005; Baker et al. 2001).

Based on all seven different types of FDOM sources, it is very important to choose “Selective Characteristic Samples” that can represent authentic FDOM components (Mostofa et al. 2019), whereas mixing all seven types of FDOM sources together in a PARAFAC model cannot evaluate authentic FDOM. Stedmon et al. (2003) firstly used the PARAFAC model on a total of 90 samples covering all major terrestrial sources of DOM, streams and estuary DOM mixed with seawater from the adjacent Kattegat, which resulted in a five component model. Since the study of Stedmon et al. (2003), the use of mixed samples in the PARAFAC model has been repeated in several other studies to date (Murphy et al., 2013; Kulkarni et al., 2017; Wu et al., 2018; Yue et al., 2019). However, artifact due to mixing. Mostofa et al. (2019) provided evidence that these components were artifacts due to mixing. “Model Validation”. Stedmon et al. (2003) and Murphy et al. (2013) have used the split-half analysis method for validation of the PARAFAC components. This method consists in dividing the EEM data set into

[Printer-friendly version](#)[Discussion paper](#)

two random, typically equal sized groups and then applying the PARAFAC model to both data halves independently (Stedmon et al., 2003). The procedure of mixing all diverse samples appears to be inappropriate to validate the PARAFAC model. Furthermore, this model validation based on split-half analysis of unselected EEM data does not allow to identify and distinguish the authentic fluorescent components in the PARAFAC model. Thus, we believe that a correct model validation should be based on the authentic identification and characterization of fluorescent components of the individual seven sources mentioned above. In this study, we have validated the authentic fluorescent components on the basis of our earlier experiences on the identification and classification of EPS and their released photomicrobial components (Shammi et al. 2017a, 2017b, 2017c; Sheng and Yu, 2006), autochthonous humic-like substances (C-type and M-type) of phytoplankton origin (Zhang et al., 2009; Mostofa et al., 2013), as well as aromatic amino acids (e.g. tryptophan, tyrosine and phenylalanine) of phytoplankton origin and standard substances (Yamashita and Tanoue, 2003; Zhang et al., 2009; Mostofa et al., 2013, 2018a, 2018b, 2019). According to the comments above, we have revised the manuscript with addition of a new paragraph “The number of components and validation of PARAFAC model were performed based on the following criteria. Firstly, we applied ‘Selective Characteristic Sample’ EEM data to the PARAFAC model that could be continued from 1 to 8 components until achieving negative a -values (see Eqs 1-2). When two successive models (e.g. the 5- and 6-component model) showed one or two negative a -values, the model could be fitted up to 4 components and there was no need to go further by checking for a 7- or 8-component model. Then, the valid models were subjected to the next step for their validation by considering the highest number of fluorescent components and comparing with their respective EEM images and fluorescence peaks of standard and extracted substances (or FDOM) (Yamashita and Tanoue, 2003; Sheng and Yu, 2006; Zhang et al. 2009; Mostofa et al., 2013; Shammi et al. 2017a, 2017b, 2017c; Mostofa et al., 2019). For example, if we consider a 4-component model where the image and fluorescence peaks of one of the four fluorescent components does not correspond to the respective peak position appearing

[Printer-friendly version](#)[Discussion paper](#)

outside the usual DOM fluorescence (i.e. Rayleigh and water Raman peak positions), then the model should be rechecked for 3 components. Differently, if all components are verified with those of standard substances or specific field/experimental observations, the model and the respective fluorescent components are validated. Finally, the fluorescent components obtained from the model can be characterized and classified (Yamashita and Tanoue, 2003; Sheng and Yu, 2006; Zhang et al. 2009; Mostofa et al., 2013; Shammi et al. 2017a, 2017b, 2017c; Mostofa et al., 2019).” in lines 206-225. We believe that it is important to find out the various sources of FDOM which can be related to biogeochemical facts and their significances. To clarify the principles on which the PARAFAC model is based, we have also added an additional paragraph “The parallel factor (PARAFAC) analysis is a three-way multivariate method that can be applied to EEM data of DOM to decompose them into trilinear components (Harshman. 1970; Caroll and Chang, 1970). For any fluorophore of FDOM, the emission intensity, x_{jk} , at a specific wavelength, j , which corresponds to excitation at the wavelength, k , can be expressed by the following equation (Harshman. 1970):

where a_i is the concentration (in arbitrary units, a.u.) of the analyte (fluorophore or FDOM), b_{ij} is the relative emission at the wavelength j , and c_{ik} is the relative amount of light absorbed at the excitation wavelength k . For any number of analytes and samples, the PARAFAC model can be expressed as a set of trilinear terms and a residual array as (Stedmon et al., 2003):

where x_{ijk} is the fluorescence intensity of the i th sample at the emission wavelength j and excitation wavelength k , a_{if} is directly proportional to the concentration (in a.u.) of the analyte f in sample i . b_{jf} is directly proportional to the quantum efficiency of fluorescence of the analyte f at the emission wavelength j , c_{kf} is linearly related to the specific absorption coefficient at the excitation wavelength k , F is the number of components in the model, and ϵ_{ijk} is the residual matrix that indicates the variability not accounted for by the model.” in lines 182-198.

(2) The authors concluded that some components were disappeared over a 24-h diur-

[Printer-friendly version](#)[Discussion paper](#)

nal period (e.g., lines 29-32) from the qualitative analysis by PARAFAC. I do not agree with this conclusion, because it may be results of artifact by PARAFAC modeling. For example, if the authors apply new PARAFAC with slightly large number of samples, the disappeared components may appear. If the authors want to conclude disappearance of components quantitatively, the authors should use single PARAFAC model and determine diurnal changes in fluorescence intensity of individual PARAFAC components. I also disagree the conclusion of “degraded into four FDOM components (lines 32-35)” with the same reason. R-2. We have extensively discussed above that this study does not include any artifact components. A large number of water samples from different sources can produce artifact components from their mixing (Mostofa et al. 2019). The disappearance of FDOM occurred only in July samples under strong sunlight and high water/air temperature conditions and did not occur in October and June samples. Disappearance of FDOM is possibly due to its conversion into low molecular weight DOM which usually do not show any fluorescence. Furthermore, terrestrial humic-like substances (C-type) do not degrade and only autochthonous FDOM can degrade as we showed in this study. The July changes in FDOM can be further evidenced from inconsistent changes of DOC concentrations that significantly increased at noon (18Furthermore, EPS is the first component that we have comprehensively discussed in the manuscript. Earlier results (Shammi et al. 2017a, 2017b) demonstrated that raw EPS (A: 3 components) can convert rapidly into three new components in one to six-hours irradiation, which is supported by results of this current study. Then, after 18 to 54 hours, two components are completely degraded. In our July samples, degradation occurred at the highest air temperature (41.13 °C) and water temperature range (29.2-33.5 °C). To better clarify these aspects, we have revised the relevant sentence “at the highest air and water temperatures (respectively, 41.1 and 33.5°C)” in lines 31-32. Full sentence: In this work sunlight-mediated high-temperature was shown to accelerate the production of FDOM, but also its complete disappearance over a 24-h diurnal period in July at the highest air and water temperatures (respectively, 41.1 and 33.5°C), but not in lower temperature months). (3) The authors described temporal

[Printer-friendly version](#)[Discussion paper](#)

changes in PARAFAC components quantitatively (e.g., lines 232-234). However, I cannot follow the time periods which involving the temporal changes. Again, it is difficult to quantitatively compare abundance of PARAFAC components derived from different PARAFAC models. R-3. We have rearranged the sentences and revised properly “In particular: (a) newly-released PLS with a peak T at 280/351 nm appeared and the TUV peak disappeared, whereas the intensity of peak T of newly-released PLS increased by 49%. The changes of FDOM components occur due to microbial respiration/ sunlight degradation at every moment, which simultaneously altered nutrients (NO₃⁻, NH₄⁺, etc.) and DOC contents. FDOM components are changing with changes in sunlight intensity during day time and microbial degradation at night time. Therefore, PARAFAC model on time-specific water samples are important, as well as the seven-specific FDOM components, to conduct PARAFAC analysis individually. Otherwise, we cannot observe the biogeochemical changes of FDOM components which are related to changes in nutrients, DOC concentration and phytoplankton. (4) The authors seemed to conclude that EPS produced during early morning partially degraded at midday and remineralized at night time (lines 32-35). Is this conclusion consistent with temporal changes in DOC concentrations and nutrients concentrations (Fig. 6)? I don't think so. To verify the authors' conclusion, careful, quantitative, and statistical comparisons of diurnal changes in PARAFAC components with diurnal changes in DOC and nutrients concentrations should be made. R-4. We have partly clarified this issue in the reply to comment 2. This issue is based on the various biogeochemical changes that occur at each timescale and involves the following processes: • Phytoplankton via photosynthesis is constantly produced during sunlight period. • Photoinduced and microbial respiration can release various FDOM from phytoplankton, which changes (generally increases) DOC and DON contents. • The occurrence of photodegradation by photo-induced reactive oxygen species (O₂⁻, H₂O₂ and •OH) during day time varies depending on sunlight intensity. • The occurrence of microbial degradation of FDOM can alter FDOM, nutrients and DOC. • The conversion of FDOM by photochemical and microbial processes into low molecular weight (LMW)-non-fluorescent

[Printer-friendly version](#)[Discussion paper](#)

DOC and nutrients. The complete mineralization of DOC occurs constantly under photo-microbial processes. Nitrification, denitrification and anaerobic ammonium oxidation processes are occurring constantly. As all of these processes are occurring actively at every moment, the FDOM, DOC and nutrients we have measured at specific times (e.g. 10 am) are the net existing amounts deriving from all the mentioned biogeochemical changes, in “4.2 Biogeochemical processes involving DOC and nutrients in lake Jingye” In July, DOC concentration varied hourly from a minimum of 815 μM C achieved in the night (from 21.00 to 6.00) to a maximum of 963 μM C reached during the day (from 10.00 to 16.00), with the highest DOC fluctuation of 18. The highest production of nutrients in October than in July was estimated to be approximately of 16.0, 28, 4.0 and 23.8-fold respectively for NO_3^- , NO_2^- , NH_4^+ , and PO_4^{3-} (Fig. 6). In particular, in October the highest NH_4^+ concentration paralleled the highest DOC concentration detected at the same time (14.00) (Table S2, Fig. 6), which suggested a rapid biogeochemical transformation of organic matter into DOC and nutrients under ambient lake conditions. Apparently, the NH_4^+ formation in October was followed by its subsequent rapid nitrification and denitrification at same time (14.00) (Table S2, Fig. 6). Similarly, the lowest level of NH_4^+ at 10.00 was consistent with the lowest level of NO_3^- . Further, the occurrence of nitrification was confirmed by the decrease of NH_4^+ contents occurring simultaneously to the increase of NO_3^- in early morning samples (Table S2, Fig. 6). Further evidence of photosynthetic activities of phytoplankton was provided by the significant shifting of the $\delta^{15}\text{N}$ value of NO_3^- . In particular, in July the $\delta^{15}\text{N}$ value decreased from +0.67‰ at 10:00 to -0.02‰ at 12:00, which corresponded to the increase of SI from 2.33 MJ m^{-2} to 2.95 MJ m^{-2} (for a total of 8.14 MJ m^{-2}), whereas the $\delta^{18}\text{O}$ value of NO_3^- increased from +5.28‰ at 10:00 to +9.3‰ at 12:00 under conditions of elevated SI and high WT (36.5-39.9 °C) and AT (31.0-32.7 °C). The decrease of $\delta^{15}\text{N}$ and increase of $\delta^{18}\text{O}$ in NO_3^- with increasing SI from 10:00 to 12:00 indicated the uptake of lighter ^{16}O -containing NO_3^- by phytoplankton with increasing photosynthesis. This effect was confirmed by the 20We have also partly discussed this issue based on the reviewer2 comments in the new addition “4.4. High-

[Printer-friendly version](#)[Discussion paper](#)

temperature production of FDOM from phytoplankton and its sequential entire degradation at 24-h diurnal scale. The complete degradation of FDOM after its formation from phytoplankton can be ascribed to the gradual increase in SI (from 1.36 to 2.97 MJ/m² for a total of 15.26 MJ/m²) at the highest WT (30.2-33.7 °C) and AT (35.4-41.8 °C) (Fig. 2). These environmental conditions are directly responsible of the rapid production of FDOM from phytoplankton and its corresponding rapid degradation by day-time sunlight-induced and night-time microbially-induced processes. Such entire sequential diurnal degradation of FDOM was not observed in October, May and June samples (Figs. 3-5) when solar intensity, AT and WT were relatively low compared to July samples (Fig. 2). As discussed previously, the production and degradation of FDOM are related to simultaneous significant fluctuations in the contents of NO₃⁻, NH₄⁺, NO₂⁻, DON, PO₄³⁻, DSi and DOC (Fig. 6; Table S2). Photochemical and microbial processes can transform EPS into various forms of FDOM (Shammi et al., 2017a, 2017b) that are related to field observations, which show that different components of EPS vary over different timescales and temperatures (Shammi et al., 2017c; Sheng and Yu, 2006). Many studies provided evidence that photochemical and microbial degradation processes increase with increasing temperature and light intensity (Matsumoto et al., 2007; Weston and Joye, 2005; Malinverno and Martinez, 2015; Whelan and Rhew, 2015; McKay and Rosario-Ortiz, 2015; Grannas et al., 2006; Farias et al., 2007). In turn, these results would indicate that high SI, WT and AT would accelerate the complete transformation of autochthonous FDOM into LMW DOM and other mineralization end-products on a 24-h diurnal cycle, which could be further increased by the influence of future GW. Such changes can to be reasonably expected to occur in the future on the basis of increasing water temperature, extended summer season and increased water stratification in response to the predicted GW from 1.5 to 2.0°C (Huisman et al., 2006; Watanabe et al., 2011; Rogelj et al., 2019).”

(5) It seems that the authors generally consider/discuss degradation of EPS to explain daytime changes in PARAFAC components. Why don't the authors consider DOM production including the PARAFAC components with primary production during daytime?

[Printer-friendly version](#)[Discussion paper](#)

R-5. DOM is well known to comprise many types of organic substances, which include EPS that can gradually convert into various FDOM components, non-fluorescent photoproducts that can also be produced from FDOM, non-fluorescent microbial products, non-fluorescent LMW substances (aldehydes, ketones), etc. Besides terrestrial sources, autochthonous DOM is a key fraction of DOM compositions in lakes and reservoirs, but it is still uncertain how it is released from phytoplankton. Our research group has gained a wide experience on EPS extraction from large volume of surface waters, and its photo-microbial products, which is the key to better understand the relation between FDOM and DOM. We believe that results of this study will contribute to understand the sequential release of FDOM from EPS and their subsequent degradation (Figs. 8-9; this study). As commonly reported in all other studies, in this manuscript DOM is discussed as DOC changes.

Other comments I listed some other comments below. Similar errors with some of the comments were found throughout the manuscript. Lines 69-72: What is “key DOM components”? It’s not clear. R. The sentence has been revised according to this comment with addition of “Key DOM components include terrestrial humic substances (fulvic and humic acids), EPS, and aromatic amino acids (Coble, 1996; Yamashina and Tanoue, 2003; Yamashina and Tanoue, 2004; Shammi et al. 2017a, 2017b, 2017c; Zhang et al., 2009). Lines 99-101: I could not understand the message of the sentence. The authors would like to mention “diurnal degradation processes depended on the photosynthetic activity of primary producers”? R: Yes. The sentence has been rearranged according to this comment as “Diurnal day-time (sunlight) and night-time (microbial) degradation processes are a natural phenomenon that depends on the photosynthetic activity of primary producers in surface waters which is ultimately related to daily biogeochemical changes of C, N and P cycling”.

Lines 102-105: “a significant decrease of its fluorescence intensity with increasing water depth” is due to photoinduced degradation? I don’t agree with it. R. Yes, we agree. The sentence has been revised properly in lines 108-109. Line 105: Which

[Printer-friendly version](#)[Discussion paper](#)

overall day-night degradation of FDOM corresponds “such”? R. The sentence has been moved earlier sentence, as it is connected with the previous sentence. “Diurnal day-time (sunlight) and night-time (microbial) degradation processes are a natural phenomenon that depends on the photosynthetic activity of primary producers in surface waters which is ultimately related to daily biogeochemical changes of C, N and P cycling (Guidi et al., 2016; Segschneider and Bendtsen., 2013; Huisman et al., 2006; Carpenter et al., 1998). Such overall day-night degradation of FDOM is caused from diurnal photo-microbial transformations.” Fig. S1: The figure is not clear. It may be better to show a campus map with map of Tianjin city. R. The maps have been replaced as suggested (see Figure 9). Line 134: Do seagrasses occur in inland closed lakes R. Yes, there are some seagrasses in the Qingnian lake. Line 149: Add filter name and company name supplied it. R. Revised as suggested (....0.45- μ m glass-fiber filter (GF/F, Shanghai Xin Ya Purification Equipment Co. Ltd) previously cleaned....). Lines 152-153: “a combustion total organic carbon (TOC) auto-sampler analyzer” is awkward. R. Revised as “.... using a total organic carbon (TOC) analyzer (OI Analytical Aurora,.....”. Lines 154-155 168-169: The authors described a method for UV-Vis absorbance twice. In addition, the authors described different spectrophotometers, respectively. R. Revised as “A UV-VIS spectrophotometer (UV-2700, Shimadzu) was used to estimate absorption properties of chromophoric DOM (CDOM). Ultra-pure water was simultaneously used in the reference cell as a blank.”. We deleted the duplication in former lines 167-168. Lines 171-180: How did the authors calibrate the fluorescence intensity? With quinine sulfate? Also, what is “700 v” in line 174. R. Arbitrary units have been used in this study. ‘700 v’ has been revised as “...and a photomultiplier voltage of 700 v.” in lines 175-176. Line 209: Define “early stage DOM”. R. The text has been revised as suggested by addition of “Generally, EPS are considered an early-stage DOM, defined as “dissolved organic substances newly-formed and not yet converted to individual organic components”, which”. Lines 223-226: I think no reference cited here describe “FDOM production by degradation of EPS would occur during the gradual increase of SI”. R. All references in line 261 have been deleted

[Printer-friendly version](#)[Discussion paper](#)

from the mentioned sentences “FDOM production by degradation of EPS would occur during the gradual increase of SI (from 1.36 to 2.97 MJ/m² for a total = 15.26 MJ/m²) at WT of 30.2-33.7 °C and AT of 35.4-41.8 °C (Fig. 2).”.

Lines 245-247: I could not follow how the authors estimate the abundance of EPS quantitatively. R. The text has been revised as “. . . . In particular, the decrease in fluorescence intensity of EPS in this time period was approximately 3Line 254: The citation of Ma and Green (2004) is not appropriate for the sentence, because their work was carried out Lake Superior. R. Reference deleted in line 288-289 and revised according to our own results with addition of “(b) night-time extended microbial degradation, which was also supported by the net decrease of DOC along with nutrients at night period, which will be comprehensively discussed in the next section.”. Line 415: What is “photoinduced respiration of phytoplankton”? In addition, it seems that the authors discuss the photosynthesis rather than respiration at the rest of this paragraph. R. The text has been revised as “Further evidence of photosynthetic activities of phytoplankton was provided by the significant shifting of the $\delta^{15}\text{N}$ value of NO_3^- .”. Lines 439-465: I think the authors do not discuss “A global view of production and degradation pathways of FDOM in lake waters” in this paragraph. R. For reviewer’s kind information. Our group published previously on EPS extraction at large scales using 40 liters of surface waters and on the subsequent photo-microbial experiments that can sequentially release FDOM from EPS (Shammi et al. 2017a, 2017b, 2017c). The results of our current study, as well as some unpublished data, appear to confirm all previous results on FDOM, EPS and their byproducts. Although it is till now unclear how autochthonous FDOM originates from phytoplankton, the results of this study demonstrated that EPS are produced in the morning and then convert into many autochthonous FDOM components that are ubiquitously detected in surface waters, as previously reported in experimental observations (Zhang et al., 2009). The paragraph in question attempts to provide a conceptual model based on our research findings by describing the sequential release of FDOM, which ultimately describes dynamics of DOM that originates from phytoplankton. This global view contributes to further under-

standing autochthonous FDOM and its degradation processes. Line 435: I could not
And Figs 12 and 13. R. Sorry, these are Fig. 8 and 9 as “.can be illustrated in the
pathway shown in Fig. 7 and detailed in Figs. 8 and 9.”. Fig. 2: The figure caption,
in particular about (b), does not explain the figure. R. Figure caption corrected. Fig.
6: Describe which Y-axis corresponds to each plot in all figures except the figure
(e). Why does the Y-axis on right side of Fig. 6d show negative values? R. Figure 6
has been revised as suggested (see Figure 10), and negative values for dissolved Si
have been deleted.

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2020-145>, 2020.

BGD

Interactive
comment

Printer-friendly version

Discussion paper



Figure 1. Review response-1 (R-1) examples for “Number of Samples”.




Fig. 1. Figure 1. Review response-1 (R-1) examples for “Number of Samples”.

Figure 2. Review response-1 (R-1) examples for “Number of Samples”.




Fig. 2. Figure 2. Review response-1 (R-1) examples for “Number of Samples”.

Figure 3. Review response-1 (R-1) examples for “Number of Samples”.




Fig. 3. Figure 3. Review response-1 (R-1) examples for “Number of Samples”.

Printer-friendly version

Discussion paper



Figure 4. Author response (R-1) for ‘Components and Model Validation’.




Fig. 4. Figure 4. Author response (R-1) for ‘Components and Model Validation’.

Figure 5. Author response (R-1) for ‘Components and Model Validation’.




Fig. 5. Figure 5. Author response (R-1) for ‘Components and Model Validation’.

Figure 6. Author response (R-1) for ‘Components and Model Validation’.




Fig. 6. Figure 6. Author response (R-1) for ‘Components and Model Validation’.

Figure 7. Author response (R-1) for ‘Components and Model Validation’.




Fig. 7. Figure 7. Author response (R-1) for ‘Components and Model Validation’.

Figure 8. Author response (R-2) for review comments by Reviewer#1




Fig. 8. Figure 8. Author response (R-2) for review comments by Reviewer#1

Figure 9. Author response for specific comment by reviewer#1




Fig. 9. Figure 9. Author response for specific comment by reviewer#1

Printer-friendly version

Discussion paper



Figure 10. Author response for specific comment by reviewer#1




Fig. 10. Figure 10. Author response for specific comment by reviewer#1

Printer-friendly version

Discussion paper

