

Biogeosciences Discuss., author comment AC3  
<https://doi.org/10.5194/bg-2021-234-AC3>, 2021  
© Author(s) 2021. This work is distributed under  
the Creative Commons Attribution 4.0 License.



## Reply on RC3

Yuanyuan Zhang et al.

---

Author comment on "Methane accumulation affected by particulate organic carbon in upper Yangtze deep valley cascade reservoirs, China" by Yuanyuan Zhang et al., Biogeosciences Discuss., <https://doi.org/10.5194/bg-2021-234-AC3>, 2021

---

We feel sorry for the misunderstanding and inaccurate response about this question and thank the Referee #2 for pointing out this mistake.

At first, autochthonous organic matter was derived from aquatic production. Our research site, i.e., the Xiangjiaba Reservoir, is a deep-valley large reservoir located in the upper Yangtze River, China. Unlike streams, creeks or some other shallow lakes where dominant primary producers are periphyton communities (Piggott et al., 2015; Trochine et al., 2017), phytoplankton communities contributed almost all the primary productivity and autochthonous organic matter in water column in large water bodies, such as the Xiangjiaba Reservoir. Even there would be periphyton communities who contributed partial autochthonous organic matter in the reservoir, we believed that the relative abundance of periphyton-derived organic matter was negligible because of such large water bodies.

In our sampling campaign, as indicated in the manuscript, water samples were collected at the main channel of the reservoir. The phytoplankton and zooplankton communities were collected in water by phytoplankton net (diameter of mesh pore 64  $\mu\text{m}$ ) and sorted under microscope to represent the endmember of the autochthonous particulate organic matter. This approach was also discussed in some of the previous studies (Grasset et al., 2018; Chen et al., 2018; Tittel et al., 2019; Bueno et al., 2020).

Here, we would like to withdraw our previous mistake, and keep the term "plankton" in the manuscript. If there will be any questions, please don't hesitate to contact us at [lizhe@cigit.ac.cn](mailto:lizhe@cigit.ac.cn).

## References

Bueno, C.C., Frascareli, D., Gontijo, E.S.J., van Geldern, R., Rosa, A.H., & Friese, K., Barth, J.A.C.: Dominance of in situ produced particulate organic carbon in a subtropical reservoir inferred from carbon stable isotopes, *Scientific Reports*, 10: 13187, 2020.

Chen, J., Yang, H., Zeng, Y., Guo, J., Song, Y., & Ding, W.: Combined use of radiocarbon and stable carbon isotope to constrain the sources and cycling of particulate organic carbon in a large freshwater lake, China, *Science of the Total Environment*, 625, 27-38, 2018.

Grasset, C., Mendonça, R., Villamor Saucedo, G., Bastviken, D., Roland, F., & Sobek, S.: Large but variable methane production in anoxic freshwater sediment upon addition of allochthonous and autochthonous organic matter, *Limnology and Oceanography*, 63, 1488-1501, 2018.

Piggott, J. J., Salis, R. K., Lear, G., Townsend, C. R., & Matthaei, C. D.: Climate warming and agricultural stressors interact to determine stream periphyton community composition, *Global Change Biology*, 21, 206-222, 2015.

Tittel, J., Hüls, M., & Koschorreck, M.: Terrestrial vegetation drives methane production in the sediments of two German reservoirs, *Scientific reports*, 9, 1-10, 2019.

Trochine, C., Guerrieri, M., Liboriussen, L., Willems, P., Lauridsen, T. L., Søndergaard, M., & Jeppesen, E.: Factors controlling the stable isotope composition and C: N ratio of seston and periphyton in shallow lake mesocosms with contrasting nutrient loadings and temperatures, *Freshwater Biology*, 62, 1596-1613, 2017.