

Biogeosciences Discuss., community comment CC1
<https://doi.org/10.5194/bg-2021-3-CC1>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Comment on bg-2021-3

Antti Rissanen

Community comment on "Methane oxidation in the waters of a humic-rich boreal lake stimulated by photosynthesis, nitrite, Fe(III) and humics" by Sigrid van Grinsven et al., Biogeosciences Discuss., <https://doi.org/10.5194/bg-2021-3-CC1>, 2021

It was really interesting to read a study on methane oxidation conducted in the same study lake as our recent study (Rissanen et al. FEMS Microb Ecol, Volume 97, Issue 2, February 2021, f1aa252). While our study was focused on the genetic potential of methanotrophs, this study provides very valuable novel information on the various factors affecting the activity of methanotrophs in boreal lakes. Here are some minor suggestions, which I think could further improve the manuscript:

Line 49-52. It is perhaps worth to mention here study by Zheng et al. 2020 on extracellular electron transfer from methane to Fe-mineral by *Methylomonas* in hypoxic conditions.

<https://pubs.acs.org/doi/abs/10.1021/acs.estlett.0c00436>

Line 69-70. Either "can play" or "are likely to play"

Line 73-74 and in especially discussion. Maybe it would be relevant to mention and discuss your results also in the light of our previous study from 2018, which was conducted in the nearby humic boreal lakes, where we also studied lake water methane oxidation with amendments of various EAs (incl. NO₃, metals, organic EAs) and in different light conditions, and also studied the community structure and genetic potential of methanotroph communities (Rissanen et al. 2018):

<https://www.int-res.com/abstracts/ame/v81/n3/p257-276/>

Furthermore, study by Kallistova et al. (2018) might be also relevant to mention and discuss. They also studied methane oxidation and MOB communities in water column of a boreal lake.

<https://www.int-res.com/abstracts/ame/v82/n1/p1-18/>

Line 87-. Study site. It is maybe worth to mention the historical anthropogenic effects, the

soaking of flax and hemp, which potentially have contributed to the (chemical) stratification in the lake. In the Finnish publication (Tolonen et al. 1976), it is mentioned in Finnish that "Hampun ja myöhemmin myös pellavan liotus nopeuttivat läheisen Lovojärven pilaantumista jo rautakaudella (Huttunen & Tolonen 1975)." = Soaking of hemp and later also soaking of flax accelerated the pollution of nearby Lake Lovojärvi already during Iron Age".

Tolonen K, Tolonen M, Honkasalo L et al. Esihistoriallisen ja historiallisen maankäytön vaikutuksesta Lammin Lampellonjärven kehitykseen. Luonnon Tutkija. 1976;80:1-15 (in Finnish with English abstract):

https://www.researchgate.net/publication/311665698_Esihistoriallisen_ja_historiallisen_maankayton_vaikutus_Lammin_Lampellonjarven_kehitykseen_The_influence_of_of_prehistoric_and_historic_land_use_on_Lake_Lampellonjarvi_South_Finland

Unfortunately, I could not find the original reference of Huttunen & Tolonen 1975.

Line 118. were fixed

Line 382- What about archaea-driven methanogenesis in anoxic micro-niches?

Also more generally, there are recent studies suggesting that also methanogenic archaea can oxidize methane anaerobically, e.g. via extracellular electron transfer to solid EAs (iron minerals, organic EAs, anode in bioelectrochemical systems). Maybe worth to mention and discuss. See, e.g.

<https://www.sciencedirect.com/science/article/abs/pii/S1385894720328199>

<https://pubmed.ncbi.nlm.nih.gov/28965392/>

Line 468-470. Our study in the same study lake (which has been cited but not in this context) detected genetic potential for nitrate/nitrite/NO⁻ reduction as well as for extracellular electron transfer (to metal minerals and organic EAs) in metagenome assembled genomes of Methylococcales (incl. Methylobacter sp. and the Crenothrix-type MOB), which supports the results of this study on enhancement of methane oxidation by these various EAs. See:

<https://academic.oup.com/femsec/article/97/2/fiaa252/6034011>

Line 473-481. Microbial interactions. Maybe worth to include and discuss also the results by Cabrol et al. 2020. They studied anaerobic methane oxidation (AOM) and MOB communities in water columns of northern lakes and found correlation between Methylococcales and OTUs within Methylothera, Geothrix and Geobacter genera which indicated that AOM might occur in an interaction between MOB, denitrifiers and iron-cycling partners.

<https://www.sciencedirect.com/science/article/pii/S0048969720331053>

Figure 1. Is there a slight increase in O₂ towards the deepest depths (from appr. 15 m to the deepest depth)? If there is, what is causing it?

References:

Kortelainen et al. 2000. Numbers as subscripts for CH₄, CO₂ and N₂O

Mutyaba 2012. Maybe it could be mentioned that it is Master of Science thesis. And perhaps provide a link to it. <https://jyx.jyu.fi/handle/123456789/40735>

Rissanen et al. DOI-link is missing.