

Biogeosciences Discuss., community comment CC1
<https://doi.org/10.5194/bg-2021-69-CC1>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Downstream contribution to C mass balance

Kleiton Rabelo de Araújo

Community comment on "Partitioning carbon sources between wetland and well-drained ecosystems to a tropical first-order stream – implications for carbon cycling at the watershed scale (Nyong, Cameroon)" by Moussa Moustapha et al., Biogeosciences Discuss., <https://doi.org/10.5194/bg-2021-69-CC1>, 2021

The manuscript "Partitioning carbon sources in a tropical watershed (Nyong River, Cameroon) between wetlands and terrestrial ecosystems – Do CO₂ emissions from tropical rivers offset the terrestrial carbon sink?" by Moustapha et al. discusses CO₂ emissions in tropical rivers and the role of floodplains as organic C importers along with groundwater. Since tropical continental areas are hotspots of C emission, the evaluation of mechanisms associated to C degradation and transport in these regions are essential to determine its implications on C budget. Therefore, it is an important work to the field with unquestionable publication interest. However, part of the experimental design is still not clear. The most downstream tributaries are neglect from the sampling, however a C mass balance of organic C exported to ocean is presented. There is a thesis, Nkoue-ndondo, 2008, embaying that Olama was the most downstream site with representative contribution to C export. However, the argument is not robust enough to support that downstream rivers can be neglected. I would suggest the authors to present a brief description about downstream C deposition and degradation. If the downstream geomorphology and discharge does not favor deposition and C oxidation, downstream tributaries may be neglected. Otherwise, the C mass balance should be adjusted.