

Biogeosciences Discuss., referee comment RC1
<https://doi.org/10.5194/bg-2022-31-RC1>, 2022
© Author(s) 2022. This work is distributed under
the Creative Commons Attribution 4.0 License.

Comment on bg-2022-31

Anonymous Referee #1

Referee comment on "Modelling the effects of benthic fauna on carbon, nitrogen and phosphorus dynamics in the Baltic Sea" by Eva Ehrnsten et al., Biogeosciences Discuss., <https://doi.org/10.5194/bg-2022-31-RC1>, 2022

General comments

The manuscript investigates the effects of benthic fauna on the biogeochemical transformations of major elements in the Baltic Sea via direct (metabolic) and indirect (bioturbation) pathways, by coupling a recently developed model of benthic fauna to an established hydrodynamic-biogeochemical model. Within regional ecosystem models, benthic components are often represented in a simplified way, where their role is largely limited to returning remineralised nutrients back into the water column. In the Baltic Sea, models accounting for indirect impacts of benthic fauna have been used for a while now, incentivised recently by the need to study the effects of bioturbation by invasive species. However, these models did not consider benthic faunal biomass explicitly, and thus did not allow a separation of effects of bioturbation and direct effects of faunal metabolism on benthic-pelagic dynamics. This manuscript bridges this gap and provides a detailed look at the role benthic fauna plays in shaping biogeochemical fluxes in the Baltic Sea. This study builds on the authors' previous work and contributes to the advancement of benthic system modelling, which has lagged behind its pelagic counterpart in coupled regional models. Although it is an important step in the right direction, I have several criticisms I would like to bring to the attention of the authors. Foremost they are related to the "large-scale" approach taken for this work. It should not serve as a substitute to thorough comparison of model results with data, nor to the detailed presentation and discussion of results (providing quantitative estimates and aligning text narrative with figures). Limitations of the model, unique contributions of this work to the understanding of the Baltic Sea biogeochemistry beyond state-of-the-art knowledge should be appropriately discussed. Below are my detailed comments and questions, that I hope will improve the quality of the manuscript.

Specific comments

Title

1. The title should clearly indicate that it is a model-based study.

Abstract

1. It would be informative to add quantitative estimates of changes due to direct and indirect contributions of benthic fauna. "Small proportion of seafloor organic stocks", "decreases denitrification", "increases P retention", "reduction in N fixation" etc all should be accompanied with quantitative estimates (% change). The way information is currently presented, it is hard to see what a unique contribution of this work is to the understanding of biogeochemical impacts of benthic fauna in the Baltic Sea, as the effects of bioturbation *via* sediment oxygenation on N, P dynamics in the Baltic Sea are generally well known from previous studies.
2. Line 16: It should be clear from the text that bioturbation affects denitrification, P retention etc *indirectly*, e.g., increasing oxygen penetration depth and availability.
3. Line 20: "chain of indirect effects" is opposed to "direct effects of faunal respiration, excretion and bioturbation". This create some confusion over which effects are "direct" and which are "indirect" ones.

Introduction

1. Lines 43-44. "The combined effects of animal bioturbation and metabolism have seldom been studied together." Several references support this statement, but could the authors elaborate on what are the main reasons for that gap in research, despite recognition of the importance of benthic processes?
2. Could you provide a more solid justification for your approach? Why do we need to implement modelling – does it address the knowledge gap? Why did biogeochemical models of the Baltic Sea so far did not include benthic fauna?
3. When specifying reasons for using Baltic Sea as a model area, to what extent also relatively simple benthic community composition play role?

Materials and methods

1. Line 74: the focus of the study is "on the Baltic Proper and the Gulf of Riga". Could you please explain your choice of focus areas in more detail? While the focus on Baltic Proper as the largest and deepest basin is straightforward, why Gulf of Riga? It is the only sub-basin to the east of the Danish Straits for which there was no validation data available apart from some literature-based values, so it is not possible to adequately validate the model for this region, and consecutively the confidence in model performance here is lower than in other areas, both in terms of absolute biomasses of benthic fauna and its impact on pelagic biogeochemistry.
2. Figure 1: As the text often refers to different regions of the Baltic Sea, a legend for all of them should be provided in addition to numbers. Moreover, model results are presented

at intervals of 0-30, 30-70, 70-120 (and 120+) meters, so bathymetry on the figure should use the same gradation.

3. Section 2.2: BALTSEM simulates Baltic Sea as 13 horizontally homogeneous boxes. In the Results and/or Discussion section, it should be elaborated on how this affects model results. Some of the Baltic Sea sub-basins have strong gradients in nutrient distributions, which in turn leads to gradients in productivity and in distributions of benthic fauna. For instance, the eastern part of the Gulf of Finland is heavily influenced by riverine nutrient inputs and have higher primary production rates than its western part. How are these gradients accounted for?

4. Lines 128-130: why does the model consider respiration and excretion of benthic fauna to contribute *ammonia* and phosphate directly to the water column? At least in case of deposit feeders one might expect part of the excreted ammonia to be oxidised directly within sediments and to be released into the water column in form of nitrate, and some of the excreted phosphate to be bound within sediment, the same way as it is considered for microbially remineralised nutrients in the model?

5. Does the model consider sediment resuspension? It seems to be an important factor mediating organic matter availability in the sediments in coastal seas, especially shallow regions such as Gulf of Riga? Omitting it might have important consequences both for model parameterisation and its results?

6. Line 134: "degradation" - what type of degradation? At this point, it should be specified in more concrete terms.

7. Around line 152: how are silicate transformations handled in the benthos?

8. Line 167: is nitrification considered a sink for oxygen in the model?

9. Equation 8, page 7: is sequestered phosphorus considered a state variable in the model, as it is not listed among the state variables in Table A1?

10. Line 176: what is a definition of "severely hypoxic"?

11. Section 2.3: please provide more detail on model forcing and setup in addition to the references, so the interested reader does not need to look for those details within several previous papers. How was benthic fauna initialised?

12. Line 184-185. Aggregating the results to means and standard deviations obscures a lot of detail about model capabilities, which cannot be justified by stating that "the purpose of this study was to evaluate large-scale dynamics". Does the model capture seasonal dynamics of benthic fauna? Are there long-term trends in the model *or* in the data? There is a general lack of knowledge about benthic processes, so more detailed and varied comparison with data would allow to identify critical gaps and steer discussions which will help to identify directions for improvement. This is especially important for a new model implementation, as presented in the manuscript.

13. Lines 187-188. This is an important topic for benthic modelling and should be discussed in some more detail. What should observational scientists measure to help constrain the model?

Results

1. The model run is 1970-2020. Why was validation of pelagic state variables performed for 1970-2015?

2. Figure 8 shows that there was a noticeable impact of inclusion of benthic fauna/bioturbation on primary productivity and nitrogen fixation, so even if overall relative bias remained almost unchanged, there was an impact on pelagic dynamics. Please provide more detail on how the models with BMM and without BMM compare. Given the impacts of fauna and its activity on nutrient fluxes is the main scope of the paper, a more detailed assessment of changes in pelagic environment is justified.

3. Line 214: how is "reasonable accuracy" defined in case of benthic variables? As the

authors demonstrate, benthic fauna biomass shows high variability, hence large standard deviations, especially when aggregated over long time and large areas, while standard deviations in the modelled fauna are relatively small. Does this high std in data really justify using cost function (Appendix C) as a validation metric?

4. Line 221-224: see my comment above: why focus on the Gulf of Riga?

5. Section 3.3, from line 247: this is a key section in the manuscript, as it demonstrates the direct and indirect effects of fauna. Yet it is very short and just skims through the results and through the figures. It should be more detailed, which would also make it easier to relate figures 7-9 to the text. It would also benefit from detailed quantitative information, in particular regarding relative changes.

6. Line 264: for this, a figure similar to figure 5 could be provided.

7. Lines 265-271: figure 11 (as well as several other figures of the manuscript) contain multiple subfigures, so it would be very helpful to provide relevant pointers in the text, e.g., reference to Fig 11a on line 268, Fig 11b on line 269 etc, so the text and figures could complement each other.

Discussion

1. Line 284: both terms "long-term" and "large-scale" have been used multiple times in the manuscript without proper definition. It creates overall impression of vagueness. Does "long-term" stand for "multidecadal" or "long-term mean"? The words "large-scale" could be omitted altogether without impact on meaning.

2. Line 285: as it currently stands, it should be "Baltic Sea", not "coastal sea", as the manuscript shows the model is not yet applicable in high-salinity regions.

3. Line 288-289: to support the statement that benthic fauna can alleviate the 'vicious circle' of eutrophication, could you show the differences in extent of hypoxic area (e.g. with Figure 8) for simulations with different levels of bioturbation?

4. Similarly, what about primary production and N-fixation for the two future scenarios (Fig 10)? I think these would also support the discussion on faunal impacts on 'vicious circle'?

5. From line 307: as the overall model performance was not improved by adding benthic fauna, and related processes are not well understood, could the authors discuss on what understanding is currently lacking and which model improvements are desired.

6. Lines 313-316: same as the previous comment, which data is currently lacking? Which mechanisms are not sufficiently understood? The sentence on "the main strength of this study" is too general ("many interlinkages") – could it be replaced with something more concrete and relevant to the actual modelling work?

7. Lines 326-334: please keep referencing to figures and sub-figures as discussion requires.

8. Line 361: is it worth presenting fast running time as an advantage, since the manuscript presents results from only several simulations with sensitivity analysis limited to a single parameter?

Conclusions and outlook

1. Line 397: "much-studied bioturbation" - I suggest replacing this with "bioturbation, relatively more studied in the modelling context".

2. Could the authors provide some outlook on the potential future directions of benthic

modelling, in the context of Baltic Sea in particular?

3. Lines 399-402: These conclusions are far too general. Could it be something emerging from the study rather than vague statement that benthic-pelagic coupling is "modified by multiple drivers, which may change over time"?

4. Based on your work, has the time arrived for the regional models (of the Baltic Sea) to extend their formulations to explicitly include benthic fauna?

Appendix A, line 817: why is mortality rate chosen to be linear for *Limecola balthica*, and quadratic for the other two groups?

Technical corrections

1. Line 89: "extension" - should be "extent"
2. Line 110: "Ponotporeia" - should be "Pontoporeia"
3. In equation 1, should "Uci" be "UBFiC" in the denominator?
4. Page 6: equation 6 is missing.
5. Line 175 and equation 10: the fitting constants 5-8 are not featuring in the equation.
6. Figures 7 and 11: in both cases, "phosphorus (c, d)" should be "phosphorus (e, f)".
7. Line 319: "suggests" - should be "suggest"
8. Table 1: what is "total" biomass? Should it be "mean" instead?
9. Line 712: "and" - should probably be "an"
10. Line 742: Nitrite and nitrate should be "NO₂" and "NO₃", respectively.
11. Line 784: "and" at the end of the line should be "are"?
12. Line 794: "feces" - "faeces".
13. Line 907: "120-120" should be "70-120"
14. Figure C2: please use a different y-scale for the Bothnian Bay.