

Biogeosciences Discuss., author comment AC1  
<https://doi.org/10.5194/bg-2022-48-AC1>, 2022  
© Author(s) 2022. This work is distributed under  
the Creative Commons Attribution 4.0 License.



## Reply on RC1

Malcolm E. Scully et al.

---

Author comment on "Unprecedented Summer Hypoxia in Southern Cape Cod Bay: An Ecological Response to Regional Climate Change?" by Malcolm E. Scully et al., Biogeosciences Discuss., <https://doi.org/10.5194/bg-2022-48-AC1>, 2022

---

### Reviewer #1

General comment

Ongoing environmental changes such as warming and deoxygenation in the open and coastal oceans are altering biogeochemical cycles and ecosystems. However, a comprehensive understanding of the causes and consequences of deoxygenation in coastal areas is still lacking. The manuscript under review presents new results from a study on environmental changes recently observed in the ecosystem of the southern part of the Cape Cod Bay (CCB). The manuscript is well written and the main conclusion is justified by the presented data. (The answer to the question from the title is: yes it is.) However, I have some critical comments about the statistical analysis of the time-series data presented. Therefore, I can recommend publication only after minor revisions.

Specific comments:

- Methods: Wind speed data from two sources (Chapin Beach and NDBC buoy 44013) are presented at several places in the manuscript. But, unfortunately, I could not find any description of the wind speed data at all: How have they been measured? What was the time resolution of the measurements at Chapin Beach? How representative/comparable are the measurements from the two sites? The only reference for the source of the wind speed data from Chapin Beach ('courtesy of Weather Flow, Inc.') is given in the Figure 1 caption. So, add some text describing the wind speed data from Chapin Beach and the buoy 44013, please. And: is there no time-series available for the Chapin Beach site? (it would make more sense to analyze wind speed data which have been measured closer to the sites of the oceanographic measurements.)

***We only have very limited data from the Chapin Beach site (August–October 2019 and 2020). This data came from Weather Flow Inc. and is not publicly available. The data from NDBC buoy 44013 is highly consistent with the data from Chapin Beach data and is publicly available going back to 1984. Therefore, we will remove the Chapin Beach data from the manuscript and solely use the NDBC buoy 44013 data for consistency. We will include a brief description of the how the wind speed data is measured at buoy 44013 to the methods.***

- Methods: page 4, line 116: Add the details for the calibration of the DO sensor in the lab.

***The DO sensor is calibrated using a two-point calibration using water that has 100% saturation (obtained using aerating bubbler) and 0% saturation (obtained by sparging off all O<sub>2</sub> by continuous bubbling of nitrogen gas). Language will be added describing this process.***

- P8L195 and Figure 7d: I found it confusing that throughout the text 'winds from the north' and winds from the south' are mentioned and used to explain downwelling and upwelling, but in Figure 7d winds from SW and NE are shown. I do not see the point to 'change' the wind direction for Figure 7d.

***When analyzing the long-term wind data from NDBC buoy 44013, the strongest changes in summer winds are for winds from the SW and NE, so these are highlighted in Figure 7d. As discussed below, it is not clear if upwelling/downwelling in CCB is driven locally by across isobath winds (e.g., N-S) or remotely by the larger-scale response of the New England shelf to NE-SW winds. Moreover, we do not have enough in-situ data to determine what wind direction causes the maximal upwelling/downwelling response (N-S or NE-SW). However, we observe a downwelling response when the winds are generally from the north and an upwelling response when winds are generally from the south and this basic classification would not change if the reference angle for our winds was rotate +/- 45 degrees.***

- P8L196/197: Upwelling occurs when there are winds parallel to the coast, which won't be S or N in this case (please note: the hypoxia occur along the south coast of the CCB, not at the west coast of the CCB). The CCB seems to be a 'semi-enclosed embayment' open to the north. So, I wondering whether winds from the south may just lead to a decrease in the sea level (pushing the water away from the south coast). In turn, winds from the north just lead to an increase in the sea level (pushing water towards the south coast). I am wondering whether the authors should be more careful with their wording (i.e., down-/upwelling).

***There is evidence that the upwelling/downwelling response in a semi-enclosed embayment like Cape Cod Bay is driven primarily by the response along the open coastline to the north. Allen (1976) suggests that the upwelling/downwelling signal propagates as a coastally trapped wave, and thus may be driven by the larger scale response along the broader New England coastline. Over this region, the coastline is orientated roughly parallel to SW-NE winds, with SW winds driving upwelling and NE winds driving downwelling. Alternatively, there may be a local response in CCB where winds from the North-South are balanced by the sea-surface slope, which drives a geostrophic along-isobath interior flow. This along-isobath interior flow would drive bottom Ekman transport that results in upwelling for winds from South and downwelling for winds from North. Whether the upwelling-downwelling dynamics are driven locally or remotely is an open research question and beyond the scope of this paper. Both processes may be important. However, our data clearly demonstrate that the across isobath density field responds to winds from the North and South via downwelling and upwelling, respectively. Our intent in using the terms upwelling and downwelling is to convey the locally observed response of the density field and not represent the dynamics that are driving this response. We recognize this a little confusing and can alter the language to better reflect this intent.***

- Figure 6: *K. mikimotoi* appeared under 'normal' conditions in CCB already in 2017 and

2018. Is there an explanation why *K. mikimotoi* has appeared in CCB at all?

***We have no way knowing exactly how *K. mikimotoi* was introduced to CCB. There are a number of potential pathways including regional circulation and transport in ballast water. Its first appearance in the region was in summer 2017, and it was detected at locations as far north as Casco Bay in Maine and as far south as CCB. This nearly synchronous appearance over a very large region suggests transport by regional circulation, but we cannot say definitively. We are happy to add a little more language about the regional appearance of this species if that would be helpful. However, the key result is that blooms of significant magnitude have started to occur in recent years. Consistent with Smayda's model discussed on page 2 lines 54-57, we conclude that *K. mikimotoi* has been able to bloom in late summer in CCB because the physical environment has been significantly modified. This is one of the key results of the paper and we suggest that the emergence of *K. mikimotoi* blooms is strong evidence of significant habitat alteration. Our hypothesis is that this is either due to increased temperature or changes in wind. It is our view that 2017 and 2018 were not normal years. 2018 was the warmest summer on record and 2017 had the strongest NE winds during summer of any year. These very warm waters combined with the deep stratification that results from downwelling conditions are likely what allowed this new species to start blooming.***

- P13L296/297: This statement about the data presented in Figure 7b is misleading: There is only one year (i.e., 2018) with >40 days with temperatures higher than 20°C in the period 2018-2020. But there was indeed another year (i.e., 2016) with >40 days with temperatures >20°C before that period (i.e., between 1986-2017). The authors should be more careful with their wording here. Please rephrase.

***2018, 2019 and 2020 had 53.4, 40.3 and 40.0 days > 20°C, respectively. The only other year with > 40 days was 2016 (43.6). Perhaps this is not clear from the graph, so we can add a red dashed line at 40 days and modify the language, so this is clearer.***

- Figure 7c): I am wondering whether the significance (trend) of the linear regression is only driven by the data points from 2018-2020. Please check.

***Yes, if we exclude the 2019 and 2020 data, the regression is no longer significant. The slope is still negative, but there is sufficient scatter, and with relatively few data points (n=8), the slope is not significant without these two points (p>0.1). We will make sure this is clearly stated in the text.***

- Figure 7d): There are two points about the wind speed data which seem to be interesting but which are not discussed at all: (i) There is an obvious regime shift around 1995. Before 1995 the percent of the SW winds (NE winds) have been increasing (decreasing). After 1995 the percent of the SW winds show no trend with time anymore, but the NE winds show an increasing trend; and (ii) Between 2014-2020 the variability of the percent of SW winds was surprisingly low (compared to the rest of the data) which obviously corresponds very nicely with the astonishing increase in days with temp. >20°C (see Figure 7b) from 2015-2020. So, I think that there were shifts in the wind regimes in 1995/6 and 2014/5. The later one may have caused the appearance of *K. mikimotoi* in CCB in 2017.

***There may have been a regime shift in the % SW winds after 1993 (the slope of the regression for 1994-2020 is not significantly different than zero @ p<0.05). So, the significance of the overall slope appears to be driven by differences pre/post 1993. However, this is not true for NE summer winds or for surface***

**temperature. There is no statistical support for a regime shift around 2014/2015. In fact, both NE winds and surface temperature have increased relatively steadily over the period of record. The dramatic increase in the number of days > 20°C simply reflects this long-term increase in temperature. So, there is not statistical support to suggest that a regime shift has caused the appearance of *K. mikimotoi*. More likely either the steadily increasing temperatures and/or increasing summer NE winds have resulted in an environment that crossed some threshold allowing this new species to bloom.**

- Are there any water current data (i.e. ADCP data) available? It would be worth to see whether circulation patterns (-> ventilation, residence time of the bottom water) in the CCB have changed as well.

**There is no long term ADCP data available in CCB to our knowledge. There is ADCP data from the Gulf of Maine Ocean Observing System (GOMOOS) just southeast from Gloucester, MA (GOMOOS buoy A), but this is unlikely to capture the circulation within CCB.**

- Figure 7: It would make sense to calculate anomalies (i.e. the difference between the actual value and the average value from a reference period) for the data in a) and c). This will give a more robust idea about exceptionally warm or cold years (in a) or surface-bottom differences in c).

**We will change the presentation in figures 7a and 7c to show anomalies.**

- Nutrient concentrations can be significantly affected by other inputs such as rain and groundwater discharge (rain increases groundwater discharge) as well: Are there rain data (from the meteorological station at Chapin Bay?) and/or groundwater discharge data available?

**For the greater Gulf of Maine, Townsend (1998) estimated that advective inputs of nitrogen exceeded riverine and atmospheric inputs by nearly 20 times. While his budget did not include groundwater inputs, they are likely considerably smaller than riverine inputs. We expect similar dynamics in CCB with advective fluxes of nutrients from deeper offshore waters far exceeding atmospheric or groundwater inputs. Our data suggest that the intense blooms that are resulting in hypoxia in CCB are formed by a motile phytoplankton species utilizing the deep reservoir of nutrients below the thermocline. The drawdown of sub-pycnocline nutrients in 2019 and 2020 supports this basic hypothesis. Furthermore, these data also suggest that there were ample bottom nutrients in previous years when no blooms formed. We conclude that these intense blooms were not primarily controlled by variations in nutrient inputs, but by changes in the physical environment that allowed these deep nutrients to be utilized.**

- Would it make sense to add a brief outlook in order to speculate about the time point when CCB might becoming seasonally anoxic and the resulting consequences for fisheries and tourism?

**We are reluctant to speculate about whether or not CCB is going to become seasonally anoxic given the uncertainties in what is driving hypoxia. Our primary hypothesis is that intense blooms of *K. mikimotoi* provided the biomass that ultimately resulted in hypoxia in 2019 and 2020. While physical conditions are changing in CCB and such blooms could become more common, predicting when/if a specific species will bloom is very challenging. There is simply too much uncertainty to say with any confidence hypoxia is likely to occur moving forward.**

