

Clim. Past Discuss., author comment AC2
<https://doi.org/10.5194/cp-2020-163-AC2>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Reply on RC2

Gabriella M. Weiss et al.

Author comment on "Co-evolution of the terrestrial and aquatic ecosystem in the Holocene Baltic Sea" by Gabriella M. Weiss et al., Clim. Past Discuss.,
<https://doi.org/10.5194/cp-2020-163-AC2>, 2021

The article by Weiss et al. is well written as well as well illustrated and the record is based on a core from a particularly interesting region of the Baltic Sea for which new biogeochemical data is to be appreciated. I am not an expert on this kind of data myself, but think that this publication very well and precisely documents the used approaches, and I think that the results and the discussion are worth being published to broaden the view on the Holocene in the southern Baltic Sea, even though the relatively low temporal resolution of the record makes detailed comparison with other records difficult (see below). I also think that the age model could be explained in more detail since one has not only to visit the supplementary material of Weiss et al. 2020 but ideally also Warden et al. 2016 to get a precise idea how it was generated. Another aspect I see a little sceptical are the results concerning the terrestrial ecosystem structure since I think that with the used proxies, it can only be well estimated concerning a few certain plant types (such as *Sphagnum* indicated by the C₂₃ n-alkane). In case of several statements concerning the vegetation/climate development, I think the literature cited for comparison is only partly suited to support the results based on the biogeochemical proxies – probably better references for direct comparison can be found (see below).

We thank the reviewer for these positive comments. As mentioned in response to reviewer one we will include more detail on the age model as well as the vegetation changes in the region in the revised manuscript. See the answer to reviewer one: "The core was dated by correlating XRF data to two nearby cores described in Warden et al. (2016). The ¹⁴C ages were corrected using local marine reservoir values from Lougheed et al. (2013; doi:10.5194/cp-9-1015-2013). We will add the two Berglund et al. (2008) findings into our discussion of regional vegetation. These two pollen records show an increase in *Pinus* between 9.2 and 9.5 ka (Berglund et al. 2008a – doi: 10.1016/j.quaint.2007.09.018, and Berglund et al., 2008b – doi: 10.1007/s00334-007-0094-x) and revise the discussion to emphasize the dominance of woody species between ~11 and 3 ka." The dynamics of *Sphagnum* specifically can be difficult to determine in a mixed (sub)Arctic and temperate drainage basin as the dominant chain length of their *n*-alkanes has been reported to shift from C₂₃ in temperate environments to C₃₁ Scandinavian (sub)Arctic environments (see Vonk and Gustavsson, 2009).

Detailed remarks

Title and Abstract: I personally think that "structure" in the title implies more than the paper can provide in the end. Consider that in the abstract only a water source shift and a suggested vegetation diversification are mentioned concerning terrestrial ecosystem structure. General, the abstract could give less methodology and more own environment-related results.

We are sorry the reviewer feels this way, but we respectfully disagree. The abstract should reflect the work presented in the manuscript and this type of work is strongly based on analytical chemistry and advances therein. Hence the methodology in the abstract.

Sections 1 to 3.2

I have almost no remarks to sections 1 to 3.2, because they are generally very well written and are of appropriate length and focus. One remark only: In line 33 the expression "the SIS melted, exposed the land ..." sounds a little odd to me. I cannot say very much concerning the method sections since this is out of my expertise, but as far as I can tell this is also well done.

We will rephrase the sentence on line 33 in the revised manuscript.

Section 3.3

I am not sure if there is an inconsistency with the discussion here – maybe it should be mentioned here already that the C₂₈ 1,14-diol was present in high amounts during the MB phase as mentioned in section 4.4 - the expression "only present" does not imply high amounts in my opinion.

We will add that the 1,14-diols (including the C₂₈ 1,14-diol) were present in a relatively high amount in the MB phase to section 3.3 of the revised manuscript.

Line 199: Checking Kothhoff et al. 2017, it appears to me that the Yoldia Sea (YS) is not reflected in the record described there, a comparison should not start before ca. 8000 yr BP.

As mentioned in our response to reviewer one, "it is true that the record of Kothhoff et al. (2017) only goes back to 7.4 ka. As the only other record of temperatures reconstructed using LDI in the area, we felt it should be included. Our record has similar reconstructed temperatures to those presented in Kothhoff et al. (2017)." We will make it clear in the revised manuscript that the comparisons of the two records are from 7.4 ka and younger.

Section 4

Section 4.2 line 234 and following: it is implied here that the pollen records mentioned in line 236 are nearby, but I think the records used un Seppä and Birks 2001 are quite far away (some pollen records from northern Germany, Southern Sweden, Denmark or Poland would be nearer).

Section 4.2.2

First paragraph: Again, I am not sure if Seppä and Birks 2001 is well fitting here.

See previous answer and our response to reviewer 1.

In line 269 in the same paragraph it is stated that “the maximum extension of *Pinus* and *Juniperus* was recorded at 9.2 ka in these regional lakes”... Checking the cited literature, I can only partly agree: There are only minor increases of *Juniperus* pollen in the related time interval in Seppä et al. 2005/Digerfeldt 1977 and the *Pinus* peak is earlier. In Antonsson and Seppä 2007, there is a peak in *Pinus* percentages at 9.2 ka, but *Juniperus* percentages are significantly higher during the late Holocene and one could not speak of a maximum extension of this taxon around 9.2 ka. The sentence in line 270 and the following lines is correct concerning *Alnus*, but the pollen diagram of Lake Trehörningen depicted in Antonsson and Seppä 2007 does much more imply a decrease in *Pinus* (this one is very clear) than in *Juniperus* percentages (which does not seem to be consistently present between 11 and 5 ka anyway). I would think in this context that the attribution of the C isotope shifts to *Juniperus* shrub extension is not supported by the cited pollen records.

A time lag can be expected between the pollen deposited in lakes (usually from nearby catchment) and alkanes deposited in the Baltic Sea sediments (after being transported over relatively long distances). We are talking about each Baltic Sea phase independently, hence when we talk about the “maximum extension of *Juniperus*” we refer to the relatively higher peak for the specific phase (here the Ancylus lake). *Pinus* do not produce enough alkanes to have a real impact on the total alkane concentration found in the Baltic Sea (Diefendorf et al., 2011, 2015; Diefendorf and Freimuth, 2017), hence other gymnosperms such as *Juniperus* are likely the main driver of the carbon isotope composition of the *n*-alkane. As *Juniperus* produce a large amount of alkanes, even small shifts will have an impact, especially if the population of angiosperm trees such as *Alnus* are shifting as well. We do not claim that one type of vegetation is driving the carbon isotope signature of the alkanes but rather the relative increase/decrease of angiosperm versus gymnosperm vegetation. We will rephrase in the revised manuscript and add more comparison sites.

It would be really nice to have closer pollen records for comparison in which coeval *Juniperus* increases were present to support the interpretation concerning the C isotope shifts (e.g. the record shown by Yu et al. 2005 is quite near, but does not reveal such a signal), or if possible to see pollen data from core 64PE410-S7 itself.

Unfortunately, we do not have a pollen record for this core; however, we will add more comparison sites as noted above and in response to reviewer one.

Line 294: A citation for the 8.2 ka event would be good, particularly if one could be found for the research area.

We cite Fleitmann et al. (2008) for the multiple cold events recorded in the northern hemisphere. We will check for a more regional citation to add to the revised manuscript.

Conclusions:

Here, the 8.2 ka event is mentioned again (and marked in the figures, too) while it was said in 4.2.2 that no such event could be found in the own record and no citation was given in the discussions for such an event. If mentioned in the conclusions, this should be discussed in more detail and with a citation that there was indeed a cold event around that time in the research area. Generally, it would be good if conclusions based on own results would be better indicated.

The 8.2 ka event (as well as 9.2 ka and 10.2 ka) is marked in the figures as a reference point since it can be found in multiple northern hemisphere records.

Our conclusions are derived from the data presented in this paper and compared with previous studies.