



EGUsphere, referee comment RC1
<https://doi.org/10.5194/egusphere-2022-163-RC1>, 2022
© Author(s) 2022. This work is distributed under
the Creative Commons Attribution 4.0 License.

Comment on egusphere-2022-163

Anonymous Referee #1

Referee comment on "Constraining the particle-scale diversity of black carbon light absorption using a unified framework" by Payton Beeler and Rajan K. Chakrabarty, EGU sphere, <https://doi.org/10.5194/egusphere-2022-163-RC1>, 2022

The manuscript "**Constraining the particle-scale diversity of black carbon light absorption using a unified framework**" by Payton Beeler et al. reveals the effect of morphology of BC core on light absorption enhancement of BC due to the "lensing effect". The study used the Amsterdam Discrete Dipole Approximation (ADDA) method to calculate the particle absorption with varying BC core morphology and different coating compositions. Based on the results of the studies the authors found a factor (phase shift parameter) to describe the increase and decrease of absorption enhancements of BC caused by its morphology, and formulate universal scaling laws centered on the phase shift parameter. This study also provides physics-based insights regarding core-shell approximation overestimating BC light absorption. The presentation is concise and clear, and the topic fits well into the scope of the journal. The manuscript could be considered for publication with the following concerns being addressed.

- The novelty of the manuscript is not well presented. There have been a number of numerical studies on optical properties of BC with complex morphology using DDA and T-matrix method. Either the fractal aggregate model or coating scheme has been considered before.. It is recommended to explain the advance specifically for this work at least in the introduction section.
- In this study, ρ_{BC} (the phase shift parameter of BC core) shows the influence of BC core morphology on its light enhancement, but ρ_{BC} was determined not only by the morphology, but the size of BC can also influence ρ_{BC} according to formula (1). In 2.1

section, the authors state to calculate with BC core masses between 1 fg and 70fg, but the BC size calculation was missing in the results. In addition, previous fractal aggregate studies used the fractal dimension (D_f) to represent the morphology, what is the D_f for the freshly emitted, partially collapsed, and collapsed aggregate in this study?

- The absorption enhancement of BC core through the “lensing effect” was also investigated for light-absorbing coating materials like BrC, and the author notices that the total particle absorption is very sensitive to the image refractive index of the coating material. The increase of particles absorption with coating increase was a competition between the increase of BrC absorption and the decrease of the BC enhancement due to less light on the BC core. However, MAE_{BC} in this study shows the total absorption of the particle (e.g. in Fig. 4). It is recommend to subtract the absorption by the BrC shell in order to investigate the “lensing effect” of BC.

4 Section 2.1: The discussion about the influence of spherical monomer of BC aggregates on its optical properties is missing. Berry and Percival (1986) discussed that optical properties of fractal-like aggregates were determined by the primary spheres. In this study the primary sphere was chosen to be 20nm, Shetty et al., (2021) used 40nm. (<https://doi.org/10.1080/02786826.2021.1873909>).

- Section 2.2: The settings about the ADDA are not well described. The accuracy of ADDA depended on the size of the sub-volume compared to the wavelength of the incident light. What’s the resolution of dipoles per wave-length in this study?

Fig. 2: Y axis label is missing.