



EGUsphere, author comment AC2  
<https://doi.org/10.5194/egusphere-2022-335-AC2>, 2022  
© Author(s) 2022. This work is distributed under  
the Creative Commons Attribution 4.0 License.

## Reply on RC2

Ina Säumel et al.

---

Author comment on "Back to the future? Conservative grassland management can preserve soil health in the changing landscapes of Uruguay" by Ina Säumel et al., EGU Sphere, <https://doi.org/10.5194/egusphere-2022-335-AC2>, 2022

---

### Response Reviewer 2 (*Please find our replies in italic*)

A one-time survey across anthropogenic soils in Uruguay was carried out and analyzed for different soil parameters with the aim to identify interaction between land use and soil characteristics. The manuscript covers the relevant and timely issue of non-sustainable land use practices that support soil loss and degradation.

*Reply: We are happy that you see the relevance of the topic of our manuscript. Thank you for your time and willingness to review our manuscript.*

However, I am missing a substantial contribution to scientific progress in the methodology and/or results. There are long-term and recurring topsoil surveys established worldwide, results are being published in report formats on a regular basis, and in general, these data sets (especially when run over long time periods) hold valuable information for various scientific questions. However, in this manuscript, the authors did not communicate the aim, research question or hypothesis addressed with the study. A purely exploratory statistical analysis of a set of standard analyses of soil samples is not sufficiently novel or unique for a publication in SOIL. Therefore, I regret I cannot recommend the publication of this manuscript.

*Reply: We regret that you have the impression that our manuscript does not make a substantial contribution to scientific progress. Of course, we as authors and conductors of the research and the funders of our study have a different perspective. However, we take your critics seriously and highlight now better the novelty and contribution of our manuscript.*

*Long-term and recurring topsoil surveys that covers such a great variety of different land uses (different types of native forests, grassland, timber plantations, crops) are rare in the global south and especially in countries with limited resources. We need these surveys especially in those countries where land use change and overexploitation for the global market takes place right now. In Uruguay, these changes happen in remote areas with voiceless people and with governments that have no resources to control or reclaim*

*possible impacts. Independent data are crucial to discuss the local agendas and discourses of different stakeholders. We have already pushed the finger on this point please see line 62-71: "As soil degradation is extremely relevant for countries like Uruguay, which are socioeconomically dependent on their soils (Zubriggen et al. 2020), it is a topic of discussion for local farmers, academia, and the public. An actualization of the state of the art of soils and related processes is needed (García-Préchac et al. 2004; De Faccio et al. 2021), particularly as there has been little study of the impacts of the Uruguayan grassland intensification on soils properties (Beretta-Blanco et al., 2019). At the same time, while a new paradigm for grassland intensification with a wide set of means including fertilization has been proposed to increase economic and environmental sustainability (Jaurena et al., 2021), it is urgent to get more insights into the dynamics of nutrient in soils of Uruguay and their availability for crops (Beretta-Blanco et al., 2019)." And in line 77f. "We contribute to a better understanding of globally occurring degradation processes in the field of tension between desired soil productivity, yield limits, especially in erosion sensitive soils, and necessary soil conservation."*

*We reworked the part of the objectives of our study in the introduction as follows (line 81-91): "We therefore explored soil parameters describing current chemical conditions of topsoils that are parts of different soil groups and orders, and Uruguayan soil categories. Specifically, in order to explore the gains and losses of macro and micro-nutrients and soil organic carbon across landscapes and to determine the impact of land use change on acidification and trace metal mobility and related trade-offs with soil degradation and conservation. In detail we address the following question: i) how do fertility proxies such as soil organic carbon and content of nutrients, acidification (pH) and trace metals accumulation in topsoils vary across different land uses? Thus, we expand the knowledge across land uses from more natural to strongly modified uses and discuss the results in light of different degradation processes such as erosion, depletion of nutrients or carbon, acidification and accumulation of pollutants and in the light on current debates on intensification."*

*We also checked again if there have been publications on surveys that we might have overseen unintentionally before. Topsoil surveys that cover such a great variety of different land uses are missing. We explained this situation in line 62-71. "As soil degradation is extremely relevant for countries like Uruguay, which are socioeconomically dependent on their soils (Zubriggen et al. 2020), it is a topic of discussion for local farmers, academia, and the public. An actualization of the state of the art of soils and related processes is needed (García-Préchac et al. 2004; De Faccio et al. 2021), particularly as there has been little study of the impacts of the Uruguayan grassland intensification on soils properties (Beretta-Blanco et al., 2019). At the same time, while a new paradigm for grassland intensification with a wide set of means including fertilization has been proposed to increase economic and environmental sustainability (Jaurena et al., 2021), it is urgent to get more insights into the dynamics of nutrient in soils of Uruguay and their availability for crops (Beretta-Blanco et al., 2019)." Regarding existing surveys and the novelty of our data we already explained in line 310-317: "This is clearly reflected in the results of our topsoil survey, which also adds interesting insights from timber plantations, grasslands and native forests to an existing database consisting mainly of crops and pastures samples from 2002-2014, which demonstrated the loss of organic matter by 25% and an increasing loss of nutrients (Beretta-Blanco et al. 2019). We contribute deeper insights on fertility, acidification and trace metals accumulation in topsoils from a wide range of different land uses, which is, to our knowledge, unique for the region since the CONEAT classification (CONEAT Index, 1976)." Or in line 434ff: "To our knowledge, there has been no regional study of trace metals in the native riverine forests or timber plantations. Our work thus expands the evidence base for these land uses."*

*Concerning your second point of long-time data, we totally agree with you, that long term*

*data are very valuable and necessary, especially those that follows a standardized protocol. We call this always as the "gold of ecology". Regarding this we only want to state that the reality of funding and research projects looks different, research projects are often only funded for 2-3 years, in this time you can maybe establish a sampling design and do one or maximum two sampling runs and you have to publish the results immediately in high score journals to get in a new call funding for next projects. In addition, we had two years of funding without any possibility to go back to the field for sampling.*

Some points that I would like to share to reason my decision, and that might be helpful for future manuscript preparations, follow below.

*Reply: Thank you for your suggestions to improve our manuscript.*

I stumbled a bit over the title, being a question "sentence" without a verb. Titles, especially when they include questions, should be concise and meaningful.

*Reply: Thank you for this comment. Changed as follows: Back to the future? Conservative grassland management can preserve soil health in in the changed landscapes of Uruguay*

It is not clear in the abstract, what the goal of the study was (to "understand" soils), and what was done ("we analyzed ..."). From reading the abstract, I wondered whether this study is about a spatial and/or time series analysis. The results mentioned in the abstract give clues to sophisticated analysis methods: loss in nutrients, accumulation in topsoils on riverine forests, translocation, local land use trajectories. But later in the manuscript it appears that none of this was actually investigated directly.

*Reply: We see recognize that are abstract needs improvement and we changed the abstract to meet the comments also from reviewer 1: We analysed agri- and silvicultural intensification of Uruguayan grasslands in a country wide survey on fertility proxies, pH and trace metals in topsoils originating from different land uses. We are convinced that we did a sophisticated analysis of the data we have. We included the important points based on the discussion. Our discussion is based on our data and on evidence in literature.*

*Changed as follows: The 'soils of the anthropocene' are predominately agricultural. To understand them, we analysed agri- and silvicultural intensification of Uruguayan grasslands in a country wide survey on fertility proxies, pH and trace metals in topsoils originating from different land uses. We observed a loss of nutrients, trace metals and organic matter from grassland, crops and timber plantations. As an example, the cation exchange capacity was 160 percent higher in native forests compared to grasslands and lowest in timber plantations, reaching only half of the CEC in grasslands Acidification of topsoils continues as three fourth of all samples are 'extremely acidic' and 'very strongly acidic' and lowest in timber plantations. Topsoils of riverine forests accumulate more trace metals compared to the other uses. We assume an accumulation in the topsoils of riverine forests, where high levels of nutrients, trace metals and organic matter are found. The translocation of nutrients and organic matter across the landscape to the erosion base depends on local land use trajectories. Increasing soil acidification is driving a positive feedback loop, and land use intensification is leading to degradation of local black soils*

*within a few decades. Our data raises questions about the resilience and carrying capacity of Uruguayan soils with regard to currently implemented highly productive management forms, including the use of timber plantation for carbon sequestration, and supports more conservative forms of extensive management on the grassland biome.*

*This is also changed in the discussion section 4.1 as follows: Our data demonstrate the high amounts of organic carbon, nutrients and trace metals in topsoil samples from riverine forests, suggesting transport of soil particles from the surrounding land uses (e.g., grasslands, crop or timber plantations) to the borders of rivers, streams and creeks. Therefore, we assume that organic carbon, nutrients and trace metals are displaced within the landscape and accumulate in the floodplains. Regional soil erosion models estimate the loss of 2-5 tons ha<sup>-1</sup> year<sup>-1</sup> for a third of the country depending on precipitation, topography, soil erodibility and land management (Carrasco-Letelier and Beretta-Blanco 2017). One possible direct impact is the increasing eutrophication reported for larger local rivers, although the models used by these authors did not link Chlorophyll-a concentrations with agricultural land use (Beretta-Blanco and Carrasco-Letelier 2021 and replies). However, other vertical processes and differences across land uses can be hidden deeper in the soil profile, and have not been analysed in this study.*

*Regarding the trajectories of land use change we included more detailed information in the section Study area and design as follows: Line 129-143: "If the owner agreed, plot selection was stratified by different rural land use types: grassland, timber plantations of Pinus and Eucalyptus species, native forest, and crops. Native forests cover mainly riverine and park forests. The later are a savanna like transition zones between riverine forests and the open grasslands. We subdivided grassland plots according to the intensity of use: (i) undisturbed grassland (without grazing), (ii) partially grazed grasslands (with sporadic grazing and low animal charge), and (iii) highly grazed grassland (with high animal charge). Land use change from 1986 to 2017 follows basically three different trajectories: i) the expansion of timber plantations over grassland leading to a disaggregation of grassland by timber plantations; ii) cropland expansion where crop cover maintains the open landscape character of former grasslands, grassland conservation where large and regularly interconnected riverine forests in a landscape dominated by grasslands (Ramírez and Säumel 2021) and grassland intensification changing from natural grassland to so called 'improved' or artificial grasslands (Modernel et al. 2016; Jaurena et al. 2021). Fertilization and application of other agrochemicals is standard procedure in timber plantations, artificial grasslands and industrial crops."*

Some parts in the manuscript are rather misleading: In the methods it is mentioned that landowners were asked permission for a long-term monitoring of their land, but there is no further information found on this in the remaining text.

*Reply: Here we do not understand the point of the reviewer comment. We mentioned that asking the owner to get permission was import to install monitoring sites and take samples and take again samples in the future to get more data as 'gold of ecology' as discussed above. This shaped the location of the plots and sampling sites, thus we mentioned it in M&M, but there was no need to mention this again.*

The discussion actually starts with setting the background for a rationale to carry out the

survey, so it is somewhat out of place here and should be part of the introduction.

*Reply: As you suggested we highlight the need for our contribution now more pronounced in the introduction, please see also our comments and changes above.*

Some parts are misleading or hard to understand: e.g., the authors discuss the role of riverine forests, but results are shown for native forests. Sometimes, "park forests" are also mentioned without an explanation of that term.

*Reply: Thank you for these comments. We see this point and provide now more details on local ecosystems, land uses. In Uruguay the dominating native forest type are riverine forest, there are also in some parts hill forests. The park forests are a savana like transition form from riverine forest to grassland. The local discussion to which extent these park forest are natural or culturally shaped (e.g. as a result of grazing) is ongoing since decades without agreement. We added more information and references on the land use types in the methodology part as follows: (see Line 129-143: "If the owner agreed, plot selection was stratified by different rural land use types: grassland, timber plantations of Pinus and Eucalyptus species, native forest, and crops. Native forests cover mainly riverine and park forests. The later are a savanna like transition zones between riverine forests and the open grasslands. We subdivided grassland plots according to the intensity of use: (i) undisturbed grassland (without grazing), (ii) partially grazed grasslands (with sporadic grazing and low animal charge), and (iii) highly grazed grassland (with high animal charge). Land use change from 1986 to 2017 follows basically three different trajectories: i) the expansion of timber plantations over grassland leading to a disaggregation of grassland by timber plantations; ii) cropland expansion where crop cover maintains the open landscape character of former grasslands, grassland conservation where large and regularly interconnected riverine forests in a landscape dominated by grasslands (Ramírez and Säumel 2021) and grassland intensification changing from natural grassland to so called 'improved' or artificial grasslands (Modernel et al. 2016; Jaurena et al. 2021). Fertilization and application of other agrochemicals is standard procedure in timber plantations, artificial grasslands and industrial crops.").*

The CONEAT index or categories are nowhere explained, the mere numbers are meaningless to the reader not familiar with this classification.

*Reply: We provide now more details in the method section as follows: line 197ff.: "The CONEAT groups are defined by their productive capacity in terms of beef, sheep and wool expressed by an index relative to the average productive capacity of the country, to which the index 100 corresponds. The classification is based on photo-interpretation at a scale of 1:40,000, field verifications and physico-chemical analysis of the soils. The productivity indices correspond to soil groups. The CONEAT groups have been defined by the dominant and associated soils according to the Soil Classification of Uruguay. The groups are related to the units of the Soil Reconnaissance Chart of Uruguay at a scale of 1:1,000,000. For each group, some important soil properties and associated landscape characteristics are indicated. The nomenclature of the CONEAT groups correlates with the Soil Use and Management Zones of Uruguay. The Soil Groups are superimposed on the rural parcel and are represented in the CONEAT cartography at a scale of 1:20,000 (for more details see MGAP, 2020)."*

The discussion is mostly very vague and I cannot find new insights or sound conclusions

from the performed analysis results.

*Reply: We would appreciate very much if you come up with more concrete examples. We did our best to be as precise as possible and within our data and other evidence reported in literature. We changed parts of the discussion following the suggestions of reviewer 1.*

Finally, the statement of the last paragraph, that extensive management of native grassland would be better for soil fertility/health is nothing that is new or concluded from the presented study results. A final conclusion is missing.

*Reply: The paragraph 4.5 is a positioning within the debate on the future of grassland management based on the insights from our data. Grasslands are globally seen as cropland reserve e.g. to fight hunger (SDG2). Extensively used grasslands are often seen as unproductive and intensification strategies are discussed. Our data do not support intensification strategies.*

*Thank you for your suggestion and added now a new paragraph on conclusions as follows: "The land use intensification in Uruguay associated with increasing inputs of energy, nutrients and pesticides leads to an overall loss of soil fertility and increasing toxicity related to acidification, salinization and trace metal contaminants. Our data demonstrate the high amounts of organic carbon, nutrients and trace metals in topsoil samples from riverine forests, suggesting transport of soil particles from the surrounding grasslands, crop or timber plantations to the borders of rivers, streams and creeks. Of all the fertility proxies assessed, phosphorus in topsoil was most significantly affected by different land uses, being highest in native forests. Cation exchange capacity was also highest in native forests and lowest in timber plantations, where only half that of grasslands was measured. Our study highlights that soil acidification is ongoing and probably also mobilizing trace metals and their accumulation in riverine forest topsoils."*