



EGUsphere, author comment AC1
<https://doi.org/10.5194/egusphere-2022-36-AC1>, 2022
© Author(s) 2022. This work is distributed under
the Creative Commons Attribution 4.0 License.

Reply on RC1

Linghan Li et al.

Author comment on "Impact of atmospheric rivers on Arctic sea ice variations" by Linghan Li et al., EGU sphere, <https://doi.org/10.5194/egusphere-2022-36-AC1>, 2022

We thank the reviewer for detailed comments and insightful suggestions. This study has been greatly strengthened and improved by this review. We will revise the manuscript to address each of these comments. Below are our responses to each comment.

Comments on TC manuscript egusphere-2022-36:

This manuscript investigates the impact of atmospheric rivers (ARs) on partial sea ice concentration variation based on observations. The authors started with two cases in the Chukchi Sea in the summers of 2012 and 2020. The surface heat fluxes when ARs approach the ice cover are analyzed. Then, the authors expanded their analysis to the partial sea ice cover in the whole Arctic.

General comments:

The research is interesting and within the scope of TC. However, I do have some major concerns (Note that this manuscript is revisable). First, the authors state previous literature which addresses results similar to what they are presenting. For general readers, it is hard to see how this manuscript is a significant scientific advance in knowledge. I do think that there are some novel insights in this work, but the novelty should be discussed in a more explicit way. Second, the case study is somehow superficial. I do believe this study could be much in-depth. I have some specific suggestions which may be helpful to improve this study (see details below). Overall, a major revision is needed to make the manuscript publication worthy on The Cryosphere.

Specific comments:

- *As mentioned above, several previous studies, such as Woods et al. (2016), Hegyi et al. (2018), and others discussed in the Introduction, have reported that the extreme water vapor transport, including ARs, can lead to the Arctic sea ice melt through the surface fluxes (radiative and turbulent) based on case study and statistical correlation analysis. These conclusions and the analysis method are generally similar to the current study. Therefore, I think the authors need to identify how this manuscript is unique from the existing studies, especially what gap of knowledge this manuscript addresses. To my understanding, most previous studies about the role of moisture transport in sea ice melt focused on the wintertime and the Atlantic sector; while the cases studied in*

the current paper occurred in the summer the Chukchi Sea. To distinguish from existing studies, the authors may confine their analysis to the summer. The surface energy balance in the summer Arctic is very different from winter, so I think focusing on summer (mainly on the Pacific side, including the Chukchi Sea) can be regarded as a novelty. Another suggestion to polish the novelties can be seen in #3.

Novelties of this manuscript:

- This study covers all seasons for the entire Arctic Ocean as shown in the statistical analysis. The most important result is that moisture and wind components of ARs correlate well with partial sea ice concentration change on a daily timescale throughout the year almost everywhere in the Arctic Ocean. Though we only show two case studies during summertime in the Chukchi Sea of the Arctic Ocean, we also examined AR events during wintertime in the Bering Sea with similar effects on sea ice tendency (not included in this manuscript). Therefore, the homogeneous spatial pattern of correlation of AR conditions and sea ice changes and similarity across different seasons are important novelties of this study.

- ARs are one form of extreme moisture transport operating on weather timescales. Previous studies on moisture intrusion into the Arctic emphasize the important role of incoming longwave radiation. Our results show that turbulent heat fluxes are the dominant terms of surface energy balance; net longwave radiation is only moderate, suggesting that incoming longwave radiation is largely canceled out by outgoing longwave radiation.

- We explicitly separate the thermodynamic and dynamic effects of ARs on sea ice changes by partitioning moisture and wind components of ARs. Previous relevant studies mostly examine surface energy budget due to the moisture component of ARs. Our study includes the important roles of winds in driving sea ice motion, as ARs are also characterized by a low-level jet. Therefore, our study of ARs has a more comprehensive understanding of the physical processes of the interaction between ARs and sea ice.

- We develop a new method for detecting extreme moisture anomalies. Our method extracts high-frequency variations on weather timescales, removing seasonal cycles varying year to year. Previous methods remove a fixed seasonal cycle to extract anomalies, which include contributions of interannual variability and weather events. Our method is more suitable for studying extreme weather events such as ARs.

- *Analysis method:*

For the case study in Section 3.1, I suggest the authors present the daily anomalies (the departure from the daily climatology or the anomalies used in Section 3.2). Based on the analysis of original values of the surface energy fluxes, the authors argue that sensible heating is dominant. However, with the seasonal climatology in heat flux data, the conclusion might not hold. For example, given the climatological net longwave radiation in

summer is negative (downward positive), the weak positive values shown in the figures may indicate a large positive anomaly. Therefore, using the original values cannot tell us which process is the dominant one. The current figures showing the original values can be moved to the supplementary file if the authors want to keep them. In addition, the authors may show how large the magnitude of the anomaly is (for example, exceeding its 1.5/2 standard deviation or not). In this case, Section 3.2.1 and Fig.5 can be removed to save space.

We will include analysis of daily anomalies of surface heat flux terms for the two case studies in the revised manuscript. Because the definition of ARs is based on the original values of IVT, we start with presentation of original values to quantify the relative importance of each term in surface energy budget for the case studies. However, our statistical analysis over the entire Arctic for 40 years is based on anomalies.

- *The role of dynamical ice motion, i.e., ice drifting due to the southerly wind associated with ARs, needs further analysis. For a specific location, such as the small box the authors chose, I agree that the wind anomaly could contribute to the local changes in partial sea ice concentration. Considering that ice drifting is regarded as one of the key conclusions in this study, the authors should clearly show the amount of ice drifting rather than the southerly wind only. The ice drifting has been discussed in many cyclone studies but usually has been neglected in AR or extreme moisture transport studies (focusing on thermodynamical effects). Therefore, the analysis of ice drifting (if the evidence shows that it indeed matters) could be regarded as another highlight.*

Sea ice velocity is an important variable in sea ice balance. However, ERA5 does not provide sea ice velocity. We use northward wind as the driving force to approximately study ARs' dynamic effect on sea ice. The partial sea ice cover with ice concentration < 85% we consider is in free drift with small internal ice stress (Heorton et al., 2019). Sea ice motion in free drift is largely driven by wind stress. Sea ice velocity in free drift is consistent with wind forcing especially on weather timescales: ~2% in magnitude, ~30 degree of rotation in direction. Northward wind generally induces sea ice reduction due to divergent sea ice motion near the sea ice edge. Satellite observations and model outputs of sea ice velocity will be examined in future study.

Technical

- *Rank correlation: I might miss some key information here, however, the authors may state how to rank the data. Does the result sensitive to the ranking?*

Spearman's rank correlation is the Pearson correlation between ranks of variables, considering monotonic relationships between variables. Rank correlation is nonparametric and is robust to extreme values.

- *Some figures should be refined. For example, it is hard for readers to identify the digits in Fig.1b&3b.*

We will improve the figures in the revised manuscript.

- *Figures 1a & 3a: The authors may clearly state how they define the SIC change during Aug 4-6, 2012.*

Sea ice concentration change during Aug 4-6, 2012 equals to sea ice concentration on Aug 6, 2012 minus sea ice concentration on Aug 4, 2012 (Figure 1a). The figure caption will be revised.

- *Line 174: the details of the citation are needed. Is it obtained from an NSIDC webpage?*

This will be changed to be 'We study another extreme event in summer of 2020. 2020 experienced the second lowest summer sea ice extent and the lowest sea ice extent during spring, early summer, and fall in the Arctic, based on NSIDC sea ice index (Fetterer et al., 2017).'

- *Line 208-209: More evidence is needed to support the argument "moisture content is more important than wind speed".*

This is to be changed to 'For this AR event on July 27, 2020, peaking moisture content, along with high wind speed, generates peaking downward turbulent heat fluxes.'

- *Line 267-268: 15% is conventionally regarded as the edge of sea ice cover. I'm just wondering why the authors chose 85% as the upper limit. Is it an empirical choice? If so, the authors may remind the readers here and state that the conclusion is not sensitive to the choice after test.*

We consider partial sea ice cover with ice concentration between 15% and 85%. This is in general consistent with the definition of the marginal ice zone having ice concentration between 15% and 80%. Also, sea ice is in free drift with small internal ice stress for ice concentration less than 85% (Heorton et al., 2019).

- *Line 290-291: I'm wondering about the persistence of these events (say the continuous days exceed 1.5 or 2 sigma or 90% percentile). For example, if the mean persistence is 5-day, can we infer that the mean SIC melt per event is -5% at a given location? It would be great if the authors can add a figure to present the timescale of these events.*

Duration of AR events is another important consideration. The case studies show that those two major AR events last less than 1 day (18 hours and 16 hours), based on hourly time series (Figure 2 and Figure 4). In the statistical analysis, we use daily extreme moisture anomalies, consistent with daily satellite observations of sea ice concentration. The regular and consistent time intervals are necessary for calculating correlations. Responding to the specific question the reviewer asks, the mean ice concentration tendency anomalies with daily extreme moisture anomalies are the same as the mean ice concentration tendency anomalies with one AR event lasting for several days, because ice concentration tendency is ice concentration change divided by duration. Duration and other characteristics (e.g. frequency and intensity) of ARs in the Arctic contrasting with ARs at midlatitudes is a very interesting topic for future study.

- *Line 295, 308-310: Most of the content of Section 3.2 is based on extreme moisture events. I fully understand that ARs and the 90% percentile moisture extremes share similarities and overlaps. Since the topic of this manuscript is AR, showing the analysis using AR (like Appendix Fig.3) would be more consistent, right?*

We develop a new method to extract high-frequency variations on weather timescales and to identify extreme moisture anomalies as approximate ARs which are validated by Guan and Waliser's AR catalog version 3. Our method of detecting daily extreme moisture anomalies is simple and efficient for large datasets from ERA5 and agrees well with the AR catalog from 6-hourly MERRA2. As far as we know, there is no AR catalog based on hourly ERA5, and detecting ARs from ERA5 using the full criteria from Guan and Waliser could be another project.

- *Line 322-324: There are at least two effects of southerly wind associated with ARs: dynamically redistributing the ice fraction and transporting the water vapor into the Arctic. Thus, the authors may add "In addition to delivering water vapor into the Arctic" or similar words somewhere.*

Winds associated with ARs have at least three effects on sea ice: 1. Northward winds drive divergent sea ice motion near the sea ice edge to cause sea ice to decrease dynamically. 2. High wind speed enhances turbulent heat fluxes. 3. Northward winds transport water vapor and heat from lower latitudes into the Arctic. These three effects are mentioned in the results. Additionally, ocean conditions could respond to high winds related to ARs (such as enhanced ocean mixing) inducing further sea ice melt.