



EGUsphere, author comment AC2
<https://doi.org/10.5194/egusphere-2022-705-AC2>, 2022
© Author(s) 2022. This work is distributed under
the Creative Commons Attribution 4.0 License.

Reply on RC1

Melissa Sophia Schwab et al.

Author comment on "Environmental and hydrologic controls on sediment and organic carbon export from a subalpine catchment: insights from a time series" by Melissa Sophia Schwab et al., EGU sphere, <https://doi.org/10.5194/egusphere-2022-705-AC2>, 2022

We appreciate the encouraging comments and wish to thank the reviewer for the effort and time spent reviewing the manuscript.

Current literature provides several accepted protocols for handling and storing dissolved organic carbon. Storage and preservation methods include a variety of containers (borosilicate vs HDPE), biocides (e.g., HgCl₂, NaN₃, HCl, HNO₃), and storage temperatures (refrigerated vs frozen). The commonly recommended methods are frozen storage (Walker et al., 2017; Heinz and Zak, 2018) and the storage of acidified samples in cold temperatures (4° C) (Cook et al., 2016; Nachimuthu et al., 2020). However, both options are characterized by advantages and disadvantages. Thieme et al. (2016) and Walker et al. (2017) demonstrate that freezing might preserve dissolved organic carbon concentrations and isotopic compositions while Spencer et al. (2007) and Thieme et al. (2016) argue that freeze/thaw cycles likely affect chemical and optical compositions of dissolved organic matter. Opinions further differ regarding the pre-treatment of dissolved organic carbon for freshwater and marine water samples. Regardless of the preservation method, organic carbon will be subjected to decomposition and alteration with increasing storage time.

Due to the low concentrations of dissolved organic carbon concentrations in the Sihl River, 20 mL of sample material were concentrated in precombusted gas-tight 12 mL exetainer vials by repeated freeze-drying of 5 mL aliquots. The vials were stored frozen until further analysis. We have failed to include this preparation step in the methods and have revised section 2.3.

References

Cook S., Peacock M., Evans C. D., Page S. E., Whelan M., Gauci V. and Khoon K. L. (2016) Cold storage as a method for the long-term preservation of tropical dissolved organic carbon (DOC). *Mires and Peat* **18**.

Heinz M. and Zak D. (2018) Storage effects on quantity and composition of dissolved organic carbon and nitrogen of lake water, leaf leachate and peat soil water. *Water Res* **130**, 98–104.

Nachimuthu G., Watkins M. D., Hulugalle N. and Finlay L. A. (2020) Storage and initial processing of water samples for organic carbon analysis in runoff. *MethodsX* **7**.

Spencer R. G. M., Bolton L. and Baker A. (2007) Freeze/thaw and pH effects on freshwater dissolved organic matter fluorescence and absorbance properties from a number of UK locations. *Water Res* **41**, 2941–2950.

Thieme L., Graeber D., Kaupenjohann M. and Siemens J. (2016) Fast-freezing with liquid nitrogen preserves bulk dissolved organic matter concentrations, but not its composition. *Biogeosciences* **13**, 4697–4705.

Walker B. D., Griffin S. and Druffel E. R. M. (2017) Effect of Acidified Versus Frozen Storage on Marine Dissolved Organic Carbon Concentration and Isotopic Composition. In *Radiocarbon* Cambridge University Press. pp. 843–857.