



EGUsphere, author comment AC2  
<https://doi.org/10.5194/egusphere-2022-766-AC2>, 2023  
© Author(s) 2023. This work is distributed under  
the Creative Commons Attribution 4.0 License.

## Reply on RC2

Arundhuti Banerjee et al.

---

Author comment on "On Parameter Bias in Earthquake Sequence Models using Data Assimilation" by Arundhuti Banerjee et al., EGU sphere,  
<https://doi.org/10.5194/egusphere-2022-766-AC2>, 2023

---

### Response to the editorial comments for the Manuscript entitled "**On Parameter Bias in Earthquake Sequence Models using Data Assimilation**".

The authors are thankful to the editor for his insightful guidance in making us improve the quality of the manuscript and for considering us for possible publication in the esteemed journal of 'Nonlinear Processes in Geophysics'. The authors are thankful to the reviewers for their helpful comments. Following the comments of the reviewers, the necessary changes are made in the revised manuscript.

#### Comments from Reviewer 2

Comment 1:

The study uses particle filter as the data assimilation method to solve the problem. But it would be good to inform the reader which other data assimilation methods were previously used in studies related to earthquake modelling and if those were successful. If not, which were the main issues and why a particle filter would suit better in this problem compared to those. This would mainly situate the reader on the importance of your choice on the method for this study.

Response:

Several different data assimilation methods have been used previously for estimating states for earthquake models. It has been reported in the manuscript (line 25-27, page 2). However, the data assimilation methods used in these references have not been explicitly mentioned in the manuscript. This has been modified from line 32-43 in Page 2 in the text.

Comment 2:

Equations 4 and 5: What does  $j$  mean?

Response:

The authors are thankful to the reviewer for the comment.  $J$  is a typo, and it should be  $j$  which is the number of particles or realization. It has been corrected in the manuscript.

Comment 3:

The authors use a Lorentz function instead of a Gaussian to prevent filter degeneracy. In addition, a SIR step is used to further avoid this issue. Were these enough to avoid filter degeneracy or the system still presents the problem?

Response:

The authors are thankful to the reviewer for the comment. Yes, the system did not face any filter degeneracy issues. We introduce a resampling step known as sequential importance resampling (SIR). This resampling discards particles with very low weights, while duplicating particles with high weights. Additionally, we introduced jitter in the prior distribution to avoid degeneracy.

Comment 4:

The authors mention that real earthquakes are far from being periodic but, as they have considered a 0D model, their system generates periodic cycles. I wonder how far from a real state this 0D model is and why this study has not used a 1D model, in which at least a minimum spatial dimension would be considered. It will be interesting to address in the manuscript why the 0D model was chosen in this case.

Response:

The authors are thankful to the reviewer for the comment. We have addressed this comment by adding a paragraph on the use of 0D, 1D and higher models for studying earthquake cycles. It has been mentioned in the discussion (line 357-366 in page 21 in text).

Comment 5:

I suggest different colors for the trajectories in the phase diagram in Figure 2, as it is hard to distinguish between them.

Response:

This comment has been incorporated and the color has been changed in the figure.

Comment 6:

4 time steps in this model correspond to which portion of a seismic event? Please, describe it in the manuscript to situate the reader on the frequency of the assimilation steps, as earthquake cycles may not be a subject well understood by many.

Response:

Having large assimilation steps can also have a detrimental effect on the data assimilation process as it can miss characteristic variations of the earthquake cycle. A parameter bias can have a substantial effect on the evolution of the state variables, which may be difficult to correct if the assimilation step is large. Hence a short time step is to be chosen that allows the assimilation to capture the important characteristics of the earthquake cycle. An additional text has been added in the manuscript (line 207-210, page 9) to educate the readers behind selecting this time step.

Comment 7:

Have you tested the impact of the use of less particles in the filter? If so, it would be good to share these results as well.

Response:

The authors are thankful to the reviewer for the comment. Yes, the authors have used 50 and 100 particles for this study. However, using a smaller number of particles, the system faces filter degeneracy issues. The authors added a sentence to describe these results. We have addressed this comment by adding text in the manuscript (line 340-347, page 21).

Comment 8:

How do the orders of magnitude of the observation errors compare to the states? Are those the typical magnitude of the real measurement errors?

Response:

The observational errors of the synthetic observations used in the study have been mentioned in the text (line 215-216, page 10). To obtain error estimates for each measurement type, the method explained in the study by Van Dinter (2019) were used where latest state-of-the-art values from the literature were considered. These errors were then downscaled to our model setup using the analogue scaling relation developed in Corbi et al.2013 (Van Dinter (2019)).

Comment 9:

It would be good to explain in the label of Figure 5 what each of the lines in the pdf represent.

Response:

The red line represents the posterior pdf, the grey line represents the prior pdf, the magenta line shows where the observation stands, and the black line is the true state. This explanation has been added in the manuscript.

Comment 10:

The authors mention that by using different assimilation settings, it is possible to inflate the ensemble. But it seems that the ensemble spread is not exactly the problem, as the state-parameter estimation presents much better results than the other tests which have an improved ensemble spread. Can the authors explain more clearly the effects of the spread on this specific seismic system?

Response:

The authors are thankful to the reviewer for the comment. In the case of state parameter estimation, the prior distribution contains particles with different parameter values (covering the entire spectrum from  $e = 0.1$  to  $0.8$ ) as shown in Figure 9 (a-e). Hence for the data assimilation to be effective there is no need to inflate the ensemble spread to cover the (observation or) the true state. As seen from Fig 9, the range of the prior encompasses the shear stress value of the observation. On the other hand, in state estimation, the parameter  $e$  is constant, which would require inflation of the ensemble to encompass the value of the observation.

Comment 11:

Figure 8b): What happens after nearly every 250 time steps, in which the periodic behaviour is lost by a double peak? What is the influence of the double resampling in these patterns?

Response:

In Figure 8b, which represents the double resampling experiment, we observe a double peak at 250 time steps. The data assimilation analysis does not fit the observations well. A similar mismatch is observed after approximately 500 time steps in this experiment. We thank the reviewer for pointing this out. At these moments, the double resampling effectively increases the spread in the particles to such an extent, that the constraint to the shear stress observations becomes less strong. The double resampling is not as effective in increasing the ensemble spread as the increased model error is (Fig 8a). We added a sentence to the text to describe this. We have addressed this comment by adding text in the manuscript (line 269-274, page 17).

Comment 12:

Still on Figure 8, it seems that Fig 8a) presents better results than Fig 8c), which makes me wonder if the double resampling is really helping the system.

Response:

Indeed, the increased model error in Fig 8a appears to be more effective than the double resampling. We changed the text to reflect this and thank the reviewer for this observation. We have addressed this comment by adding text (line 274-279, page 17) in the manuscript.

Comment 13:

The results for the state-parameter estimation are indeed promising and I congratulate the authors for this, but I would expect a comparison of these results with any other study (if they exist) using 1D models with or without data assimilation. The manuscript lacks information on other results found by studies which used other data assimilation methods and/or models with different dimensions.

Response

Data assimilation in earthquake cycle models is still in its infancy. We are not aware of other studies that considers a model similar to the earthquake cycle model that we have used in this study. In the discussion, we have added a section which talks about the use of simplified and higher dimensional models for studying earthquakes. (line 352-362).

References:

Corbi, F., Funicello, F., Moroni, M., Van Dinther, Y., Mai, P. M., Dalguer, L. A., & Faccenna, C. (2013). The seismic cycle at subduction thrusts: 1. Insights from laboratory models. *Journal of Geophysical Research: Solid Earth*, 118(4), 1483-1501.

van Dinther, Y., Künsch, H. R., & Fichtner, A. (2019). Ensemble data assimilation for earthquake sequences: probabilistic estimation and forecasting of fault stresses. *Geophysical Journal International*, 217(3), 1453-1478.

Please also note the supplement to this comment:

<https://egusphere.copernicus.org/preprints/2022/egusphere-2022-766/egusphere-2022-766-AC2-supplement.pdf>