

## ***Interactive comment on “COSMOS-UK: National soil moisture and hydrometeorology data for empowering UK environmental science” by Hollie M. Cooper et al.***

**Hollie M. Cooper et al.**

holcoo@ceh.ac.uk

Received and published: 21 January 2021

The authors thank both referees for providing detailed and helpful comments, suggestions and questions. We will add detail and clarity to the manuscript where suggested.

With regard to comments concerning the inclusion of individual calculations and more detailed explanations of variable derivation and methodology, we believe the manuscript describes the data set at an appropriate level of detail for the intended end users who are interested in soil moisture and hydrometeorology measurements. This description paper is not aimed at researchers looking to further develop the cosmic-ray method of soil moisture measurement.

Printer-friendly version

Discussion paper



The COSMOS-UK team is in active discussion with researchers across the European COSMOS community, with the intention of producing a more detailed output to describe specific methodologies for deriving soil moisture from the CRNS.

We address each of your individual comments (RC) with our response (AR) below.

Response to Referee 1

RC1 1: (Title) "empowering UK environmental science" To me this formulation reads a bit "selfish". It may be UK-national soil moisture data, but it should be (and I believe it is) open to anyone who works in the field of environmental science, British or not. I think there are many global or continental models that include the UK and would likewise benefit from your data. So I suggest to change the title into something less "nationalistic".

AR1 We had no intention for the title to imply any such meaning and we will therefore carefully amend it. The revised title will read, "COSMOS-UK: National soil moisture and hydrometeorology data for environmental science research".

RC2 35: Could you find another citation Moene and van Dam? I think there are plenty of papers out there that give a more specific introduction to the topic of soil moisture in the Earth System, e.g., by Seneviratne et al.

AR2 Thank you for your suggestion, we will reference the work by Seneviratne et al. (2010).

RC3 40: "measurement footprint" You may already specify it here to let the reader know what to expect. E.g. "the field scale measurement footprint"

AR3 We will amend the text to "field scale measurement footprint".

RC4 57: 80cm -> blank missing

AR4 This will be corrected.

[Printer-friendly version](#)

[Discussion paper](#)



RC5 80-85: What's the reason for not having more stations in Scotland (only 2 stations in the east)? Also for Yorkshire and North-West England the network is more sparse.

AR5 Installation of COSMOS-UK sites was initially focussed in areas where soil moisture was expected to have greater variability and where practical and logistical challenges related to access and communication were minimised. Installation of sites in less-represented regions is in consideration but is dependent on the availability of resources. This will be made clearer in the manuscript.

RC6 Table 1: I think the table needs to be condensed. Since available in the metadata, you can skip the 2 columns for Easting and Northing and also End date can be moved to Start date like this: "Start(end) date" 26/11/2013(-01/10/2016) The numbers for SAAR and Altitude should be right-adjusted. For the Soil type and land cover you should define abbreviations in the table header (e.g. MS for Mineral soil, IG for Improved grassland). and Soil type should go right of Altitude and SAAR. So finally one line could be as short as: Cochno 23/08/2017 168 662 MS IG and all stations could be overviewed at a glance on a single page.

AR6 Thank you for this suggestion for Table 1. We agree that this will greatly improve the table and we will amend it as suggested.

RC7 120-122: not clear if all CRS1000B sensors have been removed from the network and CRS2000Bs are used now everywhere. If all had been changed, since when is the network pure CRS 2000B? Pls. be more specific.

AR7 The text will be revised for clarity. Only the first four sites were installed with a CRS1000B (Chimney Meadows, Sheepdrove, Waddesdon and Wytham Woods), whilst all other sites were installed with a CRS2000B sensor. Wytham Woods was decommissioned in October 2016. In February 2020, CRS2000B sensors were installed at Chimney Meadows, Sheepdrove and Waddesdon alongside the existing CRS1000B sensors which remain connected.

[Printer-friendly version](#)[Discussion paper](#)

RC8 125: Shortly explain what TDT sensors are.

AR8 The following detail will be added, “Each site includes point scale soil moisture sensors, which estimate VWC via Time Domain Transmissometry (TDT). These TDT sensors estimate point scale soil moisture by measuring the time taken for an electromagnetic wave to travel along the sensor’s closed circuit; this transmission decreases in speed with soil permittivity (Blonquist et al., 2005).”

RC9 127: What was the reason to remove the PICO profiles?

AR9 We experienced issues with this sensor as configured at our sites. Specifically the configuration resulted in a high failure rate for other sensors and resulted in loss of data. Because of this, these instruments were removed during 2019 - 2020 in order to maximise the data capture across the array of instruments. This will be made clearer in the manuscript.

RC10 139: Why don’t you use wind shields for the Pluvios? It’s so often windy in Britain.

AR10 Pluvio wind protection shields were omitted during COSMOS-UK site installations as site locations were identified as being not particularly exposed. It is acknowledged that this decision could impact the extent of wind-induced under-catch, and shields will be considered during future network improvements. This will be made clear in the manuscript.

RC11 157: Pls. list the 8 sites with extra snow measurements

AR11 We will amend the text to read, “These sites (Glensaugh, Easter Bush, Gisburn Forest, Plynlimon, Sourhope, Moor House, Cwm Garw and Cochno) were installed with...”.

RC12 169: Is there any publication that shows the superiority of the SBS500, so that you want to use it as reference to check the quality of others (like the Pluvio)?

[Printer-friendly version](#)[Discussion paper](#)

AR12 We will amend the text to include relevant citations. The design of the EML SBS500 has improved aerodynamic characteristics (Strangeways, 2004), and Colli et al. (2018) report that when compared with other assessed rain gauges, the SBS500 exhibits superior reduction in turbulence and under catch.

RC13 183-184: Why do you only consider 0-25 cm depth for calibration? Later in figure 7 you suggest much larger penetration depths.

AR13 Soil sampling depths were selected to match typical (moist) UK conditions. Ideally, subject to cost, calibrations should be repeated at different VWCs, preferably capturing the full range of VWC measured at a particular site. We will add clarity to the text and highlight that the shallow soil layers are given higher weighting than deeper layers to represent the decreasing contribution of deeper soil water (Köhli et al., 2015; Schrön et al., 2017). Figure 7 will be amended; this currently shows incorrect (higher) values for D86. We apologise for this error and we will correct the data set accordingly.

RC14 183: Pls. provide model and manufacturer for the soil augers

AR14 We will include these details.

RC15 189: Fig.3: You should also provide annuluses (25,75, 200m) for the older calibrations (e.g. by having 2 separate graphs side by side with one legend, or by additional annotation in brackets '5 (25 m')

AR15 We will amend Figure 3 to include past calibration distances.

RC16 205: Were the bulk densities also obtained with the weighting function?

AR16 We do not currently obtain the bulk densities with the weighting function, however we will consider this in future work. This will be made clear in the manuscript.

RC17 206: Did you also consider the findings of Schroen et al. 2017 <https://hess.copernicus.org/articles/21/5009/2017/> for your weighting and calibration approach? The Sheepdrove data of COSMOS-UK was used in this work.

[Printer-friendly version](#)[Discussion paper](#)

AR17 Yes these findings have been considered in our approach; when applying the soil sampling calibration, the shallow 5 cm layers are given a higher weighting than the deeper ones. Please note that the Sheepdrove COSMOS-UK site is unrelated and in a different location to the Sheepdrove monitoring station referred to by Schrön et al. (2017). We will make this clearer in the manuscript.

RC18 208-210: There should be a discussion about the magnitude of possible errors in the determination of the reference soil moisture. Is the 0.03 vol.% difference significant or may the error in sampling and thermo-gravimetry be even higher?

AR18 This is an interesting question which we are unable to answer fully at this time, however we will add detail and address this as part of future recalibrations.

RC19 211: Tab. 5: Please give standard deviations for bulk density, lattice water, and soil organic carbon (add with  $\pm$  sign)

AR19 We agree these are needed and we will add these to Table 5.

RC20 217-230: Please be more specific in the description of the derivation of the corrected neutron count signal. This important step should be reproducible by the data users. So you should provide the exact formulas and specify the constants used (e.g. for the reference pressure). Did you use the same reference pressure for all stations? Did you do or do you plan to do a cross-calibration for the different sites, to get an idea of the variations in sensitivity of the sensors?

AR20 After consideration we believe that the current description with relevant citations provides an acceptable level of detail for the data set's intended users, who are interested in soil moisture and other environmental data. We will add clarity regarding reference pressure; a value of 1000 hPa is used for all sites, whilst the instantaneous barometric attenuation length is calculated for each site. A selection of CRS2000/B sensors deployed at COSMOS-UK sites were cross-compared for sensitivity under lab conditions prior to installation. Where necessary, in situ sensitivity comparisons have

[Printer-friendly version](#)[Discussion paper](#)

been completed by collecting several months of data from adjacent sensors.

RC21 224: I think it should be "Physikalisches Institut's, University of Bern, Jungfrau-joch"

AR21 Thank you, we will correct this text.

RC22 227: What are the implications of using a neutron monitor with such a large displacement and elevation difference. I assume a big difference in the cutoff rigidity between Jungfrau-joch and your sites. Wouldn't the Kiel monitor be better suited? I think that this should also be discussed in the paper.

AR22 We will add detail and discussion to the manuscript. We use normalised count rates in the intensity correction, which are not greatly affected by their cut-off rigidity in the absence of significant space weather events. During such events, the cut-off rigidity of a specific location may change due to magnetic field disturbances, so only matching cut-off rigidity might not compensate for such events. When comparing Jungfrau-joch neutron monitor with the available monitors of similar cut-off rigidities to the COSMOS-UK sites, the normalised counts and the associated trends were in good agreement. When choosing the most suitable neutron monitor for this work, Jungfrau-joch was identified as a well-maintained monitor with a high level of data completeness, however we will continue to monitor research in this area to improve our methods where possible. The Kiel monitor is currently described as not having efficiency-corrected values available, so we will reconsider this monitor when these data are available.

RC23 241: Are the fits for all the sites performing equally well? You may discuss and add some other cases too.

AR23 There may be some misunderstanding with this question. This figure does not illustrate goodness of fit, it only illustrates the portion of the calibration curve corresponding to the observed count rates. We will make this clearer in the manuscript and add some other cases.

[Printer-friendly version](#)[Discussion paper](#)

RC24 243: Fig. 4: Please add the formula for the synthetic curve including constants to the legend of the plot

AR24 The formula will be included.

RC25 264: Fig. 5: Please add panel IDs (a, b, c). What is the reason for the counts in a) being half of those in b)? Is it a different aggregation interval? X-axes text (date) should be removed for all but the lowermost panel. Then the margins between panels can be removed in favor of larger legend font sizes. Y-axis text for e) Precipitation is too small (smaller than for other panels). Legends (font sizes) for d) VWC are too small.

AR25 Thank you for your suggestions and spotting an error in the plot. Panel IDs a, b, c will be added to Figure 5. The data in panel a will be replaced with the intended aggregation interval. Your other suggestions for improvement will also be included.

RC26 271: Fig. 6: Please add panel IDs. Resolution of figure needs to be improved. Since colors are rather hard to distinguish (especially with red-green blindness), I would recommend to use gray-scale (maybe with transparency) for the lines and symbols (point, cross, dot, ...) for the regions at the highlighted date.

AR26 We will add panel IDs to Figure 6, improve the resolution and make the different groups more distinguishable for all users.

RC27 289: "measurements" I wouldn't see this as a measurement but rather as a derived quantity. Something of "shows the assumed/computed/estimated D86..."

AR27 We will change the text to read, "...shows the estimated D86 values for a...".

RC28 293: Fig. 7: Add panel IDs. All texts are by far too small. As for Fig. 5 I suggest to remove the x-axes annotation and to cut the margin between the sub-panels. How can the Euston VWC time-series reach 0? Was that reflected by the TDT sensors?

AR28 Figure 7 will be amended as suggested. The Euston site is located on very well-draining sandy soil and during the extreme dry event in 2018 the derived soil moisture

Printer-friendly version

Discussion paper



data were consistently low. The minimum daily average TDT VWC readings between 2018-05-01 and 2018-08-31 were: 1.3, 0.7, 2.3, 0.5, 2.7, 1.6, 2.2, 0.2, 0.3 and 0.4 at the respective depths of 5, 5, 10, 10, 15, 15, 25, 25, 50 and 50 cm.

RC29 328: As you provided SnowFox derived SWE in the data set, its derivation should be documented more precisely (formula, calibration). At least provide it as supplementary material.

AR29 The data set only provides SWE derived from the CRNS (not the SnowFox). We will add clarity and additional detail alongside the citation.

RC30 334: Fig. 8: Same as before (panel IDs and description in the caption), remove x-axes between sub-panels, increase font sizes and avoid red-green coloring in one graph.

AR30 Figure 8 will be amended with your suggestions.

RC31 353: Please explain what "gauge boards" are

AR31 We will amend the manuscript to include additional detail, "In 2020 the network's first gauge board was installed at the Cwm Garw site in Wales. Gauge boards indicate height above ground level (in cm) against which vegetation height and snow depth can be estimated via PhenoCam images. Further gauge boards are planned at sites across the network."

RC32 361-386: The different data sub-products are clearly distinguished and organized in a logical structure. However, on top-level, the user doesn't want to find a 200 files long list to be downloadable click by click. So all the data should be organized as a single file archive (zip) download. The user should be able to use the data set without considering the data-description paper just by making use of the metadata. Thus, the information in table 7 should be provided within 4 additional metadata files (SH, SH...QC, Hourly, Daily). And the JSON format might be a better way to specify things. What's lacking in the metadata is the information how a certain measurement is

[Printer-friendly version](#)[Discussion paper](#)

derived does the timestamp 00 refer to the period 00-30 minutes or 30-00 minutes and has the value been obtained by averaging or as instantaneous value? The timestamp in the data files need to be ISO 8601 compliant (e.g. 2018-01-01T12:00:00Z) so that also the time-zone information is contained. The site metadata should also contain the standard deviations for BD, SOC, and lattice water (which is btw. completely missing). For the easting and northing the projection needs to be specified. Please provide also the hourly raw neutron counts (uncorrected) as well as the Snow-Fox raw and corrected counts as some people may be interested in using their own corrections.

AR32 Thank you for these suggestions. We will update the metadata to improve clarity and data accessibility for users, and enable easier download of files, which we agree will make the data set more usable. We currently provide hourly corrected neutron counts and will consider including the raw neutron counts in future uploads.

RC33 387: Fig. 10: Change to color-blind safe palette (<https://knightlab.northwestern.edu/2016/07/18/three-tools-to-help-you-makecolorblind-friendly-graphics/>)

AR33 The colours in Figure 10 will be amended.

RC34 435-447: It would be nice to have also a short outlook. What are the perspectives for COSMOS-UK? Are there plans to extend the network (e.g., for Scotland)? How long is the projected lifetime of the project. How will the CRNS sensors degrade over time? Do you plan to upgrade the network with more sensitive detectors as they become available on the market?

AR34 Thank you for your interesting questions. COSMOS-UK has been designed as a long-term monitoring network and aims to provide easily-accessible soil moisture and hydrometeorological data; this will continue to be the network's goal whilst dependent on funding. Regarding CRNS sensor degradation, we will consider performing more repeat calibrations to understand any changes. We will continue to consider sensor upgrades when available and feasible. We will amend the manuscript to reflect these

[Printer-friendly version](#)[Discussion paper](#)

points.

## References

Blonquist, J. M., Jones, S. B. and Robinson, D. A.: A time domain transmission sensor with TDR performance characteristics, *J. Hydrol.*, 314(1–4), 235–245, doi:10.1016/j.jhydrol.2005.04.005, 2005.

Colli, M., Pollock, M., Stagnaro, M., Lanza, L. G., Dutton, M. and O’Connell, E.: A Computational Fluid-Dynamics Assessment of the Improved Performance of Aerodynamic Rain Gauges, *Water Resour. Res.*, 54(2), 779–796, doi:10.1002/2017WR020549, 2018.

Köhli, M., Schrön, M., Zreda, M., Schmidt, U., Dietrich, P. and Zacharias, S.: Footprint characteristics revised for field-scale soil moisture monitoring with cosmic-ray neutrons, *Water Resour. Res.*, 51(7), 5772–5790, doi:10.1002/2015WR017169, 2015.

Schrön, M., Köhli, M., Scheiffele, L., Iwema, J., Bogena, H. R., Lv, L., Martini, E., Baroni, G., Rosolem, R., Weimar, J., Mai, J., Cuntz, M., Rebmann, C., Oswald, S. E., Dietrich, P., Schmidt, U. and Zacharias, S.: Improving calibration and validation of cosmic-ray neutron sensors in the light of spatial sensitivity, *Hydrol. Earth Syst. Sci.*, 21(10), 5009–5030, doi:10.5194/hess-21-5009-2017, 2017.

Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B. and Teuling, A. J.: Investigating soil moisture-climate interactions in a changing climate: A review, *Earth-Science Rev.*, 99(3–4), 125–161, doi:10.1016/j.earscirev.2010.02.004, 2010.

Strangeways, I.: Improving precipitation measurement, *Int. J. Climatol.*, 24(11), 1443–1460, doi:10.1002/joc.1075, 2004.

## Response to Referee 2

RC1 1. The main issue is with an absence of calibration details : Table 5 and the dis-



cussion preceding it provide elaborate details of the sampling procedure and determination of soil moisture in the lab. The effectiveness of calibration is however mentioned in just one sentence, “There was  $< 0.03 \text{ cm}^3 \text{ cm}^{-3}$  difference in VWC between the soil moisture determined from these samples and the corresponding daily VWC value derived using the site’s initial calibration data”. Table 5 only shows the reference values and not the difference between the point scale measurements and the CRNS data. In the absence of such results, it is difficult to judge how good has the calibration been. Perhaps in the supplementary material, more detailed results of calibration could be given.

AR1 We will include additional information in Table 5 and amend the text to increase clarity. In our experience the point-scale sensor measurements on the day of calibration are unreliable due to poor contact between the sensor and the soil. This contact improves over time and qualitative comparisons have been made between the CRNS VWC and point measurements. Undertaking quantitative comparisons is planned.

RC2 2. The claim that the spatial data on soil moisture leads to improved hydrometeorological forecasts needs to be substantiated, either by citing appropriate references or through a convincing argument.

AR2 We will amend the text and add references.

RC3 3. Table 1 gives details of the COSMOS-UK stations. Please consider highlighting the mountainous sites, if any among these. The altitude values shown in the table do indicate the heights at which the stations are located, but a higher altitude does not necessarily indicate a station on a mountain. It would also be interesting to see the soil moisture signatures of the mountainous regions as compared to those in the plains.

AR3 The COSMOS-UK sites are installed in non-mountainous and largely flat locations due to the logistical challenges these regions present. We will make this clearer in the manuscript.

[Printer-friendly version](#)[Discussion paper](#)

RC4 4. Is there any irrigation in the area around the CRNS stations? If yes, how has it affected the soil moisture data? Is it possible to filter out the effect of irrigation in the data?

AR4 Sites with regular irrigation were avoided when identifying locations for COSMOS-UK installations. We will be notified of any irrigation at the sites and will explore the impact on data if this occurs.

RC5 5. Table 4 shows that at the same depth two TDT point source sensors are located. How are they spaced? How is the data from the two sensors merged?

AR5 All TDT sensors at COSMOS-UK sites are installed in pairs. Each site has one pair of sensors buried at a depth of 10 cm, located 1 m apart. The additional array of 4 pairs of TDT sensors (where available) are buried at 5, 15, 25 and 50 cm depths, with each sensor located 30 cm from its paired sensor. A horizontal distance of 15 cm is ensured before burying the next pair of sensors at the desired depth. Data from each sensor are provided separately, and the user can choose whether to combine data across pairs. We will make this clearer in the manuscript.

RC6 6. Lines 183-184 : Volumetric samples are taken at five depths, upto 0.25 m bgl. However, the CRNS data is between 0.1 m to 0.8 m. Is there any calibration for depths below 0.25 m?

AR6 There is currently no adjusted calibration applied for depths below 0.25 m. Soil sampling depths were selected to match typical (moist) UK conditions. Ideally sites should have repeated calibrations at different VWCs to capture the full range of measured VWC. We will amend the text and emphasise that the shallow soil layers are given higher weighting than deeper layers to accommodate greater water contribution at shallow depths (Köhli et al., 2015; Schrön et al., 2017).

RC7 7. Fig 2 (appearing around line 272) should be Fig. 6.

AR7 Thank you for your comment. This is correct in the preprint version of the

[Printer-friendly version](#)[Discussion paper](#)

manuscript.

RC8 8. It is difficult to understand the “automatic processing” for quality checks. Table 6 provides the flags raised for various events, but how these events are identified in the data is not clear (for example, how does the automatic processing detect simultaneously, missing data and small sample of data?)

AR8 The explanation will be amended to increase clarity.

RC9 9. Fig. 8 caption may be made more descriptive. Also, the caption states that these are the observations required for PE calculation, but the last panel in fact shows the PE calculated! It would also be interesting to see the soil temperature plot along with the other variables shown here (although soil temperature is not used in PE calculation).

AR9 Thank you for your comment and suggestion. We will improve the caption for Figure 8.

RC10 10. Please also discuss how the soil moisture measured at these 51 stations may be smoothed to upscale it to the national scale (see line no. 396).

AR10 We will add this discussion to the manuscript.

## References

Köhli, M., Schrön, M., Zreda, M., Schmidt, U., Dietrich, P. and Zacharias, S.: Footprint characteristics revised for field-scale soil moisture monitoring with cosmic-ray neutrons, *Water Resour. Res.*, 51(7), 5772–5790, doi:10.1002/2015WR017169, 2015.

Schrön, M., Köhli, M., Scheiffele, L., Iwema, J., Bogena, H. R., Lv, L., Martini, E., Baroni, G., Rosolem, R., Weimar, J., Mai, J., Cuntz, M., Rebmann, C., Oswald, S. E., Dietrich, P., Schmidt, U. and Zacharias, S.: Improving calibration and validation of cosmic-ray neutron sensors in the light of spatial sensitivity, *Hydrol. Earth Syst. Sci.*, 21(10), 5009–5030, doi:10.5194/hess-21-5009-2017, 2017.

Interactive comment on Earth Syst. Sci. Data Discuss., <https://doi.org/10.5194/essd-2020-287>, 2020.

**ESSDD**

---

Interactive  
comment

Printer-friendly version

Discussion paper

