

Earth Syst. Sci. Data Discuss., author comment AC1
<https://doi.org/10.5194/essd-2021-270-AC1>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Reply on RC1

Hou Jiang et al.

Author comment on "Multi-resolution dataset for photovoltaic panel segmentation from satellite and aerial imagery" by Hou Jiang et al., Earth Syst. Sci. Data Discuss., <https://doi.org/10.5194/essd-2021-270-AC1>, 2021

Comment 1. The dataset was collected in Jiangsu Province. Considering that the changes in geographic context may affect the model's performance, how can you guarantee that the model trained with this dataset can be generalized to other regions?

Response: We compared the samples from Gaofen-2 and Beijing-2 images, and found that PV panels exhibit similar characteristic in high-resolution imagery and the main difference comes from the backgrounds. The changes in geographic context will inevitably affect the performance of deep networks. To mitigate such effects, we tried to collect PV samples covering as many backgrounds as possible to enhance the generalization ability of deep networks trained by our dataset. The installed PVs in Jiangsu province are distributed on various land covers, such as, sparse shrubs, low-density grasslands, reservoirs, ponds, saline alkali lands and rooftops, making our dataset representative for most cases. Besides, some skills in deep learning community can be adopted to guarantee the generalization ability in other regions, such as transferring learning, cross-domain feature representation. In completely different areas where direct application may fail, using a small number of samples in the target area to fine-tune the pre-trained model can also achieve satisfactory results. Therefore, although the samples were collected in Jiangsu Province, there is no need to worry too much. Furthermore, our work will continue, and we plan to collect more PV samples in China and around the world through automatic deep learning algorithm and manual discrimination.

Comment 2. The dataset is composed of Gaofen-2 and Beijing-2 imagery, which include a near-infrared band in addition to RGB bands. Does this mean that the model trained with this dataset can only be applied to the satellite imagery with a specific combination of spectrums? Are these two imagery representative of the current common satellite imagery?

Response: Actually, all samples in our dataset are composed of red, green and blue (RGB) bands of used satellite and aerial images. Our experiments (Section 4.1) also demonstrate that RGB bands are enough for deep networks to distinguish PV panels from various backgrounds. Therefore, the near-infrared band of Gaofen-2 and Beijing-2 images is not contained in our samples. We have made it clear in Section 3 as "The shapefile of polygonal annotations was converted to a raster that has the same spatial resolution as satellite or aerial images. The raster and original red, green and blue (RGB) images were then seamlessly cropped into tiles at a fixed size by referring to the sampling grids." The comparison of PV samples from Gaofen-2 and Beijing-2 imagery shows that different PV

panels exhibit similar characteristic after histogram equalization. Thus, we believe that the two satellite imagery are representative for common applications.

Please also note the supplement to this comment:

<https://essd.copernicus.org/preprints/essd-2021-270/essd-2021-270-AC1-supplement.pdf>