

Earth Syst. Sci. Data Discuss., author comment AC2
<https://doi.org/10.5194/essd-2021-82-AC2>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.



Reply on RC2

Joaquín Muñoz-Sabater et al.

Author comment on "ERA5-Land: a state-of-the-art global reanalysis dataset for land applications" by Joaquín Muñoz-Sabater et al., Earth Syst. Sci. Data Discuss., <https://doi.org/10.5194/essd-2021-82-AC2>, 2021

We would like to thank this reviewer for his many useful and relevant comments that has helped us to improve the text of the manuscript and figures. Please, find here below a point-by-point response to all your major comments and minor comments:

1) I see mainly four reasons why ERA5-Land is supposed to be superior to ERA5:

- (i) higher spatial resolution of static land model parameters
- (ii) higher spatial resolution of downscaled forcing
- (iii) reduced impact of discontinuities by longer spin-ups for the segments
- (iv) newer LSM version used in ERA5-Land

How does trend analysis benefit from these four reasons? Indeed, ERA5-Land will be used for trend analysis but since it is not included in the evaluation, what are the expected improvements relative to ERA5? The skill improvement that is currently presented was probably mainly due to the higher spatial resolution, I am wondering how relevant (for which variables) higher spatial resolution is for trend analysis? If it is decided not to provide a trend analysis in this paper, it would be useful to read about the authors' opinion as a first guidance for the user community regarding trend analysis. Thinking of trend analysis of river discharge, it should be emphasized that crucial factors like land use change are not included in the land surface model. I recommend a discussion section on implications of the current results + reanalysis setups for trend analysis.

Many thanks for your very interesting comments. We believe the comments about the trends would be very relevant if the paper would conduct a trend analysis. However, although we believe that this dataset enables trend analysis, it is out of the scope of this paper. Trend analysis deserves a dedicated study and we cannot fit it in a data presentation paper which is already quite long by itself. We have tried to make it clear in Lines 5-6 of the abstract (see response to comment (1) of reviewer-1), and also we have added a sentence in line 657 of the "conclusions and discussion" section of the revised

manuscript:

Line 657 “[...] and the consistency presented over multi-decadal time scales (that could set the basis to enable reliable trend analyses), all of them making ERA5-Land a state-of-the-art dataset for multiple land applications”.

It is difficult to provide an educated guess of how ERA5-Land trends will compare to those of ERA5 without conducting the analysis itself. This paper has shown that, based on the evaluation carried out over several key variables of the water cycle, ERA5-Land is more accurate at least at representing the spatial/temporal evolution of the hydrological cycle. The differences with ERA5 are not enormous though, so we believe that trends will not be very different at global scale. However locally, where resolution matters, it could provide more accurate trends. In particular trends at areas with low soil moisture variability, where the reduced impact of discontinuities by longer spin-ups is important, could benefit from ERA5-Land multi-decadal consistency. At this respect we have also added a sentence in the discussion part, in lines 658-660 of the revised manuscript:

“The reduced impact of discontinuities by using longer spin-up periods, could also be a crucial factor to obtain accurate trends over multi-decadal periods for variables slowly changing in time, for instance the root-zone soil moisture (see bottom panel of Fig.3)”.

We agree that factors such as land use change are crucial for accurate land surface modelling and trend analysis, and those are in the current plans of the ECMWF land surface model evolution. This point, as well as the use of a monthly (static) LAI, has already been described in the ‘perspectives’ section, in lines 697-708 of the original manuscript.

(2) One application of ERA5-Land will be to force other land models with higher temporal and spatial resolution atmospheric input. An application is indicated by the evaluation with GLEAM, though differences between ERA5 and ERA5-Land driving GLEAM were very small. Consider to extend the related discussion section (e) and include a discussion of the benefits from downscaled precipitation for other land models.

This is a really good point raised by the reviewer. A short discussion on precipitation is already included in the paper in the “perspectives” section, however we have also extended it as to comment on the benefits from downscaled precipitation. In the perspectives, after “as well as their availability in near-real-time” In lines 697-699, we have added the following paragraph:

“In addition to coarse-scale near-real-time bias-correction, high resolution downscaling could be also explored. Such a correction could be based on a climatological rescaling of precipitation based on a high resolution reference climatology (e.g. Karger et al. 2017)”

For Karger et al. 2017, the new reference is:

Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., and Kessler, M.: Climatologies at high resolution for the earth’s land surface areas, *Sci. Data*, 5, 170 122, <https://doi.org/10.1038/sdata.2017.122>, 2017.

(3) The new PET variable is included in the portfolio of ERA5-Land, which is

briefly mentioned with a note of caution on its use because the atmosphere is not affected by the water-unlimited land surface assumed for PET. That seems to me being a general issue of PET. How is this different for the ERA5-Land PET from other PET approaches e.g. discussed in <https://doi.org/10.5194/hess-23-925-2019> ?

We agree with the reviewer on this comment and we acknowledge that the main problem of the PET concept is that it is related to a hypothetical quantity defined for hypothetical environmental conditions which hampers experimental validations. The ERA5-Land PET is only one option among others such the Priestly-Taylor formulation or the radiation-based ones. A rational of the concept of evaporation under unstressed soil moisture conditions would require mapping effects on net radiation (which is accounted for in the second call to the surface model) without atmospheric feedbacks. However, there is always a hypothetical element in the atmospheric feedback assessment, as it depends on the spatial extent of the target unstressed area. Small plots in arid environments will generate advection effects, but large areas will lead to atmospheric moistening considerably suppressing evaporation, and advection effects confined to the edges of the considered area. This to emphasize the complexity of the PET estimation problem and that it would depend on the target application being dependent on the spatial (and temporal) scale and eventually on the irrigation regime.

Also to note that PET can also be estimated at a daily time scale with the Priestly-Taylor formulation using daily averages of net radiation, temperature, humidity, and wind from ERA5/ERA5-Land without need of model timestep integration (which is another option to the ERA users).

(4) Results indicate higher bias for ERA5-Land energy fluxes. Was the EC data corrected for energy balance closure gap?

The EC data was not corrected for energy balance closure. Eddy covariance In-situ flux measurements can be affected by systematic biases due to issues affecting the energy closure. However, the correction of those imbalances require making assumptions on the sources of the errors, which can also impact the fluxes. Furthermore all the measured fluxes (including ground-heat flux) are needed, and they are not available from many sites. Therefore, no correction of the EC data was considered in this study.

See an extended discussion of this point here below, in response to a comment on Line 340 of the original manuscript.

(5) I found it interesting that the river discharge skill mostly improved for ERA5-Land relative to ERA5. Based on the snow depth evaluation, I would have assumed that discharge for watersheds with significant snow melt might deteriorate due to lacking assimilation of snow obs in ERA5-Land compared to ERA5. It could be discussed more in detail why snow DA did not help ERA5 to perform better. Further, river discharge of large-scale routing models are often evaluated not at daily but at coarser resolution, e.g. monthly, due to large uncertainties of routing parameters that makes it unrealistic to predict the timing of peaks at daily resolution. To be more comparable with other routing scheme results, Figure 13 (with reduced (a) figure, see annotated pdf) could be also provided for monthly resolution.

As described in Zsoter et al. 2019, the snow data assimilation was shown to be

detrimental in much of the Northern Hemispheric snow impacted areas in the ERA5 experiment, compared with the offline simulation without coupling and land data assimilation. The average snow increments in ERA5 are negative, due mainly to the too slow snow melt in the HTESSEL, which consequently removes water from the hydrological system. This contributes to the decrease of the dominantly negative biases in a large area with generally deteriorating the hydrological performance.

To make this point more clear, we have added the following paragraph, in lines 617-622 of the revised manuscript, in the relevant part of the discussion section:

“Also, the fact that ERA5-Land does not directly assimilate any observation, as is the case for ERA5, has shown to have positive impacts on the closure of the water balance (Zsoter et al., 2019). They have shown that the snow data assimilation is detrimental to the hydrology in large parts of the snow impacted Northern Hemisphere in the ERA5 experiment, compared with the offline simulation without coupling and land data assimilation (such as ERA5-Land). The average snow increments in ERA5 are negative, due mainly to the too slow snow melt in CHTESSEL, which consequently removes water from the hydrological system. This contributes to decreasing the dominantly negative biases in a large area with subsequently deteriorating the hydrological performance. The exception is in the West US and Amazonian basins.”

With respect to the use of daily, rather than monthly; to be able to appropriately evaluate the ability of a model to simulate river discharge, especially with regards to a model’s ability to capture the timing of river discharge peaks, it is necessary to use a minimum of a daily model time-step, which is also the time resolution of the available observations. The aim of the evaluation here was to assess the ability of GloFAS-ERA5 compared to GloFAS -ERA5-Land in capturing key hydrological dynamics of timing, bias, and variability – performed with the modified Kling-Gupta Efficiency (KGE) statistical metric. This is a widely used standard methodology in hydrological model evaluation. We hence prefer to maintain the analysis of river discharge using daily time-steps.

(6) The main technical aspect of the production of ERA5-Land, that was not sufficiently explained to me, is how the "Integration of the land surface model in 24 cycles" works. Please provide longer explanation of what is done here and why.

The “24-h cycles” is a convention that has been used to make it easier the retrieval of ‘forecasted’ fields by ERA5-Land users (as it was done for ERA-Interim) and limit the accumulation period of surface fluxes to 24h. ERA5-Land is run as a continuous simulation (in 24 hour’s chunks with reproducible restarts) forced by ERA5 near-surface meteorological variables, so although theoretically there aren’t ‘analysis’ fields, some of them are grib-coded as ‘analysis’ and others as ‘forecasts’. In the C3S catalogue the ‘forecast’ fields are valid at a specific time of the day (00 to 23UTC). We could have equally selected 12h, 48h cycles or a different one, but it would have been much more complex to understand the validity time of a ‘forecast’ field.

We believe that the reason above is too technical to be explained in detail in a data presentation paper. In section 7 we have explained how the accumulation periods works, and we have directed interested readers about the technical details to the online documentation that provides further details.

(7) Regarding water management applications, it's important to mention to the

user community that ERA5-Land lacks a groundwater storage. Something that should be clarified at some place.

The reviewer is right. We are currently working with a modular, flexible system called ECLand. This system allows to develop several modelling aspects separately, such as the open water areas and river/inundation. Currently there are some ongoing developments with coupling runoff with a river discharge system (CamaFlood), and indeed in the longer term the introduction of a groundwater storage will also be considered as part of our developments. We have added this point in the perspectives section, by replacing:

“All the components discussed above are currently under research and, along with the improvement in other components of the land surface scheme (increase of the number of soil layers, introduction of a multi-layer snow scheme, reduction of model time step, etc.), provide the basis for a future new version with improved accuracy for the land states at multi-decadal timescales” by

“All the components discussed above are currently under research, but they are not the only ones. ECMWF is currently working with a flexible, modular system called ECLand (Boussetta et al., 2021), which allows to develop separately several modelling aspects of the land surface, such as the increase of the number of soil layers or the introduction of a multi-layer snow scheme, as well as progressing on other longer term perspectives such as the introduction of a groundwater storage or the reduction of the model time step. All the above ongoing developments will provide the basis for a future new version with improved accuracy for the land states at multi-decadal timescales.”.

The added reference is the following:

Boussetta, S., Balsamo, G., Arduini, G., Dutra, E., McNorton, J., Choulga, M., Agustí-Panareda, A., Beljaars, A., Wedi, N., Muñoz Sabater, J., de Rosnay, P., Sandu, I., Hadade, I., Carver, G., Mazzetti, C., Prudhomme, C., Yamazaki, D., and Zsoter, E.: ECLand: The ECMWFLand Surface Modelling System, *Atmosphere*, 12, <https://doi.org/10.3390/atmos12060723>, 2021

(8) Figure quality and captions would strongly benefit from a thorough revision, such as

- Consistent use of (a) (b) etc. instead of top/middle left/right that is also sometimes used
- Explanation of abbreviations and variable names in caption
- Same chronological order of ERA reanalysis products (currently ERA5-Land sometimes comes last, sometimes first in figures).
- Consistent use of variable names.

For related and other comments, see annotated pdf.

Thanks, we have tried to make figures more consistent by following all reviewer recommendations.

(9) The lake temperature skill of ERA5-Land depended a lot on the lake depth information. The evaluation was thus stratified by whether lake depth improved

(got more realistic) in ERA5-Land compared to ERA5. It's unclear on which information this stratification was made. How can a higher spatial resolution can lead to less realistic lake depth in ERA5-Land?

The current lake mean depth file used at ECMWF is based on the Global Lake Database (See Table S1 of the original manuscript), and it has native resolution of 1 km. It consists of more than 14,000 in situ lake mean depth observations mapped on to a gridded file, and of indirectly estimated depths for all the rest of lakes based on a geomorphological approach. This means that the mean depth for some lakes which are not common for certain geomorphological regions might be estimated incorrectly – i.e., they can be overestimated or underestimated. Once lakes depths are aggregated to a coarser resolution, the depth for uncommon lakes can change and even become closer to the observed value but for the wrong reason (due to the other lake depths in the vicinity).

(10) Figure 15: the spread of the bias values for the Bowen ratio is strongly increased for ERA5-Land. Either it's wrongly calculated/shown here or requires clarification.

The code to produce these plots has been reviewed and we can confirm there are no errors. Nonetheless, the calculation of the Bowen ratio is usually prone to error, since it results from the ratio between two fluxes which can be positive or negative (so it is numerically unconstrained). This may create differences when large pixels are compared to in situ measurements. To reduce the influence of outliers, fluxes are first aggregated to daily resolution before calculating their ratio, as recommended by Shuttleworth (2012). We acknowledge, in any case, that the spread of the bias for the Bowen ratio is a bit higher for ERA5-Land. This relates mainly to a few outliers, as the quantiles are quite similar (as shown by the dashed lines that actually match up when comparing ERA5 and ERA5-Land), and close to each other. We computed the number of outliers found that are either " $< q_{25} - 1.5 * (q_{75} - q_{25})$ " or " $> q_{25} + 1.5 * (q_{75} - q_{25})$ ". For ERA5 we found 12 observations meeting these conditions, whereas for ERA5-Land we found 20. This is not a strict definition of 'outlier', but it is analogous to the way they were computed by Martens et al. (2020).

The two references in this response:

Shuttleworth, W.J. (2012). Surface Energy Fluxes. In *Terrestrial Hydrometeorology*, W.J. Shuttleworth (Ed.). <https://doi.org/10.1002/9781119951933.ch4>

Martens, B., Schumacher, D. L., Wouters, H., Muñoz Sabater, J., Verhoest, N. E. C., and Miralles, D. G.: Evaluating the land-surface energy partitioning in ERA5, *Geosci. Model Dev.*, 13, 4159–4181, <https://doi.org/10.5194/gmd-13-4159-2020>, 2020.

(11) Several more comments are included in the annotated pdf;

L36: We have added two extra, more recent references to highlight the need for land model developments:

Vereecken, H., Weihermüller, L., Assouline, S., Šimůnek, J., Verhoef, A., Herbst, M., Archer, N., Mohanty, B., Montzka, C., Vanderborght, J., Balsamo, G., Bechtold, M., Boone, A., Chadburn, S., Cuntz, M., Decharme, B., Ducharne, A., Ek, M., Garrigues, S., Goergen, K., Ingwersen, J., Kollet, S., Lawrence, D.M., Li, Q., Or, D., Swenson, S., de Vrese, P., Walko, R., Wu, Y. and Xue, Y. (2019), Infiltration from the Pedon to Global Grid Scales:

An Overview and Outlook for Land Surface Modeling. *Vadose Zone Journal*, 18: 1-53
180191. <https://doi.org/10.2136/vzj2018.10.0191>

Boussetta, S., Balsamo, G., Arduini, G., Dutra, E., McNorton, J., Choulga, M., Agustí-Panareda, A., Beljaars, A., Wedi, N., Muñoz Sabater, J., de Rosnay, P., Sandu, I., Hadade, I., Carver, G., Mazzetti, C., Prudhomme, C., Yamazaki, D., and Zsoter, E.: ECLand: The ECMWF Land Surface Modelling System, *Atmosphere*, 12, <https://doi.org/10.3390/atmos12060723>, 2021

L38: We believe the sentence is right. Atmospheric reanalysis are produced with a single version of the coupled atmosphere-ocean-land prediction system and therefore show, despite the analysis increments, temporal consistency over the production period. This doesn't occur with the forecast models (as the IFS), which every certain period of time are updated (parameterizations, assimilation system, etc.).

L41: done, thanks.

L57: We prefer to keep the wording "climate studies" since it includes in a broader sense several inter-related investigations such as "climate change studies", "climate modelling studies" or "climate trend studies".

L86: the type of grid has been added to the main manuscript, as well as a supporting reference: "ERA5-Land produces a total of 53 variables describing the water and energy cycles over land, globally, hourly and at a spatial resolution of 9 km, matching the ECMWF triangular-cubic-octahedral (TCO1279) operational grid (Malardel et al., 2016)".

And the reference added:

Malardel, S., Wedi, N., Deconinck, W., Diamantakis, M., Kuehnlein, C., Mozdzyński, G., Hamrud, M., and Smolarkiewicz, P.: A new grid for the IFS, *ECMWF Newsletter*, 146, 23–28, <https://doi.org/10.21957/zwdu9u5i>, 2016.

L102: coordinates swapped. Thanks.

L103: ERA5-Land has been added to the figure and now it is used consistently over all the manuscript.

L117: "[...], no prior long ERA5 stream was available." has been replaced by "[...], a long, prior ERA5 stream was not available". For consistency the "prior long" in Line 110 has also been replaced by "long, prior".

Figure 3 → the figure has been improved as suggested by the reviewer.

L118: The following sentence "[...], and then letting three spin-up years." has been replaced by "[...], and then allowing the system to spin-up for three years".

L120: for the 3rd stream only one spin-up year was feasible (1949) because there wasn't ERA5 forcing available prior to Jan 1949. The available climatology (1981-2010) was used to initialize the spin-up year. This was indicated in Line 120 of the original manuscript.

L127: "approximately 10 m" → It is indeed 10 m exactly. The text has been changed to "[...], which is 10 m above the surface [...]"

L133: "conventional" refers to all conventional meteorological observations (ships,

radiosondes, SYNOP stations, etc.). We have added the word “conventional meteorological” to the main text.

L151: Fig.4 → Following the reviewer recommendation, we have included the location of the three mentioned lakes in the top-right subplot of Fig.4, and updated the caption of the figure accordingly.

L162: Specify which carbon fluxes are estimated. Is heterotrophic respiration simulated? Are longer term carbon stock changes estimated?

The carbon fluxes are the Net ecosystem exchange (NEE), the Gross Primary production (GPP) and the Ecosystem respiration (Reco). Long term carbon stock is not estimated, instead ecosystem respiration is parametrized in an NWP adapted way as a function of land-use and soil temperature. The type of carbon fluxes has been added in the main text, lines 164-165 of the revised manuscript.

Fig4.--> We have replaced ERA5L by ERA5-Land in the title of each subplot. We have also defined LST in the figure caption.

L173: Which crop? Constant parameters? I guess it's not seasonally changing and does not include harvest and bare soil period?

There is no sub-type of crops (such C3 OR C4) considered in the actual version, it is the crops and mixed farming type from the BATS classification (table 8.1 of the IFS documentation). It is parametrised with a constant canopy resistance with no consideration of seasonal change or harvest.

L183: Yes, thanks. Added in parenthesis “the latter also a component of the water cycle”.

L213: “OI” replaced by “Optimal Interpolation”.

L235: Thanks, “averaged” replaced by “average”.

Eq.1: Thanks for this comment. Indeed, the reviewer is right that the results will be equivalent either normalizing or not by the standard deviation. However, the standard definition of anomaly correlation for soil moisture is as displayed in eq.1, and for consistency with many other paper in this respect we prefer to keep it as it is.

L240-241: STDD: It is indeed equivalent to ubRMSE. We have added to the main text for clarification.

L327: Thanks to point this out. Although the reviewer is technically correct, the equation is a direct substitution of terms into the general skill score equation, so we believe it is clearer to leave it as it is in the text.

L340: gaps: The data downloaded from FLUXNET is gap-filled using different methods, each of them with some uncertainty. They also include a flag to indicate which method was used. If there was gap-filling in the time series (the flag is set) it simply means there was no measurement at that time. If it is gap-filled these data were not used as validation data, because there is some uncertain algorithm behind it. Therefore, a ‘gap’ in the text means any record that was gap-filled or has an NaN (that could not be gap-filled). We simply masked out those gap-filled records to retain only the actual measurements from the eddy-covariance sites.

We agree that the sentence in the main text is quite confusing, so we have removed the sentence in Line 346 of the original manuscript: “After quality control of in situ stations,

only records without gaps during the whole period 2001-2014 were retained.”, and we have replaced in lines 345-346: “(2) the removal of gap-filled records.” by “(2) the removal of gap-filled records to retain only the actual measurements from the eddy-covariance sites”.

Energy Balance Closure: The reviewer is quite right pointing this out. Indeed, we find that while some authors have applied energy balance closure, many others haven't done it. The reason is that there isn't clear accepted guidelines on how to do it properly, and in addition all the fluxes measured (including ground-heat flux) are needed to this end. Not every EC tower measures all the fluxes (ground heat flux is for instance often missing) which means that the sample of towers we can use for validation would drop drastically.

In order to make this point clear, we have added the following sentence in the main text, lines 348-352 of the revised manuscript: "Note that the measured energy fluxes used as reference in this paper were not corrected for energy balance closure because the number of towers used for validation would be drastically reduced, as the ground-heat flux is also needed and is not available from many towers. Some authors have already highlighted the lack of closure in the energy balance at eddy-covariance sites and a consequential tendency to underestimate the latent heat flux (Wilson et al., 2002, Ershadi et al., 2014, Jimenez et al., 2018)."

The added references are:

Wilson, K., Goldstein, A., Falge, E., Aubinet, M., Baldocchi, D., Berbigier, P., Bernhofer, C., Ceulemans, R., Dolman, H., Field, C., Grelle, A., Ibrom, A., Law, B., Kowalski, A., Meyers, T., Moncrieff, J., Monson, R., Oechel, W., Tenhunen, J., Valentini, R., and Verma, S.: Energy balance closure at FLUXNET sites, *Agric. Forest Meteorol.*, 113, 223–243, doi:10.1016/S0168-1923(02)00109-0, 2002.

Ershadi, A., McCabe, M., Evans, J., Chaney, N., and Wood, E.: Multi-site evaluation of terrestrial evaporation models using FLUXNET data, *Agric. Forest Meteorol.*, 187, 46–61, <https://doi.org/https://doi.org/10.1016/j.agrformet.2013.11.008>, 2014.

Jiménez, C., Martens, B., Miralles, D. M., Fisher, J. B., Beck, H. E., and Fernández-Prieto, D.: Exploring the merging of the global land evap-955oration WACMOS-ET products based on local tower measurements, *Hydrol. Earth Syst. Sc.*, 22, 4513–4533, <https://doi.org/10.5194/hess-22-4513-2018>, 2018.

L365: “at a daily time scales” replaced by “at daily time scales”.

L409: Thanks to spot the error, “represents” replaced by “represent”.

L417: “m a.s.l.” added here and elsewhere in the main text

Fig.9 caption updated as suggested.

L448: the Mean Absolute Error has been added to the text on top of the percent.

L456: The statistical significance was tested with the Kruskal-Wallis test by ranks (best for non-Gaussian distribution variables), as written in line 287 of the original manuscript. We have deleted the word "modest".

Fig.12 caption: We have removed the gradations and added the degree symbol, and for consistency we have done the same for Fig.11.

Fig.13. Fig. 13 summarises the distribution of scores for 1285 river discharge stations

as a CDF for both models (i.e. GloFAS-ERA5 and GloFAS-ERA5-Land). This is a valid and commonly used method for hydrological model performance comparison (e.g. Alfieri et al., 2020; Kratzert et al., 2019). The reason why panel (a) is larger is that it is the main result for the overall KGE' metric. Because the KGE' can be decomposed into three components (i.e. correlation, bias ratio and variability ratio), we added this additional information as three secondary panels (i.e. (b), (c) and (d)), to help diagnose the patterns seen in the summary KGE' in (a).

The two references in this response are:

- Alfieri, L., Lorini, V., Hirpa, F. A., Harrigan, S., Zsoter, E., Prudhomme, C., and Salamon, P.: A global streamflow reanalysis for 1980–2018, *Journal of Hydrology X*, 6, 100049, <https://doi.org/10.1016/j.hydroa.2019.100049>, 2020.
- Kratzert, F., Klotz, D., Shalev, G., Klambauer, G., Hochreiter, S., and Nearing, G.: Towards learning universal, regional, and local hydrological behaviors via machine learning applied to large-sample datasets, 23, 5089–5110, <https://doi.org/10.5194/hess-23-5089-2019>, 2019.

L483-484: Latter in line 484 it is explained that the bias show (observation – model). Anyway, following the reviewer recommendation we have moved the description of bias (and other metrics used for the energy fluxes) to section 3.7.1, and made this sentence clearer: “The left panel of Fig 15 shows the violin plots of the turbulent fluxes against in situ eddy-covariance measurements for ERA-Interim and ERA5-Land” has been replaced by “The left panel of Fig 15 shows the violin plots of the ERA-Interim and ERA5-Land turbulent fluxes (compared to in situ eddy-covariance measurements) ”

L485: anomaly correlation: We use the anomaly correlation as the temporal variability of the turbulent fluxes is strongly influenced by the seasonal cycle of its main drivers at the scales considered in this experiment, and therefore the performance of the land-surface schemes in response to anomalous weather conditions (i.e. with respect to the seasonal cycle) might be masked when raw time series are analysed. We have also used the standardised anomalies to allow us to directly compare the quality of the turbulent fluxes and the Bowen ration, despite their different order of magnitude. This was already explained in lines 346-349 of the original manuscript.

Fig. 15: Thanks; “Unlikely most metrics and variables” has been removed from the caption, making it more clear now.

We have tried to increase the font of the labels. However, since this is a side-by-side figure with two subplots, by increasing the font size the violin plots look smaller, so we decided to keep the current font size. However, we have moved the titles on top as recommended by this reviewer.

L491: We agree with the reviewer, and to make it more consistent through the main text, we have replaced all “sshf and slhf” entries by “H” and “ λpE ”, respectively.

L498-499: We already provided a plausible explanation in the “conclusions and discussion” section, in lines 617-619 of the original manuscript.

Fig.16: To generate this figure we used a projected CRS, which allows to show the background map. We have added the X (Easting) and Y (Northing) coordinates to the figure in EPSG3857 (<https://epsg.io/3857>), but not lat-lon as they just would be approximate. The result is the following (see figure attached to this comment).

We believe that they look more confusing than before, and the background map provides more context on the location of the sites than simple lat/lon coordinates. So we decided to leave the figure as it was shown originally.

L504: Please, see response to comment (10)

L506: We have revised this sentence and we have removed "and for the stations where the recorded air temperature is milder", as the figure is indeed not conclusive for air temperatures.

L512: Thanks to spot the misspelling, "term" has been replaced by "terms".

L515-516: Just to clarify: GLEAM+ERA5 is forced with T and R from ERA5, whereas GLEAM+ERA5-Land is forced with T and R from ERA5-Land, which is described in Section 3.7.2. However we agree that this sentence is confusing and therefore we have decided to remove it from the main text.

Fig.19: As recommended. by the reviewer, the ranges have been adjusted for subplots c) and d). And indeed the improvement from ERA5-Land to ERA-Interim in terms of correlation values is now much more clear. We have also increased the resolution of the figure. In addition and to be consistent, we have updated the figure S5 of the supplementary material.

Fig.19 caption. Thanks to spot the error, indeed the figure refers to ERA5-Land and not to ERA5. Corrected. In addition, we have rephrased the whole caption as follows:

"Global statistics of ERA5-Land (left column) and ERA-Interim (right column) Land Surface Temperature (LST, in K) for the time period 2003-2018. The first row are the global maps of mean LST, the second row are the correlation maps and third row are the RMSE maps, with respect to the MODIS LST average ensemble. Note that the LST here refers to the skin temperature."

Fig.20: Each triangle represents a selected pixel, but the colour hasn't got any other meaning.

The word "quantitative" has also been removed from the caption as suggested.

L552: As recommended by the reviewer, we have improved the wording of this sentence:

"Overall, the water cycle is improved in ERA5-Land compared to ERA5 according to the different variables evaluated, whereas the energy cycle variables show similar performance; both ERA5 and ERA5-Land perform substantially better than ERA-Interim. The main evaluation findings are as follows":

L554: Here we didn't use significance in a statistical sense. To clarify this point we have rephrased the whole sentence in Line 562-563:

"Significant improved correlation of the ERA5-Land (compared to ERA-Interim and ERA5) root-zone soil moisture absolute and anomaly values with respect to in situ measurements is achieved for almost all networks assessed" has been replaced by:

“Soil moisture: the boxplots show a consistent improvement of the statistical metric distributions of ERA5-Land with respect to those of ERA5, in particular the improvement is more marked for the root-zone soil moisture.”

L562: The sentence is about the minimum saturation level linked to the soil type and in the top layer both evaporation and transpiration would occur so we prefer to keep the term evapotranspiration

L590: By "lakes with more realistic depths in ERA5-Land" we mean taking into account, in the statistics, only lakes whose depth is closer to the observations using the ERA5-Land spatial grid, and not taking into account lakes whose depth remained unchanged either in ERA5 or in the ERA5-Land grid. This is because when spatial resolution changes some lake depths become closer to the in situ observations, some further from in situ, and some lake depths remain the same - for instance because it is a big lake without bathymetry. In order to make this sentence more clear, we have replaced:

“Isolating lakes with more realistic depths in ERA5-Land reduces [...]” by “Isolating lakes whose depth is more realistic using the ERA5-Land grid (i.e. only lakes whose depth in the ERA5-Land grid match better in situ observations) the LSWT errors are reduced by more than 20%”.

See also our comment to main point (9) of this reviewer.

L595-599: We apologize for the confusion created, as the sentences refer to different validation datasets (Alqueva, Finish lakes or the global inventory). We have rephrased lines 589-600 of the original manuscript to make these conclusions more clear:

“[...]. The higher resolution of ERA5-Land allows, in many cases, a more accurate specification of lake depth. Thus, isolating lakes whose depth is more realistic using the ERA5-Land grid (i.e. only lakes whose depth in the ERA5-Land grid matches better in situ observations), the LSWT MAE is reduced by more than 20%. The positive influence of a higher resolution atmospheric forcing was also verified by isolating lakes whose depth remained unchanged either using ERA5 or ERA5-Land.

Both reanalysis LSWT estimates were also compared using as reference a global inventory (1995-2009) based on summer observations from satellite sensors. In this case, the performance of both reanalysis is quite similar (excluding exceptional lakes where Flake performs poorly). The uncertainty of lake depths in the global inventory is larger and it could have an impact on these results. However, even when comparing only lakes with a depth less than 50 m (Flake was designed for medium-depth lakes) the performance of ERA5-Land LSWT does not show significant improvement compared to ERA5 (based on MAE). This might be due to the averaging technique used for the in situ measurements, which uses satellite measurements representing one instant in time rather than continuous hourly data. Nevertheless, in summer months, ERA5 LSWT biases are on average 2.2°C cooler than observations, whereas ERA5-Land LSWT is just 1.3°C cooler.”

L600: Thanks to point this out. Due to the high resolution grid of ERA5-Land, the mean depth of many lakes are more realistic in ERA5-Land, what it implies that the total volume is improved as well. So we believe that the improvement of the lake characterization also helps to improve the river discharge at global scale. We have added the word “[...] and lakes characterization [...]”

L610: In the Amazon basin, the bias term is a little bit lower in ERA5-land (so bit less water), but it is really a subtle difference. Being ERA5 precipitation a forcing parameter of ERA5-Land, we assume similar precipitation values in both reanalysis, so an overestimation of evaporation remains the most likely reason. Since the differences are, for most discharge stations, very small, we really didn't undertake a deep diagnostic though.

Consequently, we have replaced "The degradation in the Amazon is small, but it is linked to an underestimation of the surface and subsurface runoff (not shown)." by "The degradation in the Amazon is small, linked to an underestimation of the surface and subsurface runoff (not shown) that is likely caused by a slight overestimation of evaporation."

L639: We have replaced "radiative aspect" by "surface fluxes". Thanks.

L641: "land surface" replaced by "land surface modeling"

L642: We agree that a trend analysis would be beneficial, but it is out of the scope of this paper and it would require another dedicated paper only for this aspect. See also extended responses to comment (1) of reviewer 1 and 2.

L646: "international" replaced by "internationally".

L656: ";" replaced by "."

L677: Please see response to comment in line 173

L680-681: The canopy resistance for water vapour is computed with the JARVIS approach, then an A-gs based formulation is derived to compute the carbon fluxes. This modular approach is adopted in order not to deteriorate the NWP scores which are calibrated based on the JARVIS formulation. Future developments will focus on a more explicit and tight coupling of the carbon and water fluxes.

L711: We are actually exploring the improvement of both the discretisation of the soil layer as well as the soil parameters that the reviewer refers to. We have modified this paragraph to account for other ongoing developments (see response to main point 7 of this reviewer).

Please also note the supplement to this comment:

<https://essd.copernicus.org/preprints/essd-2021-82/essd-2021-82-AC2-supplement.zip>