

Geosci. Model Dev. Discuss., author comment AC1  
<https://doi.org/10.5194/gmd-2021-155-AC1>, 2021  
© Author(s) 2021. This work is distributed under  
the Creative Commons Attribution 4.0 License.

## Reply on RC1

Petra Pranić et al.

---

Author comment on "Performance of the Adriatic Sea and Coast (AdriSC) climate component – a COAWST V3.3-based one-way coupled atmosphere–ocean modelling suite: ocean results" by Petra Pranić et al., Geosci. Model Dev. Discuss., <https://doi.org/10.5194/gmd-2021-155-AC1>, 2021

---

### Response to Reviewer #1 comments

#### General comments:

*The paper of Pranic et al. entitled 'Performance of the Adriatic Sea and Coast (AdriSC) climate component – a COAWST V3.3-based coupled atmosphere-ocean modelling suite: ocean part', presents an evaluation of the ocean component of the AdriSC climate system against a huge data collection. This study complete a previous evaluation paper that was dedicated to the atmospheric component of the same numerical model run of a 31-year long period (1987-2017). The ocean evaluation is conducted for sea surface, thermohaline properties and circulation.*

My main concern, that I detail below, is about the numerical set-up. In particular I have difficulties to understand the ocean-atmosphere interface, as information about coupling/forcing are clearly missing. See also in the following for some questions about the ocean model itself.

The comparison to observations is fully described and very well presented, even if a small number of conclusions appears rapidly set. Also, some paragraphs are difficult to follow, when describing the subdomains results notably, but this appears inherent of the text insertion of the results put in the supplementary material.

That said, I suggest a minor revision will be useful to improve the paper before accepting its publication.

**Response:** Thank you very much for your detailed and constructive review. In the other sections below, concerns about the model set-up will be appropriately addressed while the text will be changed in order to simplify the reading of the manuscript.

-----

#### Main comments:

##### **Coupling:**

*The first paragraph of section 2.1 seems to stand that WRF 3km and the ROMS models are coupled. In particular, the sentence "Finally, the data exchanges ... are achieved with the Model Coupling Toolkit...". This needs some precision in my opinion.*

*Denamiel et al. 2021 introduced the fact that "the SST from ROMS grid is not prescribed to the WRF models". This means that WRF and ROMS executions are parallel but there is one-way interaction only? If so what are exactly the fields exchanged from WRF to ROMS?*

**Response:** The following sentence will be added to the paragraph in order to address the 2 above comments:

"Additionally, as no ROMS grid was set-up to entirely cover the spatial domain of the WRF 15-km grid, the ROMS sea surface temperature (SST) is not prescribed to the WRF models in order to avoid the generation of discontinuities along the border between the two-way nested WRF 15-km and WRF 3-km atmospheric grids. Consequently, the only grids exchanges in the AdriSC modelling suite consist in the WRF 3-km model providing atmospheric fields (i.e. horizontal wind at 10 m, temperature at 2 m, relative humidity at 2 m, mean sea-level pressure, downward short wave radiations, long wave radiations, rain and evaporation) to the ROMS 3-km and 1-km models which increases the efficiency of the AdriSC model."

*Do you plan to test the two-way coupled mode for future climate simulations? If yes, how will be managed the use of two ocean models? Do you think the dQ/dSST procedure will still be appropriate in the fully coupled run?*

**Response:**

For the RCP 8.5 scenario simulation following the PGW methodology soon to be completed, the one-way coupling has been kept identical in order to be able to compare the past and future simulations. This means that the MEDSEA SST is modified with a climatological change (e.g. increase of temperature up to 3.5 °C in summer) and imposed as boundary condition in the WRF grids as well as SST of reference in the dQ/dSST procedure. Consequently, (1) the WRF models do not benefit from the more accurate calculation of the future SST done in the ROMS models and (2) the tuning of the radiation fluxes is done with an approximated SST of reference. Unfortunately, this is the trade that the authors had to make in order to keep the high-resolution of the ocean models and to be able to conduct their research with their limited numerical resources.

Indeed, in the authors' opinion, as the Adriatic Sea truly needs to be described with at least a 1-km resolution (due to the complex coastline including many islands in its eastern side), the only way to achieve a full two-way coupling between WRF and ROMS models is to set-up a ROMS 9-km grid covering the same domain than the WRF 15-km grid. In this way the ROMS SST could be safely used as homogeneous (i.e. no discontinuities) boundary condition for the WRF grids. However, the AdriSC model is already extremely

slow (1 month of results per day) and the addition of a new grid as well as new grid exchanges would definitely increase the computation time. As our modelling team is composed of only one expert numerical modeller and our numerical resources only consist on the use of the ECMWF supercomputing facilities via Special Project grants, it is unrealistic for us to run such a model.

**The following paragraph will be added in the text after the previous addition already done above:**

**“Ideally, a two-way coupled system would require the use of an additional ROMS 9-km grid covering the WRF 15-km domain. However, due to limited numerical resources and the slowness of the AdriSC modelling suite, such a set-up could not be envisioned in this study. As a consequence, within the AdriSC modelling suite, the WRF models do not benefit from the more accurate simulation of the SST done with the ROMS models even for future scenario runs which only add climatological changes (e.g. increase of SST up to 3.5 °C in summer) to the SST forcing used in the evaluation run.”**

Concerning the dQ/dSST procedure, the authors think it could still be used in a fully coupled system as it just tempers with the solar radiation fluxes which are highly parameterized and not necessarily adjusted to the Adriatic Sea region. However, for scenario runs in the PGW methodology, this means that the SST of reference is just changed with a climatological modification while the real SST used as boundary condition in the WRF models is simulated with the ROMS models. This may thus have some unattended consequences in terms of fully coupled system and thus should be tested.

That said, the dQ/dSST should only be seen as a quick (dirty?) fix and not as a permanent solution. The authors believe that research on, and fine tuning of, the solar radiation penetration in the Adriatic Sea should be carried out in order to properly adjust the parametrizations in this region of the world. As suggested below by the reviewer, the use of ocean colour, turbidity and even the setup of a coupled **sediment transport component are avenues that should definitely be explored in such research. However, as mentioned before, our research group is small and limited by its access to numerical resources. Consequently, the authors did not have the resources (human, financial, numerical) to carry out such experiments.**

The following paragraph will be added in the text after the description of the dQ/dSST procedure:

“It should be noted that the use of the dQ/dSST procedure should not be seen as a permanent solution for climate studies in the Adriatic Sea. Indeed, the SST of reference used in future climate scenario runs is based either on other climate model predictions which are by nature uncertain or on approximations using climatological changes. Consequently, long-term research on the fine tuning and parametrization of the solar radiation penetration using, for example, ocean colour, turbidity or even sediment transport modelling, is thus a prerequisite to a better representation of the coupled atmosphere-

ocean dynamics in the Adriatic Sea.”

**Ocean model:**

- *I understand from run\_coaswst\_model.job0 in the package that the simulation start on 1st January 1987. This must be indicated in the text. Is there any spin-up of the ocean that affect the results for the first simulation years or the use of the MEDSEA re-analysis permits to rapidly have an equilibrium? Does the choice of starting in winter have an impact on dense water formation?*

**Response:** The authors fully agree with the reviewer that the description of the initial conditions and spin-up of the models is missing in the manuscript and the following paragraph will be added after the description of the dQ/dSST procedure:

“Fourth, the AdriSC evaluation run was initialized the 1<sup>st</sup> of November 1986 in order to have a short two-month spin-up period allowing the ocean models to reach a steady state. Indeed, short experiments have shown that rapid equilibrium is reached within the AdriSC ocean models due to (1) the use, before the 1<sup>st</sup> of January 1987, of monthly (instead of daily) MEDSEA v4.1 re-analysis products which have a relatively fine resolution (about 9-km) and assimilate all available data in the Mediterranean Sea and (2) the relatively small size of the ROMS ocean domains. Ideally, several long-term simulations should have been run with different spin-up periods in order to better quantify the impact of the initial conditions on the long-term ocean model results. However, due to numerical resources limitations, such systematic tests have not been carried out with the AdriSC climate model.”

- *I161: Are the river flows distributed homogeneously on the 20 first levels or is there any flow vertical profile?*

**Response:** A vertical profile is applied to distribute the river flow. The following sentence has been modified:

“Additionally, the river flows are linearly distributed between the 20 first sigma vertical levels – i.e. the discharge is multiplied by weights ranging from 20/210 at the surface, 19/210 at the 1<sup>st</sup> sigma level below the surface, to zero at the 20<sup>th</sup> sigma level below the surface.”

- *I162-165: It seems that it could be relevant for the Adriatic Sea to take the sea water colour and turbidity effect on the solar radiation penetration. This is also mentioned in the conclusion. Is there something in ROMS that can be tested or introduced in this direction? Or is there any interest to add the sediment component of COAWST?*

**Response:** see response in “main comments” section.

-----

**Other comments:**

▪ **Introduction**

- *I59: It can be relevant to complete with Carniel et al. (2016)’s study which is more focusing on dense water formation during the 2012 Bora event.*

**Response:** Accepted, the reference will be added.

- **Model, data and methods** (see my main comments)

- 1278: Fig. 1b □ Fig. 1c (for the 7 subdomains)

**Response:** Reference to figures 1b and 1c will be added.

- 1281-282: the sentence is cut

**Response:** Thank you, the typo problem will be fixed.

- **Results and Discussions**

- 1370-372: *In my opinion, the results summary in section 3.1.2 should be separated in two sentences to be fair.*

"... the model is capable to reproduce the BIOS, even though with a weaker intensity due to the overestimation of both seasonal and interannual signals..." "... the SST is quite well reproduced despite presenting a persistent cold bias within the Adriatic Sea."

**Response:** Accepted. Sentence will be split.

- 1400 and 403: *to avoid some confusion (like mine), I suggest to use "the CSP01 dataset" instead of "experiment".*

**Response:** Accepted. "experiment" will be replaced with "dataset"

- *Paragraphs from lines 501-516 and lines 606-616 are difficult to follow. This of course is related to the separation with the supplement material (that is a good option), but if possible, it would be better to find a clearer organization for these two parts either by describing region by region, or by separating temperature results from salinity results for the first paragraph.*

**Response:** The article will be revised in order to clarify and reorganize the text to describe the results region by region.

- 1519: *"... the analysed subdomains and mostly with a good accuracy." For me, you should delete "and".*

**Response:** The sentence will be rewritten as follow:

"In summary, the evaluation of the AdriSC ROMS 1-km thermohaline properties shows that the model is overall capable to reproduce, with mostly a good accuracy, the temperature and salinity in all of the analysed subdomains."

- 1537: *The fact that the highest biases are found around the thermocline and halocline is generally due too smoothed vertical gradients at the ocean mixed layer base. Is this what you obtain in the AdriSC ROMS models? If yes is it equivalent in the 1km-resolution than in the 3-km resolution? Is it somehow related to a similar default in the MEDSEA reanalyses that persists from initial condition or propagates through the open boundaries?*

**Response:** The authors indeed believe that the observed biases found around the thermocline and/or halocline can be attributed to smooth vertical gradients near the mixed layer base. However, to be fair, as the authors did not systematically tested different parametrizations for the vertical mixing and diffusivity in the ROMS models, they cannot discriminate whether the obtained biases are due to the ROMS model setup itself or the

MEDSEA forcing. This is an interesting point that should definitely be investigated in further studies. The following paragraph will be included in the text:

"Additionally, independently of the subdomains, the analysis of the vertical profiles shows that the temperature and salinity biases often present a peak in the vicinity of the thermocline/halocline depth which can probably be linked to an inaccurate representation of vertical diffusivity and vertical mixing in the AdriSC ROMS models. However, more in-depth work should be done to discriminate whether the vertical biases are linked to the AdriSC ROMS model set-up per se or to the MEDSEA fields used to force the initial and boundary conditions."

*-1637-640: "This highlights... the hurricane strength bora winds." This conclusion about the role of wind variability appears quite rapidly set here. Maybe you should precise the resolution of ALADIN? Did you make some tests using WRF 15km to drive the AdriSC ROMS models?*

*There is also a typo in "particular".*

**Response:** The authors forget to put the reference of their work concerning the impact of the resolution of the atmospheric models on the representation of both the bora strength and the sea-surface cooling (using ERA5, WRF 15-km, WRF 3-km and WRF 1.5-km). The paragraph will be modified as follow:

"This highlights that, in the north-eastern Adriatic, higher horizontal and vertical ocean and atmospheric model resolutions, better resolving the complex bathymetry and orography, are required to reproduce the mesoscale variability of the winds and particularly the hurricane strength bora winds as demonstrated by Denamiel et al. (2021a)."

▪ **Conclusion**

*- 1645: To be fair the word "coupled" may be changed. But yes to my knowledge too this is the first time despite several initiatives and still many challenges (see Schär et al. 2020 for instance)*

**Response:** Accepted. Coupled will be removed and the reference will be added as follow:

"In the presented study, the evaluation of the AdriSC ROMS 3-km and 1-km ocean models – forced by the already evaluated AdriSC WRF 3-km model (Denamiel et al., 2021b) – has been carried out for the 31-year long period (1987-2017). The main novelties of the work are, first, the implementation for the very first time – at least to the author's knowledge – of a kilometer scale atmosphere-ocean model for long-term climate studies which still present many challenges (Schär et al., 2020) and, second, the amount of in situ data collected to perform the evaluation of both daily thermohaline (CTD measurements) and hourly dynamical (ADCP and RCM observations) properties of the AdriSC ocean models."