

Geosci. Model Dev. Discuss., author comment AC1
<https://doi.org/10.5194/gmd-2021-73-AC1>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Author comment on gmd-2021-73

Alan J. Geer et al.

Author comment on "Bulk hydrometeor optical properties for microwave and sub-millimetre radiative transfer in RTTOV-SCATT v13.0" by Alan J. Geer et al., Geosci. Model Dev. Discuss., <https://doi.org/10.5194/gmd-2021-73-AC1>, 2021

Both reviewers are thanked for their helpful comments, copied below. Author responses follow an arrow (->)

Reviewer 1

eq 18 is weird: $\hat{\sigma}(\text{Dg})/m(\text{Dg}) \hat{m}(\text{Dg})$ -- looks like the $m(\text{Dg})$ terms would cancel out?

-> This is intentional, with the reasoning described in the text on lines 528 - 534: it is a normalisation used in Figure 6, and Eq. 18 is just Eq. 1 but multiplied and divided by the same factor, $m(\text{Dg})$. A possible improvement may be to add underbraces and captions for the two expressions that are created (the mass-normalised extinction cross-section $\sigma(\text{Dg})/m(\text{Dg})$, and the mass distribution $m(\text{Dg})n(\text{Dg})$).

Radar definition: Lines 855-860: "Here, by definition of the reflectivity factor, the backscatter cross-section is assumed to be given by the Rayleigh scattering approximation ($\hat{\sigma}_b(\text{Dg}) = z_0 D_g^6$)" This simplification shouldn't be necessary, except in that limiting case where it applies. From the particle properties, you have all of the backscattering information necessary to compute the non-Rayleigh reflectivity. Furthermore, for ice-phase hydrometeors, one would have to consider the modifications to the dielectric constant in the "z0" term, unless you're explicitly making the decision that all reflectivity factors are computed *as if* the reflecting hydrometeors are liquid water with an equivalent diameter.

-> The latter is true, all reflectivity factors are computed as if the reflecting hydrometeors are liquid water. The "z0" term is a constant, based on Rayleigh scattering. The backscatter β_b is still generated using the full capabilities of the hydrotable generator (e.g. non-Rayleigh scattering), it is then just scaled by $z_0/10^{18}$ (equation A1). The form of equation A2 was intended as a compact description of the radar reflectivity factor but it seems it is confusing. A proposed remedy is to label the reflectivity defined in equation A2 as $\beta_b^{\text{Rayleigh}}$, and then to better explain that the radar reflectivity is the Z for which β_b (computed using the full scattering capabilities of the hydratable generator) is equal to $\beta_b^{\text{Rayleigh}}$, assuming Rayleigh scattering from a liquid sphere. This is how the observable is reported for the spaceborne radar instruments such as DPR and Cloudsat.

In figure 1, what are the effective diameters of the assumed size distributions for the various hydrometeor types given the fixed water content? Similar comment for Fig. 2, particularly for consideration of the radar reflectivity, which is strongly dependent on the size of the particle.

-> A consideration of the effective diameter (the area-weighted mean diameter, or equivalently the ratio of moments M_3/M_2) could add some useful information. Before creating the revised version of the paper, the effective radii will be calculated, and potentially included in the revised manuscript.

Lines 274-275: "In Fig. 1 the frozen particles have small oscillations with frequency, particularly obvious in the radar reflectivity at lower frequencies. This is a result of interpolating away from the original temperature, size and frequency steps in the ARTS database." This seems more like an error in the interpolation routine. In every other study I've seen that provides similar plots, this behavior is not observed on interpolated points.

-> The interpolation is carried out on the single-particle scattering properties in 3 dimensions, using the trilinear approach. The trilinear interpolation routine has been carefully checked. A first point is that bilinear and trilinear interpolation only give linear results from translation along one of the interpolation axes. Along a transect that is not parallel to one of the axes, a nonlinear function is generated. A second thing to point out is that what is shown are the integrated bulk hydrometeor optical properties, so they are the integrated results of many trilinear interpolations. In figure 1, for example, the temperature is held constant, the frequency is obviously changing (it is the x-axis of the plot) but also the third coordinate of the underlying interpolation (particle size) is also implicitly changing because at different frequencies the bulk optical properties are sensitive to different parts of the PSD. Hence the figure effectively shows a transect along a direction in which both particle size and frequency are changing, where it is not guaranteed to see linear behaviour from the interpolation. This is not the case in figure 2, for example, where both the frequency and the temperature are held constant in the underlying trilinear interpolations, and any variation is along the particle size axis, so this is guaranteed to give linear variations between tie-points in the single-particle optical property database. Admittedly, the manuscript could do better to explain what is actually quite an interesting effect; maybe even a new appendix could be added to give a more complete explanation.

Equation 6: Is D_g equivalent to D_{max} for non-spherical particles, and if how is this handled in the size-integration over various D_{max} values, while maintaining the appropriate mass-dimension relationship?

-> Here, D_g is defined as in Petty and Huang (2011) as the geometric size of the particle, and for non-spherical particles this is the longest dimension of the particle (line 86 of the current manuscript). This is also referred to in some papers as D_{max} , so from that point of view D_g is equivalent to D_{max} as used by other authors. However, in the current work D_{min} and D_{max} are used as the bounds of the integration across the PSD, so D_{max} has a completely different meaning.

With regard to this as well, it wasn't clear to me (perhaps I missed it), but an accounting of the difference in density between bulk ice and liquid water need be made if one is using liquid-equivalent diameter as the baseline for all mass comparisons across hydrometeors. See, for example: Bennartz, R. and Petty, G.W., 2001. The sensitivity of microwave remote sensing observations of precipitation to ice particle size distributions. *Journal of Applied Meteorology*, 40(3), pp.345-364.

-> This is true, but the equivalent diameter is only relevant in a few places in this work (e.g the description of the McFarquhar and Heymsfield (1997) PSD). In plots such as Figure 3 the focus is on frozen hydrometeors and comparisons are not being made between liquid and frozen hydrometeors.

Line 302, units for density are incorrect / inconsistent.

-> thanks for spotting this, it was a mistake and the intended unit was kg m^{-3}

Eq 8 is the := intentional? Also, may want to mention that this is the "k-th moment" equation, and in later equations, "b = k".

-> These are helpful suggestions for making the manuscript clearer; they will be followed in the revised manuscript. However, the := was intentional to indicate a definition of the moment M_k , but on reflection any use of the := will be removed from the manuscript as it might not be 100% clear, and if the equation is a definition this will be clearly stated in the text.

Table 4: SI units?

-> Yes, this will be added to the table caption

Section 3.3.2: You can use the incomplete gamma function solution to definite integrals, this is discussed in Petty and Huang 2011. Not a recommended change, just an observation.

-> Interesting point, this could be useful for the future to avoid renormalisation. A sentence will be added in the revised manuscript.

Line 538, water content is typically expressed in kg m^{-3} ? See e.g., Line 651. Perhaps "water path" would be a more appropriate term when referring to the vertically integrated content (even if for a single layer).

-> Sorry, another typo. The intended unit was kg m^{-3} , and this was a water path.

Lines 550 area: There's some hand-wringing about D_g as the diameter variable of choice, but I think much effort would be saved using D_e instead. The logic is that D_e is a proxy for mass, by virtue of being the mass equivalent radius. This ensures equivalent mass comparisons in the PSDs, and greatly simplifies phase transitions such as melting / freezing. This is, however, not a recommendation for this paper, as it would require a complete overhaul of everything done so far.

-> Agreed. Moving to a D_e basis for the integrations was an aspiration for the v13.0 developments, but there wasn't time to try it out. This is something that could be very useful in the future. However, RTTOV is moving towards a full-spectrum approach in the next few years, it is hoped in future to generate optical properties from the microwave to the visible from the same package. Hence it is worth consideration whether a D_e basis would also suit the higher frequencies where the particle's geometric cross section is more important.

Line 665: "this warming effect" is simply due to the fact that smaller particles are more emissive than scattering at higher frequencies.

-> The text will be improved in this area to reflect this point better, i.e. that the primary

difference between "cloud ice" and "snow" is the chosen PSD, rather than the particle shape, and so indeed it is the fact that particles are in general smaller in the cloud ice category.

Line 669: "Above this frequency of maximum scattering, these clouds start emitting more radiation again and brightness 670 temperatures are higher." True, but it's also relevant to point out that surface-induced polarization effects (e.g., over ocean) ALSO decrease with increasing optical depth -- this provides a separate key piece of information content.

-> Although the use of dual-polarised channels is an important point for the wider information content of microwave radiances, it's not clear whether this refers to the (on average) increasing optical depth from water vapour as frequency increases through the sub-mm, or the point refers to the information coming from dual-polarised channels at a particular frequency. Since the discussion is based on a hypothetical slab cloud with non-polarised optical properties and a fixed warm or cold upwelling non-polarised brightness temperature from below, it is not intended to fully cover all aspects of the microwave and sub-mm information content. Perhaps the best way to address this comment is just to note in the definition of the slab cloud that it does not consider polarisation in any way.

Reviewer 2

... the choice of colors used in the plots: Perhaps red and green should not be used together at the same time in consideration of the colorblind.

-> This is a good point. The figures 2,3,9,10 and 12 do include both red and green lines, but there are so many lines they are hard to tell apart even with typical colour vision. However, the line keys have been carefully devised to follow the order of the lines at a certain point on the figure. Hence it should be possible to identify a line by its relative position on the plot, even without using colour or line style. Figures 1, 4 and 8 may be more problematic, particularly the use of green for the cloud ice category and red for snow. Hence, these will be revised with an alternative colour (or possibly line style).