

Nat. Hazards Earth Syst. Sci. Discuss., author comment AC2
<https://doi.org/10.5194/nhess-2021-403-AC2>, 2022
© Author(s) 2022. This work is distributed under
the Creative Commons Attribution 4.0 License.

Reply on RC2

Kirsty Bayliss et al.

Author comment on "Pseudo-prospective testing of 5-year earthquake forecasts for California using inlabru" by Kirsty Bayliss et al., Nat. Hazards Earth Syst. Sci. Discuss., <https://doi.org/10.5194/nhess-2021-403-AC2>, 2022

Dear Reviewer,

Thank you for your time and thoughtful comments on our article. Below, we address each of your comments.

In this paper, the authors applied the open-source inlabru method to time-independent earthquake forecasts. They used the California region, defined for the RELM experiment, as a case study. The authors described the methodology details in another scientific paper just published. In the first part of the paper, the authors described: i) the spatial models applied in broad terms, ii) the gridded forecast and the synthetic catalog obtained from the method application, and iii) the test applied for the validation of results. For the model construction, the authors examined the relative contributions of the full and declustered catalogs. In the second part, the authors analyze the results obtained with the proposed methodology applying both grid-based and synthetic catalog tests included in the PyCSEP system. Furthermore, they compared the performance obtained with their models with that produced by Helmstetter in 2006 and submitted in the RELM experiment. They concluded that: (i) the full-catalog models performed well in retrospective testing (number, magnitude, and spatial distribution) for the first period 2006-2011 and the results are comparable with those produced by the Helmstetter model; (ii) for the period 2011-2016 the declustered catalog models performed better than the full catalog models, (iii) in the period 2016-2021 the models performed better the N-Test respect to S-Test and the CL-Test and (iv) the simulated catalogs forecasts pass the consistency test more often than their grid-based forecasts.

The paper is satisfactory, and the methodological approach is partially described in the text and referred to another published paper. The data and the code are available for free. The article is well written, and it represents a development in the integration of data from different sources. The use of tests that incorporate grids and synthetic seismic catalogs is also appreciable.

I suggest that the paper must be published after minor revision.

I recommend only a few comments about the paper:

- *Why don't you use the 2005 data as input or in the testing phase?*

The testing phase was chosen to be 2006-2011 to be directly comparable with the original CSEP/RELM testing periods, in particular to allow a direct comparison with the results of Helmstetter et al (2007), which forms a benchmark as the most successful model in that first test. The input data was originally chosen to be 1984-2004 in line with our previous work. There was no intention to miss out 2005 data in both phases.

Nevertheless, we have updated the results so that models are trained on the 1985-2005 time period to avoid any confusion for the reader, though forecasts for both training periods will continue to be available through github and Zenodo. Updating to the new training period does not significantly affect the results, with minor changes in spatial performance observed in the first testing period. We have added results from the 2004-1984 data as supplementary material.

2. The imposition of $b = 1$ for the declusterized catalog probably affected the results obtained. When the catalog is declustered, b tends to be lower than one due to the lack of smaller events. What were the real values of the b -value in the complete and declustered real catalog?

We agree that the choice of b -value for the declustered forecasts may not have been ideal, though we would argue it is not a bad null hypothesis to start with. Given the small size of the catalogues, there is likely to be significant uncertainty in any single b -value used in this way.

While this choice may have had implications for the magnitude test, it should not have adversely affected other tests, and in fact the magnitude test results are acceptable in two out of three time windows despite this potential flaw. We have clarified this in the model description and testing sections of the paper and will investigate this more fully in future work.

While investigating this, we realised that the magnitude test results were incorrect, reporting the wrong quantile scores. We have now updated this in the results section, which results in a better performance for the declustered catalogues in the magnitude tests and a poorer performance for all models in the 2011-2016 testing period.

3. In the tests, you have combined various input data that you had in your possession. Why didn't you test the model with only past seismicity? In this way, it was possible to see how other data contributed to the result. It might also be interesting to see the seismic catalog alone in the synthetic catalog test.

In order to keep to a limited number of forecasts, we chose spatial models that performed best at describing California seismicity according to their DIC in Bayliss et al 2020 to develop to full time-independent forecasts. In this regard, the past seismicity input only is the second-worst performing model in that paper. We chose instead to compare our models to the Helmstetter et al (2007) past seismicity model rather than our own because of the better performance of the Helmstetter model in the RELM tests, and a desire to keep the number of different models to a reasonable size.

4. Figures 3, 5, 6, and 7: insert in the caption the various acronyms (MS, SR, FD, NK, and DC) to facilitate the reader to understand the results;

The figure captions have been updated to include a description of the acronyms.

5. Page 10, line 211: change "Figure 4" to "Figure 5".

Thank you for pointing this out, we have updated the figure number accordingly.

