

Nonlin. Processes Geophys. Discuss., author comment AC2
<https://doi.org/10.5194/npg-2021-6-AC2>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Reply on RC2

Camille Besombes et al.

Author comment on "Producing realistic climate data with generative adversarial networks" by Camille Besombes et al., Nonlin. Processes Geophys. Discuss.,
<https://doi.org/10.5194/npg-2021-6-AC2>, 2021

First of all, we would like to thank the referee for their review of our paper and for giving us the opportunity to improve it.

The structure of the response will be the detailed answers to the questions raised by the referee and then the list of the changes made compared to the preprint version :

Answers to the referees' questions :

RC2-C1 :

I feel that the manuscript can attract more readers by providing more information on the background of the study. How about adding more references about weather generators in the introduction?

A paragraph was added in the introduction that presents the different data-driven methods applied to numerical weather prediction :

"Data driven approaches and numerical weather prediction are two domains that share important similarities, Watson-Parris (2021) explains that both domains use the same methods to answer different questions. This study and Boukabara et al. (2019) also show that numerical weather prediction contains lots of interesting challenges that could be tackled by machine learning methods. It clarifies the growing literature about data driven techniques applied to weather prediction. Scher (2018) used variational autoencoders to generate the dynamics of a simple general circulation model conditioned on a weather state. Weyn et al. (2019) trained a convolutional neural network (CNN) on gridded reanalysis data in order to generate 500-hPa geopotential height fields at forecast lead times up to 3 days. Lagerquist et al. (2019) developed a CNN to identify cold and warm front and post-processing method to convert probability grids to objects. Weyn et al. (2020) built a CNN able to forecast some basic atmospheric variables using a cubed-sphere remapping in order to alleviate the task of

the CNN and impose simply boundary conditions.” (line 33, page 2)

RC2-C2.1 :

The authors evaluated the spatial distribution of the variables simulated by the generator and showed that the generator’s performance is generally good. This is good to evaluate the generator but lacks thoroughness.

The authors should evaluate how well the inter-variable relationships are simulated by the generator ?

We agree with the referee, some of the preliminary studies was done such as verification of the geostrophic balance (Fig 21.), Thermal wind balance (Fig 22, 23, 24.). It would be worth to continue this analysis but it is beyond the scope of our study which is a proof of concept.

We have added to the discussion a consideration of the inter-variable relationships that should be studied in further analysis: Omega equation, other inter-variables balances.

RC2-C2.2 : It is also better to investigate the reproducibility of the vertical structure.

? We have, to some degree illustrated that the approach reproduces some elements of the vertical structure such as Wasserstein distance of variable distributions generated by the neural network with respect to altitude (Fig. 19) and temperature and zonal wind (Fig. 22 & 24) and it would be worth to continue this analysis but it is beyond the scope of our study which is a proof of concept.

Added sentence in the manuscript : *“Authors emphasize that it is necessary to conduct more analysis of the weather situations outputted by the generator which is beyond the scope of this study. For example it would be interesting to assess if other inter-variable balances are present such as the ω -equation, vertical structures etc.”(line 364, page 29)*

RC2-C4 : Similarly, is it possible to give the generator SST fields (or any other boundary conditions (e.g. GHGs, orbital parameters, solar, etc.)) as inputs? For the generator to be used for applications you mentioned in the manuscript (e.g. risk assessment, data assimilation), such information should be included as the inputs.

Using input optimization at inference could be used to condition generation for specific properties in the generation. It is also possible to condition the generations to a specific date in the annual cycle with slight modifications in the network architecture. One could think to condition the output of the generator by a forcing field in input such as forcing field like SST fields for data assimilation application, which should be possible but with more important modifications of the network architecture and a possible impact on the speed of the training procedure.

Possible ways of conditioning our generator (by date, or input forcing fields) were added and the estimation of the changes necessary to be done to the network architecture :

"More advanced strategies, such as training a separate inference network (Chan 2019.) are also possible to apply Bayes' rule without using a particle filter. It is also possible to condition the generations to a specific date in the annual cycle with slight modifications in the network architecture. One could think to condition the output of the generator by a forcing field in input such as forcing field like SST fields for data assimilation application, which should be possible but with more important modifications of the network architecture and a possible impact on the speed of the training procedure." (line 473, page 35)

RC2-C5 : The most important advantage to use weather generators is, I believe, its low computational cost. Therefore, the authors should show how fast the weather generator is compared to the GCM used as the training data.

The GCM simulation was done in 50 minutes on 16 processors and the generator generated 36500 samples (daily sample for 100 years simulation) in 3 minutes on an NVIDIA V100.

The comparison of the computational cost between the GCM and the generator was added to the manuscript : *"The first required property for a weather generator is a low computational cost compared to the GCM that produced the data. Here the simulation with the GCM PLASIM took 50 minutes for a 100-year simulation in parallel on 16 processors, whereas the generator took 3 minutes to generate 36500 samples equivalent to a 100-year simulation on an NVIDIA Tesla V-100." (line 267, page 15)*

Minor changes :

The formatting of the equations 5, 10, 13 was reviewed.

Figure 1 modified in order to specify the layers where strided convolution were used. (see Fig. 1 in supplementary material)

Figure 2 and 3 : Captions were modified to differentiate convolutional and identity blocks. (see Fig. 2 and 3 in supplementary material)

Figure 3 was modified to introduce the parameter s representing the stride value of the convolution that change between different convolutional blocks in the critic architecture. (see Fig. 3 in supplementary material)

Figure 4 : The mention of the upsampling layers, when used, were added on the convolutional blocks. (see Fig. 4 in supplementary material)

Figure 5 : Caption modified to precise that upsampling layer is not in all convolutional blocks. (see Fig. 5 in supplementary material)

The merging of figure 10 and 11 was not done due to a loss of readability of the figures.

Figure 14 was removed because of the redundancy of the information it shows.

Figure 22, 23 and 25 : Descriptions of the panels (a), (b)... were added.

Figure 22 : *"Temperature (K) and zonal wind (m/s) latitude zonals from a boreal winter situation : the thermal wind balance. Left panels correspond to a situation taken from dataset (a) zonal temperature, (c) zonal wind, right panels correspond to a situation taken from generator (b) zonal temperature and (d) zonal wind."*

Figure 23 : *"Thermal wind balance from the boreal winter situation shown in Fig. 22 : (a) sample from the dataset, (b) sample generated by the generator. The temperature (K) is from pressure level 800 hPa and the wind (m/s) from 200 hPa."*

Figure 24 : *"Temperature (K) and zonal wind (m/s) latitude zonals averaged on 30 years subsample. Left panels correspond to a situation taken from dataset (a) zonal temperature, (c) zonal wind, right panels correspond to a situation taken from generator (b) zonal temperature and (d) zonal wind."*

Figure 30 : The equation was referenced instead of described in the caption.

Please also note the supplement to this comment:

<https://npg.copernicus.org/preprints/npg-2021-6/npg-2021-6-AC2-supplement.pdf>