

Interactive comment on “Implications of surface flooding on airborne thickness measurements of snow on sea ice” by Anja Rösel et al.

Anonymous Referee #2

Received and published: 27 October 2020

Review of

Implications of surface flooding on airborne thickness measurements of snow on sea ice

by

Rösel, A., et al.

Summary: This is a very interesting study aiming for a better quantification of the limitations of radar (altimeter) measurements over snow-covered sea ice. Such measurements have been used since two decades to obtain an estimate of the Arctic sea-ice thickness by means of satellite sensors such as CryoSat-2 and since one decade to estimate snow depth on sea ice by means of airborne sensors such as OIB. The crux

Printer-friendly version

Discussion paper



of present-day radar (altimeter) measurements is the unknown penetration depth into a snow cover of unknown depth and physical properties. While dry cold snow is not problematic, deep snow with icy layers interspersed, or wet snow in all of its forms (from surface melting or from flooding at the ice-snow interface) can cause a significant bias in both, snow depth retrieval using the OIB snow radar and sea-ice freeboard and hence sea-ice thickness retrieval using a spaceborne radar altimeter. This study - based on a convincing set of contemporary observations obtained in-situ and from OIB sensors during the N-Ice2015 expedition - quantifies biases in the quantities required for spaceborne sea-ice thickness retrieval: sea-ice freeboard and snow depth, for deep snow (about 50 cm) on relatively thin (150 cm) sea ice with a considerable portion of the ice-snow interface being flooded. Since these conditions appear to be widespread in the Atlantic sector of the Arctic Ocean and the associated peripheral seas and are likely to increase in coverage in the entire Arctic Ocean in the future, the results are of high relevance for accurate retrieval of the sea-ice thickness using satellite radar altimetry.

Given the relevance of the results I recommend to improve the current manuscript along the lines suggested in the following general comments - which are further detailed in the specific comments.

GC1: The study shines through an excellent set of observations. There are parts of the description which ask for improvement, though. One is an improved consideration / discussion of snow depth and sea-ice thickness variations between survey site (local), 5 km circle around Lance (extended in-situ survey) and 10 km circle around Lance (OIB); more details regarding this issue I give in the specific comments. The discussion about scaling and representativity issues is light and could be expanded; elements of this topic I try to express with the following questions. Can one really, as done, combine the OIB data with the in-situ data with such high accuracy? Are EM31 total thickness measurements really that accurate that one can derive highly accurate sea-ice freeboard when combining this data with snow probe snow depth observations? How does

[Printer-friendly version](#)[Discussion paper](#)

the EM31 signal respond to a possibly spatially extended area of flooded sea ice? How does the issue that OIB is known to underestimate thin snow / has issues over highly deformed sea ice influence your results? Finally, the one (or maybe two) ice salinity profiles used at the end to conclude that your observations prove that saline snow can be the main cause for the observed biases, are not very well connected to the rest of the manuscript - even though they are the dominant topic in the discussion and conclusion section. I recommend to give the description of these in situ observations more weight and demonstrate more clearly how well (hopefully) these single point measurements represent the conditions in the detailed survey area.

GC2: My impression is that the MAIN issue is that OIB data result in a sea-ice freeboard bias of 0.2 m. This is a SUBSTANTIAL bias but it is not presented and discussed in an overly prominent way in the manuscript. I recommend to be more exhaustive in the discussion of particularly these results in the context of Figures 6 through 8.

GC3: The secondary issue is the snow density of the proposed () two-layer snow setup which is discussed in the context of Figure 8, hi4. There are, to my opinion, at least two areas of improvement. The first one is superposing the drill-hole sea-ice freeboard data onto Figure 7 and discuss the results. The second one is to conduct a sensitivity study which plays around with possible snow layer depths and densities used in Equation 6 to derive hi4. A pre-requisite for these actions is an appropriate introduction of what you can the two-layer snow setup, which is not adequately described yet in the manuscript. In this context, I also kindly ask for clarification of the wording "slushy basal snow layer" versus "snow-ice basal snow layer" because snow-ice is refrozen slush hence hard while slush is soft - which has implications for the snow probe measurements and the interpretation of the measurements.

Specific Comments:

Line 179++: I am not sure your choice of denoting different variables with 1, 2, 3, 4, using hi (I) and hs (S) and hfb (Fi) and hfbs (F) to name the variables, i.e. without

[Printer-friendly version](#)[Discussion paper](#)

subscripts, and introducing subscripts such as EM, SP or IS is an optimal solution. In any case I recommend to use it in a consistent way. This means that also in the running text hi and hs should be given as used in the formulas (see Lines 174/175 for instance). In addition, I am wondering whether it is necessary to abbreviate "in situ" with IS. To me this is confusing in the zoo of short names. But this is of course your choice. If you keep IS then you need to move its definition from Line 194 to Line 179.

Equation 2: I am sure that the freeboard in this equation needs to be the snow freeboard, i.e. hfbs.; this is the classical equation to derive sea-ice thickness I from total (sea ice + snow) freeboard F and snow depth S, isn't it?

Then equation (3) would be the total freeboard as well and not the sea-ice freeboard. In order to end up with the sea-ice freeboard here I suggest to use the retrieval equation used for radar altimetry (in your notion): $hi1IS = (hs1SP * rho_s + hfb1IS * rho_water) / (rho_water - rho_ice)$

In Line 194 it needs to be "1" instead of "2" in the variable name.

Equation 4: In the context of this equation I'd like to note that you are not consistent with Figure 3. There you denote snow freeboard as hfbs while in Equation 4 and in Line 202 you write hsfb. I like hsfb more.

Line 206/207: While the accuracy for the drill-hole measurements is clear from the stated measurement accuracy of the measurement device, I am wondering whether you might want to add one sentence about the way you estimated the value of 0.06 m for the combination of snow probe and EM31 measurements.

Equation (7): I suggest to split this equation into two. Please first introduce hB without the interpretation that it can be decomposed into sea-ice freeboard, basal snow-layer thickness and an error term and subsequently, in a follow-on equation (if necessary) show the decomposition of hB. I note, after having read the entire manuscript, that there is limited reference to and usage of this equation. Or did I overlook a figure or

[Printer-friendly version](#)[Discussion paper](#)

table where you provide an estimate of the basal snow-layer thickness?

Figure 4: I am a bit confused with what I see here. It appears to me that this plot starts on the left somewhere close to the top right corner of the map shown in Figure 2, taking one of the overpasses shown towards the southwest (crossing the survey area). Fine. Then comes the aircraft turn; the main part of the radar-echogram shown in Figure 4 is related to sea-ice conditions "on ice floe south of the survey site". Now, what I have problems with is the fact that you stated the times for overpass #2 and #3 as 15:37 and 15:43 in the caption of Figure 4 while in the heading line just above the figure it specifies a time range around 15:43, suggesting that this is only overpass #3. Could it hence be that we only look at overpass #3 and that only the small part (left quarter of the echogram) is coincident with the tracks (actually only overpass #3) denoted in Figure 2? If this is the case, I am wondering whether it wouldn't make sense to expand that left quarter of the echogram because this has a direct relation to the survey area.

Lines 253-255: "In addition ..." How important is it to explicitly mention the mean of these very few (compared to the other samples) snow depth observations at the drill hole sites in the text? Would it be sufficient to only show this value in the Table? I am suggesting so because you are coming up with quite a number of different snow depth values and it begins to become confusing. Particularly, since you repeat this information in Line 273.

Line 263: What is the motivation to draw a 10 km circle around R/V Lance for the air-borne data when the in-situ observations along the five transects were carried out within a 5 km radius circle?

Line 275/276: "Three ... before drilling." → How do you know?

Figure 5: In the left panel you denote hs1 and hs3 with the respective methods while in the right panel you write "regional". My suggestion would be to be consistent. At this point the reader possibly knows that hs1 is based on snow probe data and that hs3 is based on the snow radar. Hence you could use "survey area" in the left panel and

[Printer-friendly version](#)[Discussion paper](#)

keep "regional" in the right panel.

Lines 294-298: These sentences about brine wicking etc. fall a bit from heaven. I suggest to start a new paragraph here and motivate these further considerations by again stressing that near-zero ice-freeboard supports flooding of the ice-snow interface and subsequent upward wicking of seawater and/or brine into the basal snow cover. The paragraph about the c-profile of the salinity given at the beginning of Section 3 could be placed here in a much more logical way - as this piece of information seems a bit lost where it is located currently.

Figure 7: I suggest to overplot the freeboard values shown in Table 2 of the appendix onto the maps in each panel by using, e.g., a color-filled circle, and discuss what you see.

Lines 303-313 / Figure 8:

- Line 305: I don't find hi_REGIONAL in Figure 8.

- Line 307: "slightly thinner" → It might be a matter of taste but a difference of 0.3 to 0.4 m in a thickness range between 1.1 and 1.5 m I would not call "slightly". In addition, those 5 regional survey lines, were these laid out without checking the ice conditions beforehand? What I mean by this is, that based on the large sea-ice thickness standard deviation it seems likely that these 5 lines had a substantial fraction of thin ice from re-frozen leads while the bulk of the sea ice inbetween might have had a similar thickness than the ice of the survey site. Please check and, if need be, re-phrase statements.

- I am missing commenting on hi3.

- Line 309-313: I suggest to re-write this part. hi4, if computed using Eq. 6, uses the sea-ice freeboard (hfb) which is derived from airborne ATM total freeboard (accurate) and the in-situ snow depth (accurate as well); in addition, for the snow part it uses (again) the in situ snow depth. The densities you used are those which you measured in the field = accurate. Hence, at first glance it is not clear why even with the densities

[Printer-friendly version](#)[Discussion paper](#)

you measured the h_{i4} is biased compared to h_{i1} and h_{i2} . The statement that the bias in sea-ice thickness between h_{i1} and h_{i4} of about 0.4 m is consistent with the bias in h_{fb} of 0.03 m and that this is in part due to the not sufficiently well considered "two-layer snow set up" is not backed-up yet by your writing or the figures. If, as you suggest, snow densities across this two-layer snow set up are the main cause, then I strongly suggest to play around with potential snow densities (you measured some in the field, didn't you?) and layer thicknesses to figure out whether your hypothesis is true. In other words: Which snow density values of the proposed two-layer snow setup explain the observed difference between h_{i1} and h_{i4} ? Are these realistic?

Discussion section:

General comments: I don't see a reason why to write "total snow depth" or "total snow thickness". I guess "total" can be omitted.

Line 345: This is a reminder that it might make sense to provide a scientific motivation for using a 10 km radius for the airborne data versus a 5 km radius for the in situ data. If by chance the airborne data observed a comparably large fraction of thin ice, then the airborne snow depth would naturally be smaller and hence its bias to the in situ snow depth.

Line 357: "slushy, snow-ice formation in the basal layers of the snow pack" vs. Line 367: "formation of highly saline and saturated slush in the basal snow layers" → It is a difference whether one speaks of wet / saturated slushy snow ... which is a rather soft material, or whether one speaks of snow-ice which - at least to my understanding - is refrozen ... and hence hard. It is, to my opinion, important to distinguish between those because I'd think that the SP measurements would penetrate slush and hence INCLUDE the thickness of the slush layer into the snow depth reading while these would not penetrate snow ice and exclude that part from the snow depth reading. Please be clear what you mean and observe to avoid misunderstandings.

Line 363: "the 1-m thick FYI floe" → In lines 241-244 you already provided some in-

[Printer-friendly version](#)[Discussion paper](#)

formation about these observations. I note that these are not coherent and not specific enough. What did you mean by "in the vicinity of the 2D site" in those lines? There you gave one date (March 5), here you give two dates. The thickness of that floe (is it representative?) is just 1 m, i.e. substantially less than the surveyed floe with 1.4-1.5 m thickness. I suggest to comment in your manuscript about these differences and slight misfit in information. I also interpret from your writing that snow depth (on that single floe) increased by 0.08 m between March 5 and March 23 and that that snow depth is considerably smaller than the average or modal snow depth of the 2D survey site. Comments?

Lines 379-382: I have a stupid question in this context: How do you compute the sea-ice thickness from estimates of sea-ice freeboard and snow depth using the classical equation (e.g. equation 6) when the sea-ice freeboard is negative? Then the first term in equation 6 is negative. Is the retrieval using that equation defined at all?

Line 383++: "Our study shows that saline snow conditions can ..." → I suggest to formulate more clearly how the conditions met in this study differ from those encountered on Canadian Arctic fast ice. There the snow cover was dry, the sea-ice freeboard positive and the brine concentration in the basal snow layers solely caused by the high sea-ice salinity near the ice-snow interface. Here, during N-ICE2015, the situation appears to have been completely different, with a substantial amount of negative sea-ice freeboard, hence flooding of the ice-snow interface and an (unknown?) amount of slush at the ice-snow interface from which large amounts of brine can be wicked up (how high?) into the overlying snow.

Line 407: Please explain why there are two different snow thickness values (here in the text and also in Fig. 9 a)

Line 408: I suggest to add a statement about the fact that the survey cite is located in an area where the shown CS-2 products (and also the snow depth) show large spatial variability. In light of the fact that sea ice is not static but drifts, your "verdict" about

[Printer-friendly version](#)[Discussion paper](#)

the quality of the CS-2 product could perhaps be formulated in a less harsh way. I note in this context, that you completely ignored the CPOM results published by Tilling and co-workers, and issue which I kindly ask you to amend in your manuscript.

Typos / editorial remarks:

Line 24: "wicking and saturation into" → "wicking into and saturation of". Later in the sentence: Would it be sufficient to write "causing the airborne radar signal ..."? I would read more fluently. If "more diffuse scattering and influenced" shall be kept then I suggest to split the sentence into two.

Line 33: "it may result ..." → I am not sure I understand what this refers to.

Line 43: You could add a sentence stating the importance of the thickness of the snow layer on sea ice on the in-ice and under-ice biological processes.

Line 73: I suggest to cite the work of Willat et al. (2010) here, also from the Antarctic but a different region: Digital Object Identifier 10.1109/TGRS.2009.2028237

Line 85: Perhaps switch "Atlantic Sector of the Arctic Ocean" and "Southern Ocean" since your primary focus is in the Arctic Ocean?

Lines 112/113: "... accuracy ... higher ..." → I know what you mean but a reader might stumble at first glance being surprised that the EM31 accuracy is "better" for rough and deformed ice - which, I guess, quite some readers automatically imply into "higher". Perhaps it might make sense to write "worse"?

Lines 122/123: "We use the results of the independent snow transects from Floe 2 to provide the regional context ..." → I am a bit confused. You have that 400 m by 60 m survey area on the floe. And then you have those additional (>5000) measurements within 5 km of the ship. These are - at least this is my assumption - not necessarily on floe 2. But these are the measurements which provide the regional context, am I right?

Figure 2: What is "WAV snow depth"?

Printer-friendly version

Discussion paper



Line 153: "During the survey ..." → Does this refer to the OIB survey or to the ground-based survey?

Line 161: "..., and the ..." → delete "and"

Line 171: Any measurements of the density of second-year ice?

Line 178: "for flooding" → "to flooding"

Line 190: "can calculated" → "can be calculated"

Line 250: "second mode" → "mode"

Line 254: "0.08 m than" → "0.08 m larger than"

Lines 263-265: "with the ... of the ship" is kind of a repetition of the end of the previous paragraph. I suggest to simply refer to the above-mentioned transects.

Line 273: "(FB2)" can be deleted, I guess.

Line 287: "of+-0.06 m" → two spaces are missing

Line 290: "Figure 5" → "Figure 6" "lie in the negative range, to -0.1 m." → "are negative with magnitudes up to 0.1 m."

Line 291: "Results in the same range, with ... 0.09 m, are obtained ..." → "Results in the same range are obtained ... elevation, resulting in an average value of hfb4 ... 0.09 m (FB4, see Figure 6)."

Line 296: "... elevation value" → a good place to refer to equation (7).

Line 305: "asfor" → "as for"

Lines 305/306: "calculated from a combination ..." → can you refer to one of the equations? Also, you used "SP" to denote the snow probe measurements repeatedly and can do so here as well.

Figure 7, caption: Line 326: This second "SR" needs to be "SP". I suggest to add a

[Printer-friendly version](#)[Discussion paper](#)

note that the annotation of the color bar is non-linear.

Line 356: "pushes" → "can push" as this is a function of ice thickness. I also strongly recommend to split this long sentence into at least two sentences.

Line 381: "of vertical shift" → "of a vertical shift"

Line 382: "... horizon, caused ... freeboards." → perhaps better: "... horizon, caused by slush above the ice-snow interface associated with the negative freeboard and additional wicking up of brine into the overlying snow."

Line 391: "from the sea ice surface" → "from the sea ice or slush surface"

Interactive comment on The Cryosphere Discuss., <https://doi.org/10.5194/tc-2020-168>, 2020.

[Printer-friendly version](#)[Discussion paper](#)