

The Cryosphere Discuss., referee comment RC1
<https://doi.org/10.5194/tc-2021-120-RC1>, 2021
© Author(s) 2021. This work is distributed under
the Creative Commons Attribution 4.0 License.

Comment on tc-2021-120

Anonymous Referee #1

Referee comment on "Quantifying the potential future contribution to global mean sea level from the Filchner–Ronne basin, Antarctica" by Emily A. Hill et al., The Cryosphere Discuss., <https://doi.org/10.5194/tc-2021-120-RC1>, 2021

The following is a review of "Quantifying the potential future contribution to global mean sea level from the Filchner-Ronne basin, Antarctica" By E. A. Hill, et al.

This manuscript describes a set of experiments using the ice flow model *Úa*, aimed at characterizing the spread of possible future sea level contribution from the Filchner-Ronne (FR) drainage basin. The authors create a set of ensemble simulations, under various RPC emission scenarios, that extend to the year 2300, and use these simulations to build a surrogate model. To create their ensemble, the authors sample parameters related to ocean forcing, atmospheric forcing, and ice dynamics within bounds they derive from literature, climate model ensembles of future projections, and Bayesian analysis (for the ocean forcing parameters). The authors illustrate that the surrogate model exhibits skill in predicting the sea level contribution projected by *Úa* in the FR region by year 2300, and they then use the surrogate to create a million-sample distribution to represent the possible spread of sea level contribution from FR within the same period. Surrogates are also created to derive estimates of sea-level contribution at years 2100 and 2200, for analysis of projected sea level contribution through time. Results suggest that the FR region is not likely to contribute positively to sea level contribution in the future, largely due to the modeled increase in accumulation over time in response to warming atmospheric conditions. However, significant contribution to sea level from this region is found to be possible (more than >30cm by year 2300), though this outcome is not probable. The author's analysis also allowed them to isolate contribution to uncertainty from the various model parameters sampled, and they find that atmospheric and ocean forcing account for the majority of the sea-level projection uncertainty, in agreement with past studies. The authors specifically highlight these model boundary conditions as the most important to improve models of in order to increase confidence in ice sheet model projections of sea level contribution.

Here, the authors present a novel approach to the challenge of quantifying uncertainties in ice sheet model projections. Running the number of model simulations required for robust assessment of uncertainties is, in many cases, not computationally feasible, so the design of an adequate surrogate model for this purpose is highly advantageous. Overall, I find that the authors thoroughly describe their methods, experiments, reasoning, and caveats. The discussion, in particular, highlights the care that must be taken when considering results from a single ice sheet model where specific assumptions are necessary to produce realistic model ensembles. The manuscript is well-written and the figures are highly illustrative of the methods and results. The workflow diagram (Fig. 2), is especially helpful in describing the investigation's strategy. In addition, the results are thorough and well-organized, therefore I find this manuscript highly appropriate for publication in the Cryosphere, with revisions and some supporting analysis.

I have a number of questions and comments for the authors, as listed below, for author response and discussion.

General comments:

Discussion section – The discussion is quite thorough, and you cover many important points and caveats. However, I think it would be improved if you also expand upon some interesting topics that are brought up in the results section, specifically pertaining to the advantages and disadvantages of using a surrogate model for this analysis. For example, in the results section, you note an example of an advantage of using the surrogate, can you expand upon this in the discussion to talk about what you learned in that exercise with respect to the importance of including extremes in your training set? Could you also expand upon what might be the disadvantages or pitfalls that others using your methods could encounter? (For example, is there a danger of not capturing threshold behavior or runaway retreat, as you observe in some of your extreme forcing simulations?) Is it possible that runaway retreat is more likely than your training set suggests, or do you think that your sampling space and final pdf capture the spread of possible scenarios accurately?

In addition, how important was it to use a surrogate to capture the full sample space? That is, do your final pdf's reveal a different pdf than your ensembles suggest? Perhaps

you can show some training run (ensemble) pdf's vs. the surrogate sampling pdf's in the appendix to illustrate this point.

Specific questions and suggestions:

Line 1: This opening sentence is a bit awkward to read. Maybe adding "The future ..." => "change", "behavior", or "evolution" or a similar phrase would make your point clearer.

Line 34: ... as the "combined area" of the two major drainage basins... (or something similar)

Line 55: I understand the point you are trying to make in this sentence, but it reads awkwardly. Please try rephrasing.

Line 159: Could you add a thin or dashed line to Fig 1 or its small inset that shows where the divide between the two basins sits?

Line 162: Perhaps in the supplement, an illustration of what your mesh looks like, perhaps in some key locations, would be very helpful.

Line 170: Could you please specify the settings for your initial mesh adaption, for instance: What is your minimum mesh for the adaption? Is it still 900m? Is there a maximum mesh size near the grounding line? How close to the grounding line is the adaption imposed (i.e. is there a set buffer length)?

Line 171: What is the minimum thickness value that is imposed?

Line 235: You discuss only changes to surface accumulation through time. Do your simulations have a representation of surface melt as well (i.e., a PDD scheme or something similar)? Please specify this in the text.

Line 315: The term "point" is used a number of times in the text to refer to a location in your sampling space. This terminology is easy to confuse with a point in map space. Is it possible to use a more specific term, like "sample point" or something similar throughout the manuscript to distinguish between sampling space and actual 2d map space?

Line 322: Please quantify the model drift (or spread of model drift for all of your control runs) here.

Line 328: Throughout the manuscript, you refer to this set of simulations as your "ensemble". It would be helpful for the reader if you explicitly call out here that these are the runs that you will hereafter call the "ensemble" (or create a name for this set of runs

that you can refer to later in the manuscript).

Line 339: Please add a quantitative statement with respect to the surrogate model being “close” to the ice flow model response.

Line 345: What type of algorithm is used for the sampling of these 1 million simulations? Is it Latin hypercube as in the other ensemble sampling?

Figure 5: Please note the year at which the sea GMSL represents in the plot or caption (i.e. 2300).

Line 440: Please note in the text the simulations used for this analysis (i.e., the ensemble)

Line 530: While Ritz et al., 2015 do not change surface mass balance through time, Schlegel et al., 2018 apply a step function on accumulation, so there is still a possibility for suppression due to accumulation. The difference is more likely due to your treatment of ocean forcing (e.g., PICO with Bayesian exploration extreme melt rates, which may be lower but considered more realistic, especially with consideration to the possible time lag between atmosphere and ocean warming), as well as your application of no melt on partially floating elements, adaptive grounding line mesh, and even the repetition of inversion procedures for each model simulation.

Line 535: Your results show a strong dependence on accumulation, and the discussion below gives an inclusive overview of the challenges in forcing accumulation on ice sheet models in general. Could you offer a quantitative comparison between the spread (pdf) of the change (anomaly) in precipitation that your sampling imposes on your simulations, and that that is predicted by CMIP5 models? For example, you present the spread in the ocean forcing time delay parameter from LARMIP-2, could you show similarly, maybe in the supplement, a similar pdf for your regional accumulation from various CMIP GCM's and then compare that against a pdf that represents accumulation sampling from the surrogate (or even just the ensemble?). I am curious to see this comparison, since while you sample p in a normal way (Fig. 3), this parameter is exponentially related to accumulation. Is it possible that this choice skews your sampling to higher sensitivity to temperature change? Is the imposed sampling of total increased accumulation realistic as compared to what GCM's are predicting? If not, this should be mentioned as a caveat in your discussion, because it has implications for the shape of your final GMSL pdfs (Fig. 5), and total probabilities.

Line 764: Could you quantify at what probability does the Reese et al., 2018a value sit in your distribution? Can you comment on the possible implications of your values sitting much lower in magnitude than suggested by this previous study? In Fig. S2 it appears that once the \square_{τ}^* value approaches 2×10^{-5} , that the delta GMSL contribution starts to become highly negative. Can you discuss why that is and how it influences your resulting pdf from Fig. B1?

Line 766: basal mass balance?

Figure S1: Please specify the simulation year of GMSL presented.

Figure S2: The results presented here are a bit puzzling. Are they, as other results, represented as difference from the control simulation? If so, if I am understanding your methods correctly, then I would expect all graphs to cross a value of zero Δ GMSL, when the value of the sampled parameter is equal to the value of the control run. Could you speak to this, specifically why most of the graphs show a negative contribution to sea

level for all of the sampled values?

Figure S4: Please check the lettered labels on the right-hand column plots.