

The Cryosphere Discuss., referee comment RC1 https://doi.org/10.5194/tc-2021-185-RC1, 2021 © Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.

Comment on tc-2021-185

Anonymous Referee #1

Referee comment on "Kara and Barents sea ice thickness estimation based on CryoSat-2 radar altimeter and Sentinel-1 dual-polarized synthetic aperture radar" by Juha Karvonen et al., The Cryosphere Discuss., https://doi.org/10.5194/tc-2021-185-RC1, 2021

The study "Arctic Sea Ice Thickness Estimation Based on CryoSat-2 Radar Altimeter and Sentinel-1 Dual-Polarized SAR" by Karvonen et al. puts forth a method to interpolate and extrapolate CryoSat-2 sea ice thickness data over the Kara and Barents Seas using segmented Sentinel-1 SAR data. Currently, daily-scale sea ice thickness data from altimetry include only sparse, footprint-sized swaths that are less useful for timely purposes such as maritime navigation. This method aims to increase the utility of daily CryoSat-2 thickness data by mapping the CryoSat-2 thicknesses to coincident Sentinel-1 SAR segments to create high resolution sea ice thickness maps over a given area.

This manuscript defines the proposed algorithm, explains the inter-/extrapolation of the CryoSat-2 data to neighboring SAR segments, and compares the results to different sea ice thickness products, namely the AARI ice charts and two different sea ice reanalyses. An initial high bias was found in the data, so the CS2/S1 sea ice thicknesses were remapped using another reanalysis to reduce the positive bias. While the comparisons show low-to-moderate agreement of mean thickness values over the study area, the authors put forth some discussion on potential difficulties in the comparisons as well as potential future avenues for improvement of the algorithm, so that it may be useful for future altimetry missions.

Overall, the technique presented here is novel and potentially useful, and the manuscript shows some interesting results. However, I feel the results could be strengthened by incorporating some of the comments below. Additionally, there are many grammatical errors that need to be addressed, pointed out in the technical corrections. Provided that these (at times substantial) revisions are made, I would recommend this manuscript for publication in The Cryosphere.

General Comments:
GC1: While the introduction does a good job to provide background on different retrieval techniques and limitations, I feel that it would benefit from:
 Including some applicable references that are missing (see specific comments below) Focusing the topics a bit more. The last paragraph, for example, talks a lot about the relationship between roughness and backscatter, which is not a topic covered in the rest of the paper. Providing an outline at the end that ties the introduction to the rest of the paper and outlines what will be presented in the manuscript
GC2: I feel it would be useful for this study to provide some assessment of the retrieved thickness maps with other remotely-sensed datasets. I understand the desire to compare with independent daily-resolution products (which are lacking from altimetry), however, the reanalyses simply don't capture the observed thickness distribution with enough confidence to draw robust conclusions about the performance of this algorithm.
To me, the merged CryoSat-2/SMOS product (Ricker et al. 2017; https://doi.org/10.5194/tc-11-1607-2017) would be a useful comparison, as the same input data are used (CryoSat-2) so the authors can test how well these data are mapped

To me, the merged CryoSat-2/SMOS product (Ricker et al. 2017; https://doi.org/10.5194/tc-11-1607-2017) would be a useful comparison, as the same input data are used (CryoSat-2) so the authors can test how well these data are mapped to S1 compared to the SMOS product. I understand this is a weekly product, however, I think it would be as (if not more) useful than e.g. the AARI ice charts (which are also a weekly product at a lower spatial resolution) by providing sub-polygon thickness information.

GC3: While this paper provides plenty of metrics comparing the study-area-mean thickness between the CS2/S1 and the reanalyses, I feel it does not adequately compare the retrieved spatial pattern of thickness over the study area. While the average values show moderate agreement in some cases, there is a clear discrepancy in the spatial patterns that warrants discussion. Difference maps, for example, would help to show which areas (and therefore which ice types or texture features) show better or worse agreement. Such analysis could also help to better inform the need for remapping, as the

bias may not be uniform over the region.
GC4: Parts of this study region have been found to have large snow loads that cause negative ice freeboards and wet/slushy snow-ice interfaces (Rösel et al., JGR Oceans, https://doi.org/10.1002/2017JC012865). While this paper mentions impacts of wet snow surfaces brought on by warm surface air temperatures, I would be curious to see the impacts of wet snow (and potentially slush) near the snow-ice interface mentioned in this manuscript.
GC5: Overall, the manuscript could use a closer proofread, as there are many issues with abbreviations, parentheses in citations, and grammar that partially distract from the science. Many of these are pointed out in the specific and technical comments below.
Specific Comments:
Title: While the title is descriptive and technically accurate, the method is only applied to a small area of Arctic sea ice and not tested over the whole Arctic. Therefore, I feel it may be best to modify the title and specify that the study focuses on the Kara/Barents Seas.
Lines 9-18: Appropriate references are missing from this first paragraph.













