

Metabolic tradeoffs and heterogeneity in microbial responses to temperature determine the fate of litter carbon in simulations of a warmer world

Grace Pold^{1*}, Seeta A. Sistla², and Kristen M. DeAngelis³

¹Graduate Program in Organismic and Evolutionary Biology, University of Massachusetts, Amherst, MA 01003, USA

²Natural Resources Management and Environmental Sciences, California Polytechnic State University, San Luis Obispo, CA 93407, USA

³Department of Microbiology, University of Massachusetts, Amherst, MA 01003, USA

Correspondence: Grace Pold (apold@umass.edu)

Abstract. Climate change has the potential to destabilize the Earth’s massive terrestrial carbon (C) stocks, but the degree to which models project this destabilization to occur depends on the kinds and complexities of microbial processes they simulate. Of particular note is carbon use efficiency (CUE), which determines the fraction of C processed by microbes that is anabolized into microbial biomass rather than being lost to the atmosphere as carbon dioxide. The temperature sensitivity of CUE is often modeled as an intrinsically fixed (homogeneous) property of the community, which contrasts with empirical data and has unknown impacts on projected changes to the soil C cycle under global warming. We used the DEMENT model—which simulates taxon-level litter decomposition dynamics—to explore the effects of introducing organism-level heterogeneity into the CUE response to temperature for decomposition of leaf litter under 5°C of warming. We found that allowing CUE temperature response to differ between taxa facilitated increased loss of litter C, unless fungal taxa were specifically restricted to decreasing CUE with temperature. Increased loss of litter C was observed when the growth of a larger microbial biomass pool was fueled by higher community-level average CUE at higher temperature in the heterogeneous microbial community, with effectively lower costs for extracellular enzyme production. Together these results implicate a role for diversity of taxon-level CUE responses in driving the fate of litter C in a warmer world, which should be explored within the framework of additional model structures and empirical studies.

1 Introduction

Soil heterotrophs are central to the cycling and recycling of the 60 Gigatons of organic carbon (C) that plants deposit onto and into the ground each year. How well these litter inputs are converted into relatively stable soil organic matter depends on temperature, moisture, chemical composition, and soil mineralogy, which interact to influence microbial physiology (Manzoni et al., 2012; Kallenbach et al., 2016; Oldfield et al., 2018). Predictions regarding how soil C stocks will respond to climate change are, in turn, highly sensitive to how carbon use efficiency (CUE)—or the fraction of C taken up by a cell and incorporated into biomass rather than being respired—changes with temperature (Allison et al., 2010; Wieder et al., 2013; Allison, 2014; Li et al., 2014; Sistla et al., 2014; Tang and Riley, 2015). As such, quantifying microbial decomposer CUE and its responsiveness to environmental change has been subject to intensive study (Devêvre and Horwáth, 2000; Frey et al., 2013; Blagodatskaya et al., 2014; Lee and Schmidt, 2014; Spohn et al., 2016a, b; Öquist et al., 2017; Malik et al., 2018; Geyer et al., 2019; Malik et al., 2019; Zheng et al., 2019).

Soil microbial communities show considerable differences in how their metabolisms respond to elevated temperatures, with their CUE increasing (Öquist et al., 2017; Zheng et al., 2019), decreasing (Devêvre and Horwáth, 2000; Frey et al., 2013; Öquist et al., 2017; Li et al., 2018; Zheng et al., 2019) or remaining unaffected by warming (Dijkstra et al., 2011b; Öquist et al., 2017; Walker et al., 2018; Zheng et al., 2019). However, models of the soil C cycle generally assume either no change (Allison et al., 2010; Li et al., 2014; Wieder et al., 2014) or a fixed decrease in CUE with temperature (Allison et al., 2010; Wieder et al., 2013; Allison, 2014; Li et al., 2014). When CUE is allowed to directly increase with temperature, this temperature response is fixed across taxa (Frey et al., 2013; Ye et al., 2019). In other instances, CUE may be modeled as fixed within taxa, such that dynamic changes in community-level CUE with warming are the result of shifts in the dominant group or groups of organisms present as a function of their dietary preferences and/or C: nitrogen (N) ratio (Wieder et al., 2013; Sistla et al., 2014), rather than any inherent differences in the temperature sensitivity of the constitutive community members. This may occur, for instance, if large C-rich fungi show less-positive responses to warming than small, N-rich bacteria do (DeAngelis et al., 2015). Therefore, models have thus far insufficiently accounted for how the temperature sensitivity of central metabolism may differ between microbes, such that intrinsic differences in efficiency between taxa ("heterogeneity") above and beyond temperature-driven differences in substrate supply may also drive microbial community trajectories.

Heterogeneity in the temperature sensitivity of growth efficiency across taxa could be driven by differences in the rate-limiting step of central metabolic pathways (Dijkstra et al., 2011a), or in how well the proteins responsible for the extracellular processing and uptake of environmental nutrients are able to maintain activity as temperature increases (Allison et al., 2018; Alster et al., 2018). For instance, there is some evidence that bacteria benefit more than fungi from an increase in temperature, as their growth rate was observed to decrease less rapidly with temperature above its optimum than a fungal community in an incubation study (Pietikäinen et al., 2005). The respiration rate for the two groups could not be isolated in that study, and so differential effects of temperature on CUE could not be parsed out. However, the CUE has been observed to differ in how sensitive it is to nutrient limitation for bacteria and fungi, such that differences in the temperature sensitivity of CUE between the two groups may play out as a consequence of changing nutrient demands (Keiblinger et al., 2010; Sinsabaugh et al., 2016).

50 Therefore, the temperature range over which an an organism can maintain efficient growth is one important dimension of its niche (Cavicchioli, 2016), and, when combined with other “response traits” determining how they react to the environment (Lavorel and Garnier, 2002), can impact their success in the environment.

These microbial response traits may in turn be linked to “effect traits” determining how an organism alters its environment, such as extracellular enzyme production (Martiny et al., 2013; Treseder and Lennon, 2015; Amend et al., 2016). For instance, 55 taxa capable of growth at higher temperatures may need to produce a broader suite of enzymes and attack a wider range of substrates to support this rapid growth than slower-growing taxa. Extracellular enzyme production may impose substantial metabolic costs on the cell because C which could otherwise be allocated to growing the cell must instead be spent producing amino acids and expending ATP to synthesize the enzymes (Kaleta et al., 2013; Kafri et al., 2016). As such, extracellular enzyme production is inferred to reduce the CUE of soil microbial communities (Allison, 2014; Malik et al., 2019).

60 We explored whether interactions between the temperature sensitivity of intracellular (i.e. CUE) and extracellular (i.e. litter decomposing enzyme) metabolic processes of cells can explain why CUE is observed to increase with temperature in some soils, and decrease in others. We used the litter decomposition model DEMENT (Allison, 2012) to evaluate four hypotheses: 1) allowing temperature response of CUE to differ (be heterogeneous) between taxa increases uncertainty in projected litter decomposition dynamics because more diverse phenotypic combinations exist for competitive selection (i.e. species sorting) to 65 act upon; 2) this heterogeneity favors a community with higher CUE, in turn leading to higher microbial biomass and greater litter C loss with warming; 3) forcing the temperature response of CUE to positively co-vary with the number of enzymes an organism produces causes greater litter C loss than when the two factors vary independently, because increasing CUE with temperature offsets the increased costs against CUE associated with copious enzyme production; and 4) the magnitude of litter C loss with warming is greater when the CUE of the C-rich fungal functional group increases with temperature than if only the 70 N-rich bacterial functional group does.

2 Methods

2.1 DEMENT Background and Model Design

DEMENT (Allison, 2012) is a litter decomposition model designed to simulate the loss of leaf C through time. The principal advancement of DEMENT over its predecessors is that it is both microbially- and spatially-explicit. The model is able to 75 simulate inter- and intraspecific microbial interactions, with a primary focus on the tradeoff between the ability to take up and digest substrates, and the metabolic costs of creating and maintaining the machinery required to do so. Because these tradeoffs are both explicit and variable across taxa, DEMENT is an ideal model for evaluating how the physiology and ecology of microbes affects C stocks in a changing world. Furthermore, DEMENT allows for consideration of how diversity in responses across taxa (rather than using some cross-taxon mean) can facilitate soil C responses to climate change. Full details about the 80 setup and execution of DEMENT are available elsewhere (Allison, 2012, 2014; Allison and Goulden, 2017); here we describe the model controls on CUE which are relevant to our study.

Intrinsic CUE—the maximum CUE an organism could attain under ideal temperature and stoichiometry—is calculated for each taxon as a function of the baseline CUE at 15 °C (C_t), and the number of enzymes and transporters the taxon can produce. In turn, how much CUE is decreased due to enzyme and transporter production depends on the cost per enzyme (C_e) and cost per transporter (C_u) (Table 1). The C used in enzyme synthesis is considered a loss from the cell, and is therefore not reported as microbial biomass C. The intrinsic CUE of each taxon is adjusted for temperature, decreasing by 0.016°C^{-1} by default (i.e. $C_t = -0.016^\circ\text{C}^{-1}$), consistent with a global meta-analysis (Qiao et al., 2019). Therefore, CUE is calculated as $CUE = C_i + (\text{temperature} - 288) * C_t + C_e * N_{enz} + C_u * N_{trans}$, where temperature is in kelvin and N_{enz} and N_{trans} are the number of enzyme genes and uptake transporter genes a taxon has, respectively.

90 2.2 Modifications to DEMENT

DEMENT v0.7.2 was downloaded from GitHub (<https://github.com/stevenallison/DEMENT>), and modified as follows. C_t was adjusted downwards from its original published value of 0.58 to 0.38 at 15°C; this not only improved model stability (Table S1), but is also consistent with a comparative modeling study completed by Li et al. (2014), several $^{18}\text{O}\text{-H}_2\text{O}$ based CUE measurements (Spohn et al., 2016a, b; Geyer et al., 2019), and for the structural components of litter modeled by MIMICS (Wieder et al., 2015a). By default, the temperature sensitivity of CUE (C_t) is fixed to take on the same value (ie is homogeneous) for all taxa; therefore we modified the model so C_t could vary around the mean in different ways (Figure 1). In the first set of scenarios, C_t varied independent of the taxonomic identity or number of enzymes a taxon produced. In the second scenario, C_t was limited to either increasing or decreasing as a function of the number of enzymes a given taxon had. In the third, bacteria and fungi were constrained to both have a positive C_t , both a negative C_t , or one a positive and the other a negative C_t . In all instances, C_t was selected at random from a uniform distribution bounded by $\pm 0.022^\circ\text{C}^{-1}$ at the upper and/or lower limits. These values are within the range of temperature sensitivities observed for both bacterial cultures in the lab and for field communities (Figure 2), as well as values inferred based on modeling CUE against mean annual temperature on a global basis (Sinsabaugh et al., 2017; Ye et al., 2019). It was necessary to force the temperature sensitivity of CUE to take on a zero-centered uniform distribution so that simulation outputs in which extracellular enzyme counts were linked to the C_t could be compared to those scenarios where they were not linked, without changing the distribution of extracellular enzyme counts present in the community.

2.3 Running DEMENT

DEMENT was run on the Massachusetts Green High Performance Computing Cluster for 6,000 model days using 59 different independent starting seeds and a 100x100 grid size. “Control” runs were completed at 15°C (equivalent to April to November mean soil temperature for a northern mid-latitude temperate deciduous forest (Boose, 2001)), while “heated” runs were completed at 20°C (Allison, 2014). The first 1000 days of each resultant output file was excluded from the analysis because of rapid shifts in the microbial community during this time. In addition, outputs were filtered to exclude any seeds where the substrate pool was two or more times greater at the end of the model run than the median during the preceding 5000 days, indicating unrealistic, unconstrained litter accumulation. R version 3.4.0 was used for all runs and analyses (R Core Team, 2016). A full

115 set of parameters and the model used to run all these simulations can be found in the supplementary files "params.txt" and
"DEMENTmodel.R", respectively.

2.4 Analysis of Outputs

The model outputs of interest were litter organic matter (LOM), microbial biomass carbon (MBC), respiration rate, richness
and diversity of the surviving community, median number of enzymes per taxon for taxa alive during the 5000-day simulation,
120 fungal:bacterial biomass ratio for surviving taxa, and biomass-weighted CUE at 15 and 20°C. Richness was calculated as the
number of taxa surviving to the end of the 5000 day run, while diversity was calculated as the median daily Shannon's H for the
duration of the simulation using the vegan package (Oksanen et al., 2017). Biomass-weighted CUE was reported for communi-
ties at 15°C ("reference temperature CUE") or at the simulation temperature ("simulation temperature CUE"), calculated using
the formula $\sum_{d=1000}^{6000} \sum_{i=1}^{100} C_r + C_e * N_{enz_i} + C_u * N_{trans_i} + C_{ti} * (temperature - 288) * biomass_i / \sum_{i=1}^{ntaxa} biomass_i$,
125 where 1000 and 6000 are the days of the simulation the outputs were examined over; 100 is the number of taxa the model was
initiated with; N_{enz_i} is the number of enzyme genes taxon i has; and N_{trans_i} is the number of uptake transporter genes the
taxon has; and temperature is the simulation temperature in °K.

In order to determine whether warming and model parameterization affected model outputs, we used mixed effect models
with starting seed as a random effect and warming or simulation scenario as fixed effects using lmer in lme4 v 1.1-17 (Bates
130 et al., 2015). Data were visually assessed for normality and homoskedasticity using qqplots and residual plots following log-
transformation. Significantly different pairwise differences were subsequently identified using emmeans v.1.3.0 (Lenth et al.,
2019), with a stringent Bonferoni-corrected p-value cutoff of $P < 0.0001$. Warming effect sizes are plotted as the natural log
ratio of model outputs in heated:control scenarios. Figures were generated using ggplot2 (Wickham, 2009), and asterisks and
letters denoting significant effects of experimental factors were added in Inkscape (Inkscape'sContributors, 2003–2019).

135 3 Results and discussion

LOM and MBC content were both generally higher than observed in environmental samples, leading to MBC:LOM ratios at
the high end of ranges observed in the field (2-11% vs. 1-5% (Santos et al., 2012; Xu et al., 2013)). LOM and MBC values
were within the range previously observed for simulations using DEMENT with daily litter inputs (Allison et al., 2014), but
greater than those with just a single litter pulse (Allison, 2012; Allison and Goulden, 2017; Evans et al., 2017), indicating that
140 these high biomass and litter C values can be attributed to these substrate inputs.

3.1 Intertaxon variability

To evaluate the effect of intertaxon CUE variability on LOM stocks, we ran the model at 20°C ("heated") under two scenarios,
and then compared the results to runs at 15°C ("control"). In the first "homogeneous" scenario all taxa had an identical C_t , equal
to 0°C^{-1} (Figure 1A). In the second "heterogeneous" scenario, C_t was assigned from a random uniform distribution bounded

145 by -0.022 and $0.022^{\circ}\text{C}^{-1}$ (Figure 1B). Therefore, the mean C_t of the heterogeneous scenario was 0°C^{-1} , matching that of the homogeneous scenario.

Introducing intertaxon differences in CUE temperature response caused the characteristics of the initial microbial community (starting seed) to have a greater impact on litter decomposition than when all taxa had an identical temperature response (Table 2). This contrasts with the dampening effect proposed to explain instability in small-scale microbially-explicit models compared to their macroscale counterparts (Wieder et al., 2015b), whereby the additive effect of increased physiological diversity was to increase, rather than decrease, uncertainty in the present simulations. The median-standardized interquartile ranges of both MBC (0.25 vs. 0.15) and LOM (0.28 vs. 0.14) increased with the introduction of a variable C_t . Through species sorting, this heterogeneously-responding microbial community became more uneven with warming, with similar richness but lower diversity than the control community (Table 2). The communities characterized by heterogeneous C_t maintained a higher median microbial biomass—driving two and a half times more LOM loss—than the communities characterized by a homogeneous C_t (Figure 3). Intriguingly, neither litter ($r=0.16$, $p=0.23$) nor microbial biomass pool sizes ($r=-0.44$, $P < 0.001$) positively correlated with extracellular enzyme investment; thus, a (non-significant) 28% increase in the median enzyme count is unlikely to have driven the increased decomposition under the heterogeneous scenario. Instead, increased decomposition and increased biomass are likely the consequence of elevated CUE under warming conditions.

160 The homogeneous community scenario tested here is akin to the fixed “no adaptation of CUE” scenario reported in a number of other studies (Allison et al., 2010; Li et al., 2014; Sistla et al., 2014), because the cross-taxon mean used is zero temperature response. However, the effect of inter-taxon differences in C_t (“heterogeneous”) is less studied. Our results of reduced LOM loss in the absence of acclimation are consistent with two previous studies, but contrast with others. In an ecosystem-level model parameterized for an arctic tundra system, Sistla *et al.* (2014) found that greater soil organic matter (SOM) loss occurred with warming when the microbial community was able to dynamically acclimate its CN ratio (and in turn efficiency), than when the CUE was effectively fixed. Likewise, Allison (2014) found greater potential for increased LOM accumulation under warming when there was greater absolute variation in CUE across taxa (i.e. C_t from 0.18 to 0.58 rather than 0.38 to 0.58) (Allison, 2014), although C_t was always consistent across taxa in these simulations. On the other hand, a comparison of models where taxon-level differences in CUE or C_t are not possible (i.e. C_t is intrinsically homogeneous at the community level) showed that soil organic matter loss increases when organisms do not adapt (Li et al., 2014). Similarly, Wieder *et al.* found that greater SOM loss occurred if the CUE was directly insensitive to temperature than when C_t was negative (Wieder et al., 2014). These microbially-explicit decomposition models vary in if and how they link CUE to microbial traits, and so our findings support the concept that nuances in how different components of CUE respond to warming is an important control on the fate of litter C (Hagerty et al., 2018).

175 3.2 Confirming the role of C_t as an additional niche dimension

We allowed for CUE to increase with temperature for a subset of taxa in a way that most previous modeling efforts have not, and so it is possible that our results deviate from those of prior studies not because of variation in C_t , but rather because our simulations explore novel (positive C_t) parameter space. To facilitate comparison with previous decomposition modeling

studies, we ran DEMENT simulations to test the effect of C_t being homogeneous vs. heterogeneous when CUE was either
180 always positive (homogeneous $C_t = 0.011 \text{ } ^\circ\text{C}^{-1}$ (Figure S1A_i), heterogeneous $C_t = 0$ to $0.022 \text{ } ^\circ\text{C}^{-1}$ (Figure S1G)) or always
negative (homogeneous $C_t = -0.011 \text{ } ^\circ\text{C}^{-1}$ (Figure S1A_{ii}), heterogeneous $C_t = -0.022$ to $0 \text{ } ^\circ\text{C}^{-1}$ (Figure S1H). In contrast to
when C_t was allowed to vary over the whole spectrum of values, introducing heterogeneity in CUE did not increase inter-
run uncertainty in LOM or MBC pools (Table S2). We also found that less LOM accumulated when CUE showed a variable
185 decrease with warming than a fixed one (Figure S2), which could be attributed to a reduction in MBC. By contrast, the
homogeneous zero-centered and homogeneous positive C_t scenarios, and the heterogeneous zero-centered and heterogeneous
positive C_t , behaved more similarly to one-another in that warming decreased LOM while increasing MBC and CUE to a
greater degree in the heterogeneous than homogeneous scenarios (Table S2). This finding reinforces the idea that if warming
favors decomposer taxa capable of maintaining efficient growth, then soil C loss will be accelerated. Nonetheless, the strongly
190 selected-for positive CUE response is rarely observed in complex soil communities. This indicates that additional tradeoffs
with CUE temperature response are likely at play when CUE is either unaffected or decreases with temperature, but that these
tradeoffs are missing in the formulation of DEMENT used in this scenario. One such tradeoff possible to explore within the
framework of DEMENT is the allocation of resources to extracellular enzyme activity.

3.3 Linkages between CUE temperature response and extracellular enzyme allocation

Microbes depend upon extracellular enzymes to break down substrates in the environment into digestible pieces, and enzyme
195 activities are, like CUE, responsive to temperature (Wallenstein et al., 2010; German et al., 2012; Allison et al., 2018). Soil
extracellular enzymes often are active *in-situ* at temperatures much below their activity optima (German et al., 2012; Pold
et al., 2017; Alster et al., 2018). Therefore, warming enables them to process substrates at a higher rate, increasing the supply
of growth substrates to microbes. However, the affinity of enzymes for their substrates also decreases as temperature increases
(German et al., 2012; Allison et al., 2018); unless enzyme V_{\max} increases faster with temperature than K_m , additional resources
200 must be diverted from growth to enzyme production to maintain microbial growth substrate supply rate. Therefore, taxa may
differentially-allocate resources to enzymes and so demonstrate a relationship between the temperature sensitivity of CUE and
the number of enzymes they produce.

We evaluated whether litter decomposition changed its trajectory when the organisms with the greatest genomic potential to
break the litter down (i.e. enzyme counts) also showed the most- or least-positive growth efficiency response to warming. In the
205 “increase” scenario, we simulated a positive relationship between temperature sensitivity of CUE and extracellular enzymes,
where C_t increased linearly from $-0.022^\circ\text{C}^{-1}$ for organisms with no extracellular enzyme production potential to 0.022°C^{-1} for
those organisms capable of producing the model maximum of 40 enzymes (Figure 1C). In the “decrease” scenario, we simulated
a negative relationship between temperature sensitivity of CUE and extracellular enzymes, where the opposite relationship was
imposed with C_t decreasing with enzyme counts (Figure 1D). These scenarios were then compared to the “heterogeneous”
210 scenario (aka “no relation”, as described above), where C_t varied across the same range, but independently of the number of
enzymes an organism could produce. Therefore, the starting distribution of C_t and enzymes per taxon was identical across
scenarios, and only their relationship with one-another changed.

More taxa survived to the end of the simulation when warming was applied under the “increase” scenario than either the “decrease” or “no relation” scenario (median of 13 versus 7 and 8, respectively, median absolute deviation = 2.97 in all cases).
215 Under the “increase” scenario, taxa had 70% more enzymes each than the “no relation” scenario, and more than three times as much as the “decrease” scenario (Table 2). This relationship caused the CUE of surviving taxa to be 20-37% lower at 15°C for the “increase” scenario compared to the others, but this deficit was diminished at 20°C. As a result, the “increase” scenario led to higher respiration and a greater LOM loss under warming than under the “decrease” scenario, despite an overall smaller microbial biomass pool (Figure 4).

220 How the relationship between C_i and extracellular enzymes drives favorable trait combinations can also be observed in Figure 5. Surviving taxa retained a median enzyme count of at least 30 and a realized CUE temperature response of no less than $0.0158^{\circ}\text{C}^{-1}$ under the “increasing” scenario ($\rho=0.64$, $P < 0.001$), but there was no relationship between realized CUE temperature response and enzyme production under either the “decrease” or “no relation” scenarios. The selection for community capable of maintaining high CUE at high temperatures was much weaker when it was associated with reduced enzyme produc-
225 tion. When there was no relationship between C_i and extracellular enzyme production costs, however, communities were able to attain a high realized CUE temperature response over a much wider range of median enzyme costs.

These findings indicate that response traits—which determine how an organism reacts to changes in temperature (e.g. CUE temperature response)—and effect traits—which determine how an organism alters its environment (e.g. litter decomposition potential)—interact to determine the fate of organic C. However, contrary to our hypothesis, adjusting DEMENT to allow
230 for this tradeoff did not substantially alter how community-level CUE responds to temperature. The observation that LOM is reduced further when enzyme production is effectively cheaper contrasts with earlier work with DEMENT (Allison, 2014) showing smaller litter C pools under both ambient and elevated temperature when enzymes and transporters were cheaper to produce. However, our results are consistent in that microbial biomass was lower when enzyme costs are high, and that the microbial community was able to maintain a higher CUE under warming no matter the enzyme costs. The mechanisms
235 underlying these phenomenological similarities differ, however, due to differences in how C_i was parameterized in the two sets of model simulations. Specifically, although microbes were able to attain high CUE at elevated temperatures in our simulations by balancing the benefits of elevated CUE at higher temperatures with the costs of enzyme production against CUE, CUE always decreased with temperature in earlier work with DEMENT (Allison, 2014). Furthermore, enzyme production costs varied both with and independently of enzyme counts in previous DEMENT simulations (Allison, 2014).

240 Increased CUE is likely needed to offset the costs of extracellular enzyme production if taxa are to remain competitive at elevated temperatures within the framework of DEMENT, but empirical support for correlations between temperature sensitivity of CUE and enzyme investment are needed to best put these inferences to use. By examining correlations between the number of enzymes an organism can produce and its CUE temperature response at the end of the DEMENT model run, we see that there is likely to be either no or a positive correlation between the two variables, rather than a negative one (Figure 6).
245 Limited data from bacterial isolates grown in the lab also support this, whereby CUE temperature response is either positively correlated or uncorrelated with the number of enzymes produced (Pold et al., in prep). Furthermore, although the mechanisms underlying the isolate response remain unclear, it is consistent with the scenario of C_i and enzyme counts being positively cor-

related. Specifically, we found isolates with lower CUE at 15°C (more extracellular enzymes in DEMENT) were more likely to have a positive CUE temperature response than those with a higher CUE (fewer enzymes in DEMENT). Together, these insights support a synergism between CUE temperature response and enzyme production, rather than a tradeoff. Because our DEMENT simulations indicate that selection for organisms characterized by high, positive CUE temperature responses with warming can alter both the directionality and extent of projected C loss, we propose it is important for other models to explore how possible increases—rather than just decreases—in C_t affect terrestrial C projections.

3.4 Ecological relevance of microbial metabolic diversity—bacteria vs. fungi

Across scenarios, fungi generally dominated the microbial biomass C pool (Table 2), as is typical for litter decomposition (Chapman et al. (2013), and references therein). This pattern occurred despite generally lower biomass-weighted CUE for surviving fungal taxa, and preferential loss of fungal taxa across most scenarios (Table 2). The lower CUE for surviving fungi was not driven by higher enzyme costs than for bacteria, as median biomass-weighted enzyme costs were not statistically different (pairwise t-test $P > 0.4$) and differed by less than one enzyme for the two groups. To test whether modeled differences in fungal vs. bacterial cell sizes and stoichiometry were driving this pattern, we tested how forcing differences in C_t in the two groups would impact the decomposition rate.

DEMENT was run with CUE simulated to respond to temperature: 1) negatively for all bacteria and positively for all fungi (F+B-; Figure 1E); 2) negatively for all fungi and positively for all bacteria (F-B+; Figure 1F); 3) positively for all bacterial and fungal taxa (F+B+; Figure 1G) or 4) negatively for all bacterial and fungal taxa (F-B-; Figure 1F). Minimum and maximum C_t were set to -0.022 °C and 0.022 °C, respectively.

Litter C accumulated at higher rates with warming when fungal C_t was negative, regardless of the bacterial C_t (Figure 7, blue, orange). This is consistent with the observation that the CUE of surviving fungi was lower at the simulation temperature in seven out of the eleven scenarios. Because fungi have higher CNP ratios than bacteria (and thus higher C demands per unit biomass), we predicted that if fungi have a negative CUE temperature response, they would be weaker competitors at higher temperature than bacterial taxa, reducing their C demand and mitigating the warming effect on SOC stocks. While this contrasts with the premise that fungi should have a higher CUE (Six et al., 2006) due to their higher CN ratio (Zak et al., 1996), it is consistent with a growing body of literature indicating that substrate quality—rather than the F:B ratio correlated with it—is the underlying driver of differences in CUE between soils (Frey et al., 2013; Thiet et al., 2006; Malik et al., 2018; Soares and Rousk, 2019). Despite the low nutrient content of the daily inputs to the model (92:0.26:0.02 C:N:P), microbes did not show evidence for nutrient limitation as biomass CN and CP ratios were lower than are typical for soil communities (Xu et al., 2013) (4.1 and 36.7 vs. 7.6 and 42.4, respectively). Thus C was limiting, which could have further disfavored the highly C-demanding fungi when their C_t was negative. The lower CUE (and increased sensitivity to warming) for fungi compared to bacteria under a given scenario was also not driven by increased metabolic costs for enzyme production in fungi, as median biomass-weighted enzyme counts were statistically indistinguishable from those in bacteria.

The litter C pool decreased when fungal CUE increased with temperature (Figure 7), correlating with a smaller microbial biomass pool when bacterial CUE also decreased with temperature. By contrast, the LOM and MBC responses to warming were

similar when only the C_t of bacteria was changed, indicating that it is the fungal warming response which really drives changes in litter decomposition in DEMENT. This result is interesting because no *a priori* differences in decomposition or uptake potential were imposed on the two groups, and fungal and bacterial richness was initially equivalent. Warming decreased the enzyme costs when fungal C_t was positive but bacterial C_t was negative, and decreased them under the opposing scenario, as evidenced by an increase in C_r in the former and decrease in the latter. Nonetheless, as long as both bacterial and fungal CUE did not both decrease their CUE with temperature, community level CUE remained higher at 20°C than it was at 15°C.

Empirical evidence for high-level differences in the temperature sensitivity of CUE in bacteria and fungi is currently mixed, but indicate CUE temperature response for fungi is unlikely to be more positive than that for bacteria. Zheng et al. (2019) did not find a correlation between the lipid-based fungal:bacterial ratio and Q_{10} of CUE over a range of soils. However, we (Pold et al., in prep) and our colleagues (Eric Morrison, personal comment) have found that fungi tend to show a stronger negative CUE response with warming than do bacteria when examining them in isolation in the lab. This is consistent with the observation that fungal CUE decreases more strongly with warming than bacterial CUE does when C_t is restricted to negative values (Figure 7). It is also consistent with the premise that bacterial growth benefits more from elevated temperature than does fungal growth in some soils (Pietikäinen et al., 2005). Greater empirical insight into the taxonomic drivers of the temperature sensitivity of CUE will assist with constraining the parameterization and projections of microbially-explicit decomposition models such as DEMENT.

3.5 Comparison to empirical warming studies

Litter decomposition is typically observed to accelerate under warming (Lu et al., 2013). However, both the chemical composition of the litter and the identity of the living plant community at the site of decomposition are important for the magnitude of this response (Cornelissen et al., 2007; Ward et al., 2015). Consistent with these empirical studies—but inconsistent with a previous publication using DEMENT (Allison, 2014)—we found that litter decomposition was accelerated by warming in seven of ten scenarios. The range of losses and gains of litter C we observed with warming (-62% (scenario C) to +42% (scenario A_i)) approximates the -65% to +36% change observed in field experiments (Lu et al., 2013), with the upper limit only being exceeded when C_t is constrained to negative values. Likewise, values for simulated litter respiration response (-5 to +6%) fell within those observed for soil respiration in the field (-48 to +178%), and responses for microbial biomass C were also within the observed range (-25 to +58% vs -47 to +86%) (Lu et al., 2013). Our modeled responses to warming thus suggest that one possible explanation for differences in terrestrial C pool responses to warming may be diverse temperature sensitivities of underlying decomposer communities. Nonetheless, a number of additional factors must be taken into consideration when interpreting our results in the context of global climate change, including soil mineral-mediated modulation of substrate supply (Schimel and Schaeffer, 2012; Coward et al., 2018), plant-microbe feedbacks (Melillo et al., 2011; Sistla et al., 2014; Suseela and Tharayil, 2018), and temporal variation in temperature.

While we explored how different relationships between temperature sensitivity of CUE and enzyme count or taxonomic affiliation affects the C cycle, empirical support for these scenarios is lacking. By including additional dimensions to the warming response we showed that species effectively sort to make communities with differing impacts on litter decomposition.

Following up on the direction and magnitude of these underlying metabolic scenarios in the soil and litter can help us better constrain our model results on reality. New and emerging work indicates that the temperature sensitivity of CUE is weakly negatively correlated with the temperature sensitivity of extracellular enzyme activity, and with bacterial—but not fungal—biomass on the community level (Zheng et al., 2019). However, different kinds of bacteria and fungi are able to show positive, 320 negative, or no temperature sensitivity of CUE, and so high-level assumptions about how these groups respond are unlikely to resolve uncertainties about the magnitude of soil C loss under warming. Finally, CUE and its temperature response are not fixed parameters, but rather a mathematical outcome of a complex balance between the diverse maintenance and growth costs that microbes experience. For instance, similar CUE and C_t can appear for two taxa when they allocate carbon to processes such as exopolysaccharide or extracellular enzyme production with very different impacts on decomposition (Hagerty et al., 325 2018). Allowing CUE and C_t to vary as the outcome of explicit biomass allocation functions is a necessary next step to validate our results.

4 Conclusions

Our results indicate that accounting for heterogeneous temperature response increases uncertainty regarding future litter C stocks, but only when C_t does not differ from zero on average. However, by combining simulations, empirical studies, and 330 literature searches, we can conclude that microbes with high enzyme costs are likely to have larger increases in intrinsic CUE with temperature; that taxa can sort on a CUE temperature response axis; and that fungi are more likely to increase CUE with warming than bacteria. The simulations meeting each or all of these criteria lead to loss of litter C under warming, indicating that litter is likely to become a net atmospheric C source in a warmer world. We encourage models functioning on larger scales to explore the effect of including heterogeneity in the temperature response of CUE in order to determine the robustness of 335 our conclusions to other model structures. However, ultimately increased integration of the growing body of literature on the temperature sensitivity of CUE must be explored for root causes of heterogeneity in temperature sensitivity of CUE in taxa under *in situ* conditions.

Code and data availability. The data and modified version of DEMENT used to generate it are available at OSF (https://osf.io/cwep9/?view_only=de0e6ba7b7da493f96531f398ca62c2c) under DOI 10.17605/OSF.IO/CWEP9

340 *Author contributions.* GP SAS and KMD conceived the experiment; GP completed the experiment and analyzed data; GP wrote the first draft of the manuscript; all authors contributed to revising the manuscript and interpreting the results

Competing interests. The authors declare no competing interests

Acknowledgements. The authors wish to thank Steve Allison for sharing DEMENT and providing insight into troubleshooting the model. Funding for this project came from the US Department of Energy grant DE-SC0016590 to KMD and SAS, and from an American Association
345 of University Women American Fellowship to GP.

References

- Allison, S.: A trait-based approach for modelling microbial litter decomposition, *Ecol. Lett.*, 15, 1058–1070, 2012.
- Allison, S. D.: Modeling adaptation of carbon use efficiency in microbial communities, *Front Microbiol*, 5, 571, 2014.
- Allison, S. D. and Goulden, M. L.: Consequences of drought tolerance traits for microbial decomposition in the DEMENT model, *Soil Biol Biochem*, 107, 104–113, 2017.
- Allison, S. D., Wallenstein, M. D., and Bradford, M. A.: Soil-carbon response to warming dependent on microbial physiology, *Nat Geosci*, 3, 336, 2010.
- Allison, S. D., Lu, L., Kent, A. G., and Martiny, A. C.: Extracellular enzyme production and cheating in *Pseudomonas fluorescens* depend on diffusion rates, *Front Microbiol*, 5, 2014.
- Allison, S. D., Romero-Olivares, A. L., Lu, Y., Taylor, J. W., and Treseder, K. K.: Temperature sensitivities of extracellular enzyme V_{max} and K_m across thermal environments, *Global Change Biol*, 24, 2884–2897, 2018.
- Alster, C. J., Weller, Z. D., and von Fischer, J. C.: A Meta-Analysis of Temperature Sensitivity as a Microbial Trait, *Global Change Biol*, 2018.
- Amend, A. S., Martiny, A. C., Allison, S. D., Berlemont, R., Goulden, M. L., Lu, Y., Treseder, K. K., Weihe, C., and Martiny, J. B.: Microbial response to simulated global change is phylogenetically conserved and linked with functional potential, *ISME J*, 10, 109–118, 2016.
- Bates, D., Mächler, M., Bolker, B., and Walker, S.: Fitting Linear Mixed-Effects Models Using lme4, *J Stat Softw*, 67, 1–48, <https://doi.org/10.18637/jss.v067.i01>, 2015.
- Blagodatskaya, E., Blagodatsky, S., Anderson, T.-H., and Kuzyakov, Y.: Microbial growth and carbon use efficiency in the rhizosphere and root-free soil, *PLoS one*, 9, e93282, 2014.
- Boose, E.: Fisher Meteorological Station at Harvard Forest since 2001. Harvard Forest Data Archive: HF001, doi = doi:10.6073/pasta/04076dfd30b286c6c29301b6345a63f5, howpublished= <http://harvardforest.fas.harvard.edu:8080/exist/apps/datasets/showData.html?id=hf001>, 2001.
- Cavicchioli, R.: On the concept of a psychrophile, *ISME J*, 10, 793, 2016.
- Chapman, S. K., Newman, G. S., Hart, S. C., Schweitzer, J. A., and Koch, G. W.: Leaf litter mixtures alter microbial community development: mechanisms for non-additive effects in litter decomposition, *PLoS One*, 8, e62671, 2013.
- Cornelissen, J. H., Van Bodegom, P. M., Aerts, R., Callaghan, T. V., Van Logtestijn, R. S., Alatalo, J., Stuart Chapin, F., Gerdol, R., Gudmundsson, J., Gwynn-Jones, D., et al.: Global negative vegetation feedback to climate warming responses of leaf litter decomposition rates in cold biomes, *Ecol. Lett.*, 10, 619–627, 2007.
- Coward, E. K., Ohno, T., and Sparks, D. L.: Direct Evidence for Temporal Molecular Fractionation of Dissolved Organic Matter at the Iron Oxyhydroxide Interface, *Environmental Sci Technol*, 53, 642–650, 2018.
- DeAngelis, K. M., Pold, G., Topçuoğlu, B. D., van Diepen, L. T., Varney, R. M., Blanchard, J. L., Melillo, J., and Frey, S. D.: Long-term forest soil warming alters microbial communities in temperate forest soils, *Front Microbiol*, 6, 104, 2015.
- Devêvre, O. C. and Horwáth, W. R.: Decomposition of rice straw and microbial carbon use efficiency under different soil temperatures and moistures, *Soil Biol Biochem*, 32, 1773–1785, 2000.
- Dijkstra, P., Dalder, J. J., Selmants, P. C., Hart, S. C., Koch, G. W., Schwartz, E., and Hungate, B. A.: Modeling soil metabolic processes using isotopologue pairs of position-specific ^{13}C -labeled glucose and pyruvate, *Soil Biol Biochem*, 43, 1848–1857, 2011a.

- Dijkstra, P., Thomas, S. C., Heinrich, P. L., Koch, G. W., Schwartz, E., and Hungate, B. A.: Effect of temperature on metabolic activity of intact microbial communities: evidence for altered metabolic pathway activity but not for increased maintenance respiration and reduced carbon use efficiency, *Soil Biol Biochem*, 43, 2023–2031, 2011b.
- 385 Evans, S., Martiny, J. B., and Allison, S. D.: Effects of dispersal and selection on stochastic assembly in microbial communities, *ISME J*, 11, 176, 2017.
- Frey, S. D., Lee, J., Melillo, J. M., and Six, J.: The temperature response of soil microbial efficiency and its feedback to climate, *Nat Clim Change*, 3, 395, 2013.
- German, D. P., Marcelo, K. R., Stone, M. M., and Allison, S. D.: The Michaelis–Menten kinetics of soil extracellular enzymes in response
390 to temperature: a cross-latitudinal study, *Global Change Biol*, 18, 1468–1479, 2012.
- Geyer, K. M., Dijkstra, P., Sinsabaugh, R., and Frey, S. D.: Clarifying the interpretation of carbon use efficiency in soil through methods comparison, *Soil Biol Biochem*, 128, 79–88, 2019.
- Hagerty, S. B., Allison, S. D., and Schimel, J. P.: Evaluating soil microbial carbon use efficiency explicitly as a function of cellular processes: implications for measurements and models, *Biogeochemistry*, 140, 269–283, 2018.
- 395 Inkscape’sContributors: Inkscape, <https://inkscape.org>, 2003–2019.
- Kafri, M., Metzl-Raz, E., Jona, G., and Barkai, N.: The cost of protein production, *Cell reports*, 14, 22–31, 2016.
- Kaleta, C., Schäuble, S., Rinas, U., and Schuster, S.: Metabolic costs of amino acid and protein production in *Escherichia coli*, *Biotechnol J*, 8, 1105–1114, 2013.
- Kallenbach, C. M., Frey, S. D., and Grandy, A. S.: Direct evidence for microbial-derived soil organic matter formation and its ecophysiological controls, *Nat Commun*, 7, 13 630, 2016.
- 400 Keiblinger, K. M., Hall, E. K., Wanek, W., Szukics, U., Hämmerle, I., Ellersdorfer, G., Böck, S., Strauss, J., Sterflinger, K., Richter, A., et al.: The effect of resource quantity and resource stoichiometry on microbial carbon-use-efficiency, *FEMS Microbiol Ecol*, 73, 430–440, 2010.
- Lavorel, S. and Garnier, É.: Predicting changes in community composition and ecosystem functioning from plant traits: revisiting the Holy Grail, *Funct Ecol*, 16, 545–556, 2002.
- 405 Lee, Z. M. and Schmidt, T. M.: Bacterial growth efficiency varies in soils under different land management practices, *Soil Biol Biochem*, 69, 282–290, 2014.
- Lenth, R., Henrik, S., Jonathon, L., Paul, B., and Herve, M.: emmeans, Iowa State, <https://cran.r-project.org/web/packages/emmeans/index.html>, 2019.
- Li, J., Wang, G., Allison, S. D., Mayes, M. A., and Luo, Y.: Soil carbon sensitivity to temperature and carbon use efficiency compared across
410 microbial-ecosystem models of varying complexity, *Biogeochemistry*, 119, 67–84, 2014.
- Li, J., Wang, G., Mayes, M. A., Allison, S. D., Frey, S. D., Shi, Z., Hu, X.-M., Luo, Y., and Melillo, J. M.: Reduced carbon use efficiency and increased microbial turnover with soil warming, *Global Change Biol*, 2018.
- Lu, M., Zhou, X., Yang, Q., Li, H., Luo, Y., Fang, C., Chen, J., Yang, X., and Li, B.: Responses of ecosystem carbon cycle to experimental warming: a meta-analysis, *Ecology*, 94, 726–738, 2013.
- 415 Malik, A. A., Puissant, J., Buckeridge, K. M., Goodall, T., Jehmlich, N., Chowdhury, S., Gweon, H. S., Peyton, J. M., Mason, K. E., van Agtmaal, M., et al.: Land use driven change in soil pH affects microbial carbon cycling processes, *Nat Commun*, 9, 3591, 2018.
- Malik, A. A., Puissant, J., Goodall, T., Allison, S. D., and Griffiths, R. I.: Soil microbial communities with greater investment in resource acquisition have lower growth yield, *Soil Biol Biochem*, 2019.

- Manzoni, S., Taylor, P., Richter, A., Porporato, A., and Ågren, G. I.: Environmental and stoichiometric controls on microbial carbon-use efficiency in soils, *New Phytol*, 196, 79–91, 2012.
- Martiny, A. C., Treseder, K., and Pusch, G.: Phylogenetic conservatism of functional traits in microorganisms, *ISME J*, 7, 830, 2013.
- Melillo, J. M., Butler, S., Johnson, J., Mohan, J., Steudler, P., Lux, H., Burrows, E., Bowles, F., Smith, R., Scott, L., Vario, C., Hill, T., Burton, A., Zhou, Y.-M., and Tang, J.: Soil warming, carbon–nitrogen interactions, and forest carbon budgets, *P Natl Acad Sci USA*, 108, 9508–9512, 2011.
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P. R., O’Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., and Wagner, H.: *vegan: Community Ecology Package*, <https://CRAN.R-project.org/package=vegan>, r package version 2.4-2, 2017.
- Oldfield, E. E., Crowther, T. W., and Bradford, M. A.: Substrate identity and amount overwhelm temperature effects on soil carbon formation, *Soil Biol Biochem*, 124, 218–226, 2018.
- Öquist, M. G., Erhagen, B., Haei, M., Sparrman, T., Ilstedt, U., Schleucher, J., and Nilsson, M. B.: The effect of temperature and substrate quality on the carbon use efficiency of saprotrophic decomposition, *Plant Soil*, 414, 113–125, 2017.
- Pietikäinen, J., Pettersson, M., and Bååth, E.: Comparison of temperature effects on soil respiration and bacterial and fungal growth rates, *FEMS Microbiol Ecol*, 52, 49–58, 2005.
- Pold, G., Grandy, A. S., Melillo, J. M., and DeAngelis, K. M.: Changes in substrate availability drive carbon cycle response to chronic warming, *Soil Biol Biochem*, 110, 68–78, 2017.
- Pold, G., Bales, A., Domeignoz-Horta, L. A., Sistla, S. A., Morisson, E., Frey, S. D., and DeAngelis, K.: Carbon use efficiency and its sensitivity to temperature co-vary in soil bacteria, in prep.
- Qiao, Y., Wang, J., Liang, G., Du, Z., Zhou, J., Zhu, C., Huang, K., Zhou, X., Luo, Y., Yan, L., et al.: Global variation of soil microbial carbon-use efficiency in relation to growth temperature and substrate supply, *Sci Rep*, 9, 5621, 2019.
- R Core Team: *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, 2016.
- Santos, V. B., Araújo, A. S., Leite, L. F., Nunes, L. A., and Melo, W. J.: Soil microbial biomass and organic matter fractions during transition from conventional to organic farming systems, *Geoderma*, 170, 227–231, 2012.
- Schimel, J. and Schaeffer, S. M.: Microbial control over carbon cycling in soil, *Front Microbiol*, 3, 348, 2012.
- Sinsabaugh, R. L., Turner, B. L., Talbot, J. M., Waring, B. G., Powers, J. S., Kuske, C. R., Moorhead, D. L., and Follstad Shah, J. J.: Stoichiometry of microbial carbon use efficiency in soils, *Ecol Monogr*, 86, 172–189, 2016.
- Sinsabaugh, R. L., Moorhead, D. L., Xu, X., and Litvak, M. E.: Plant, microbial and ecosystem carbon use efficiencies interact to stabilize microbial growth as a fraction of gross primary production, *New Phytol*, 214, 1518–1526, 2017.
- Sistla, S. A., Rastetter, E. B., and Schimel, J. P.: Responses of a tundra system to warming using SCAMPS: a stoichiometrically coupled, acclimating microbe–plant–soil model, *Ecol Monogr*, 84, 151–170, 2014.
- Six, J., Frey, S., Thiet, R., and Batten, K.: Bacterial and fungal contributions to carbon sequestration in agroecosystems, *Soil Sci Soc Am J*, 70, 555–569, 2006.
- Soares, M. and Rousk, J.: Microbial growth and carbon use efficiency in soil: Links to fungal-bacterial dominance, SOC-quality and stoichiometry, *Soil Biol Biochem*, 2019.
- Spohn, M., Klaus, K., Wanek, W., and Richter, A.: Microbial carbon use efficiency and biomass turnover times depending on soil depth—Implications for carbon cycling, *Soil Biol Biochem*, 96, 74–81, 2016a.

- Spohn, M., Pötsch, E. M., Eichorst, S. A., Wobken, D., Wanek, W., and Richter, A.: Soil microbial carbon use efficiency and biomass turnover in a long-term fertilization experiment in a temperate grassland, *Soil Biol Biochem*, 97, 168–175, 2016b.
- 460 Suseela, V. and Tharayil, N.: Decoupling the direct and indirect effects of climate on plant litter decomposition: Accounting for stress-induced modifications in plant chemistry, *Global Change Biol*, 24, 1428–1451, 2018.
- Tang, J. and Riley, W. J.: Weaker soil carbon-climate feedbacks resulting from microbial and abiotic interactions, *Nat Clim Change*, 5, 56–60, 2015.
- Thiet, R. K., Frey, S. D., and Six, J.: Do growth yield efficiencies differ between soil microbial communities differing in fungal: bacterial ratios? Reality check and methodological issues, *Soil Biol Biochem*, 38, 837–844, 2006.
- 465 Treseder, K. K. and Lennon, J. T.: Fungal traits that drive ecosystem dynamics on land, *Microbiology and Molecular Biology Reviews*, 79, 243–262, 2015.
- Walker, T. W., Kaiser, C., Strasser, F., Herbold, C. W., Leblans, N. I., Wobken, D., Janssens, I. A., Sigurdsson, B. D., and Richter, A.: Microbial temperature sensitivity and biomass change explain soil carbon loss with warming, *Nat Clim Change*, 8, 885, 2018.
- Wallenstein, M., Allison, S. D., Ernakovich, J., Steinweg, J. M., and Sinsabaugh, R.: Controls on the temperature sensitivity of soil enzymes: a key driver of in situ enzyme activity rates, in: *Soil enzymology*, pp. 245–258, Springer, 2010.
- 470 Ward, S. E., Orwin, K. H., Ostle, N. J., Briones, M. J., Thomson, B. C., Griffiths, R. I., Oakley, S., Quirk, H., and Bardgett, R. D.: Vegetation exerts a greater control on litter decomposition than climate warming in peatlands, *Ecology*, 96, 113–123, 2015.
- Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag New York, <http://ggplot2.org>, 2009.
- Wieder, W., Grandy, A., Kallenbach, C., and Bonan, G.: Integrating microbial physiology and physio-chemical principles in soils with the 475 Microbial-MIneral Carbon Stabilization (MIMICS) model, *Biogeosciences*, 11, 3899–3917, 2014.
- Wieder, W., Grandy, A., Kallenbach, C., Taylor, P., and Bonan, G.: Representing life in the Earth system with soil microbial functional traits in the MIMICS model, *Geoscientific Model Development*, 8, 1789–1808, 2015a.
- Wieder, W. R., Bonan, G. B., and Allison, S. D.: Global soil carbon projections are improved by modelling microbial processes, *Nat Clim Change*, 3, 909, 2013.
- 480 Wieder, W. R., Allison, S. D., Davidson, E. A., Georgiou, K., Hararuk, O., He, Y., Hopkins, F., Luo, Y., Smith, M. J., Sulman, B., et al.: Explicitly representing soil microbial processes in Earth system models, *Global Biogeochemical Cycles*, 29, 1782–1800, 2015b.
- Xu, X., Thornton, P. E., and Post, W. M.: A global analysis of soil microbial biomass carbon, nitrogen and phosphorus in terrestrial ecosystems, *Global Ecol Biogeogr*, 22, 737–749, 2013.
- Ye, J., Bradford, M. A., Dacal, M., Maestre, F. T., and García-Palacios, P.: Increasing microbial carbon use efficiency with warming predicts 485 soil heterotrophic respiration globally, *Global Change Biol*, <https://doi.org/doi:10.1111/gcb.14738>, 2019.
- Zak, D. R., Ringelberg, D. B., Pregitzer, K. S., Randlett, D. L., White, D. C., and Curtis, P. S.: Soil microbial communities beneath *Populus grandidentata* grown under elevated atmospheric CO₂, *Ecol Appl*, 6, 257–262, 1996.
- Zheng, Q., Hu, Y., Zhang, S., Noll, L., Böckle, T., Richter, A., and Wanek, W.: Growth explains microbial carbon use efficiency across soils differing in land use and geology, *Soil Biol Biochem*, 128, 45–55, 2019.




Figure 1. Schematic of experimental design used in this study, where CUE temperature response (C_t) varies as a function of the number of enzymes and/or taxonomic affiliation of organisms. Graphs show the effects of having homogeneous (A) or heterogeneous (B) C_t across taxa; the effect of forcing a positive (C) or negative (D) correlation between the number of enzymes and C_t ; bacteria showing a positive C_t and fungi a negative C_t (E); fungi showing an positive C_t and bacteria a negative C_t (F); or fungi and bacteria both having positive (G) or negative (H) C_t . Horizontal dashed lines indicate a C_t of zero, and clusters of points above and below this line denote when CUE tends to increase or decrease with increasing temperature. The letters F and B in the x-axis of individual graphs denote sensitivities for fungi and bacteria, respectively. Figure prepared in BioRender




Figure 2. Density plot of observed CUE temperature response for 23 soil bacterial isolates grown between 15 and 25°C on four different liquid media types in the lab (n=160 datapoints, grey), and for soil microbial communities grown with various different substrates and temperatures based on a literature search (n=141 datapoints, brown). Vertical lines are placed at 0 (no change in CUE with temperature) as well as the ± 0.022 °C⁻¹ upper and lower limits used in the present study. Contributing datapoints are primarily derived from Qiao et al. (2019) and can be found in the "Ct_literatureValues.txt" file in OSF.




Figure 3. Effect of warming 5°C on C stocks and flows in simulations, reported as the natural log of the ratio of values in heated compared to control conditions. CUE temperature response either varied between taxa (“heterogeneous”) or took on the fixed cross-sample mean temperature response of zero. Values above the zero line indicate warming increased the value, and values below indicate a decrease with warming. Boxplots denote 1st to 3rd quartiles with the median. Asterisks denote significant warming effect based on a paired Wilcoxon test at Bonferoni-corrected $P < 0.0001$. Letters denote warmed scenarios which are significantly different from one-another by the same criteria. “Reference temperature CUE” denotes the ratio of CUE for the surviving community at 15°C in warmed to the ratio of CUE for the surviving community at 15°C control scenarios, while “Simulation temperature CUE” denotes the ratio of CUE in heated scenarios run at 20°C to control scenarios run at 15°C.




Figure 4. Effect of warming 5°C on C stocks and flows in simulations, reported as the natural log of the ratio of values in heated compared to control conditions. CUE temperature response was forced to increase, decrease, or remain independent of the number of enzymes a taxon could produce. Values above the zero line indicate warming increased the value, and values below indicate a decrease with warming. Boxplots denote 1st to 3rd quartiles with the median. Asterisks denote significant warming effect at $P < 0.0001$ after correcting for multiple comparisons using the Bonferoni method. Letters denote warmed scenarios which are significantly different from one-another by the same criteria. “Reference temperature CUE” denotes the ratio of CUE for the surviving community at 15°C in warmed to the ratio of CUE for the surviving community at 15°C control scenarios, while “Simulation temperature CUE” denotes the ratio of CUE in heated scenarios run at 20°C to control scenarios run at 15°C.




Figure 5. Relationship between the enzyme cost and temperature sensitivity of surviving taxa when the relationship between C_t and enzyme count is set to be positive (increase), negative (decrease), or non-existent. Each point represents the value for the median across all surviving taxa for one of 59 starting communities.




Figure 6. Frequency histograms of Pearson correlation coefficients between the number of enzymes taxa which survive to the end of the 6000 day run and their C_r . Each panel corresponds to a different scenario shown elsewhere in the manuscript, with the “counts” in each bar corresponding to a single starting seed (community) under that scenario. Vertical lines denote the mean correlation coefficient for the scenario.




Figure 7. Effect of 5°C of warming on components of the litter C cycle when bacteria and fungi show dissimilar (F-B+ (orange), F+B- (yellow)) or similar (F+B+ (red), F-B- (blue)) CUE responses to temperature. Values above the zero line indicate warming increased the value (log ratio positive), and values below indicate a decrease with warming. Boxplots denote 1st to 3rd quartiles with the median. Asterisks denote significant warming effect at $P < 0.0001$ after correcting for multiple comparisons using the Bonferroni method. Letters denote warmed scenarios which are significantly different from one-another by the same criteria. “Reference temperature CUE” denotes the ratio of CUE for the surviving community at 15°C in warmed to the ratio of CUE for the surviving community at 15°C control scenarios, while “Simulation temperature CUE” denotes the ratio of CUE in heated scenarios run at 20°C to control scenarios run at 15°C. Values for “simulation temperature CUE” are also shown for bacteria and fungi separately.

Table 1. CUE-related model parameters mentioned in this paper. The complete set of parameters can be found in the params.txt file in the supplement.

Parameter	Value	Units	Description	Reference
C_r	0.38	dimensionless	CUE at 15°C for a taxon with no transporters or enzymes	this paper
C_e	-0.0025	enzyme ⁻¹	change in CUE per extracellular enzyme gene	Allison, 2014
C_u	-0.0071	transporter ⁻¹	change in CUE per transporter gene	Allison, 2014
C_t	-0.022 to 0.022	°C ⁻¹	change in CUE per degree change in temperature from 15°C	this paper

Table 2. Median (or median-standardized interquartile range (IQRm)) output values for various DEMENT model runs, marked according to warming effect (+/-) and model structure effects (letters) determined using Bonferroni-corrected post-hoc tests following linear mixed effect models. Symbols: "+" warming increased value; "-" warming decreased value). Letters: differences between warmed scenarios. All heated scenarios were compared to values in the "control" column to determine the warming effect. Only values within boxes defined by vertical lines were evaluated for significance differences of warming scenarios, as different seeds needed to be excluded for failing to constrain litter accumulation in the two boxes. NA indicates that fungi died out completely in many instances (i.e. median fungal biomass of zero), so the parameter output could not be determined. The number in brackets denotes the median excluding the scenarios where all the fungi died out.

Fig.1 scenario	NON-HEATED			CUE heterogeneity			CUE-Enzyme relationship			Fungi and bacteria differ			
	NA	A	B	C	D	E	F	G	H				
C_i ($^{\circ}C^{-1}$)	NA	0	-0.022 to 0.022	-0.022 to 0.022	-0.022 to 0.022	-0.022 to 0 (F) 0 to 0.022 (B)	-0.022 to 0 (B)	0 to 0.022 (F)	0 to 0.022 (F)				
enz_cost/ C_i relationship	NA	ind	ind	pos	neg	ind	ind	ind	ind				
LOM IQRm	0.2	0.14	0.28	0.33	0.15	0.39	0.29	0.23	0.3				
MBC IQRm	0.16	0.16	0.25	0.02	0.24	0.14	0.16	0.15	0.19				
Surviving taxa	9.5	11a	8b	13a+	7b-	12a	5c-	9b	8b-				
Enzyme count	16	16b	20.5b	34.5a+	10.8c-	17.5a	20.5a+	19a	18a				
Shannon's H	1.81	2.01a	1.58b-	2.11a+	1.31b-	1.93a	1.21c-	1.61a	1.64a-				
MBC ($mg\ cm^{-3}$)	20.9	21.0b	30.5a+	15.6c-	29.3a+	20.3b	33.0a+	32.3a+	17.2c-				
LOM ($mg\ cm^{-3}$)	583.2	496.0b-	361.5c-	220.1d-	603.2a	746.0a+	300b-	334.6b-	706.5a+				
CUE at 15°C	0.23	0.23b	0.23b	0.19c-	0.25a+	0.25a+	0.22c-	0.23b	0.23b				
CUE at 20°C	NA	0.23c	0.32a+	0.28b+	0.29b+	0.32bc+	0.31c+	0.32ab+	0.22d-				
FB biomass ratio	0.86	0.84b	0.81b-	0.00c-	0.88a	0.12b-	0.96a+	0.85a	0.73a-				
Respiration ($mg\ cm^{-3}\ day^{-1}$)	0.93	0.94b	0.96ab	0.99a+	0.88c-	0.91a	0.94a	0.96a	0.94a				
FB richness ratio	0.45	0.48a	0.44a	0.00b-	0.50a	0.14c-	0.67a+	0.44b	0.40b				
CUE bacteria	0.27	0.27b	0.26c-	0.18d-	0.36a+	0.35b+	0.26c-	0.36a+	0.25d-				
CUE fungus	0.22	0.23c	0.31a+	NA(0.28)b+	0.28b+	NA(0.18)b-	0.30a+	0.29a+	0.18b-				