



The last interglacial sea-level record of New Zealand (Aotearoa)

Deirdre D. Ryan^{1*}, Alastair J.H. Clement², Nathan R. Jankowski^{3,4}, Paolo Stocchi⁵

¹MARUM – Center for Marine Environmental Sciences, University of Bremen, Bremen, Germany

²School of Agriculture and Environment, Massey University, Palmerston North, New Zealand

5 ³Centre for Archeological Science, School of Earth, Atmospheric and Life Sciences, University of Wollongong, Wollongong, Australia

⁴Australian Research Council (ARC) Centre of Excellence for Australian Biodiversity and Heritage, University of Wollongong, Wollongong, Australia

10 ⁵NIOZ, Royal Netherlands Institute for Sea Research, Coastal Systems Department, and Utrecht University, PO Box 59 1790 AB Den Burg (Texel), The Netherlands

Correspondence to: Deirdre D. Ryan (dryan@marum.de)

Abstract: This paper presents the current state-of-knowledge of the New Zealand (Aotearoa) last interglacial (MIS 5 *sensu lato*) sea-level record compiled within the framework of the World Atlas of Last Interglacial Shorelines (WALIS) database. Seventy-seven total relative sea-level (RSL) indicators (direct, marine-, and terrestrial-limiting points), commonly in association with marine terraces, were identified from over 120 studies reviewed. Extensive coastal deformation around New Zealand has resulted in a significant range of elevation measurements on both the North Island (276.8 to 280.2 msl) and South Island (173.1 to 179.0 msl) and prompted the use of RSL indicators to estimate rates of vertical land movement; however, indicators lack adequate description and age constraint. Identified RSL indicators are correlated with MIS 5, MIS 5e, MIS 5c, and MIS 5a and indicate the potential for the New Zealand sea-level record to inform sea-level fluctuation and climatic change within MIS 5 (*sensu lato*). The Northland (North Island) and Otago (South Island) regions, historically considered stable, have the potential to provide a regional sea-level curve in a remote location of the South Pacific across broad degrees of latitude. Future work requires modern analogue information, heights above a defined sea-level datum, better stratigraphic descriptions, and use of improved geochronological methods.

25 The database presented in this study is available open-access at this link: <http://doi.org/10.5281/zenodo.4056376> (Ryan et al., 2020a).

1 Introduction

The New Zealand nation (Aotearoa in the Māori language) is an archipelago comprised of two large ‘main’ islands, nine outlying island groups, and hundreds of additional islands. The distribution of islands extends southward from the sub-tropics to sub-polar regions in the mid-latitudes of the Southern Hemisphere. The main islands, North Island (113,729 km²) and South Island (151,215 km²), are a small area of Zealandia micro-continental crust straddling the convergent boundary of the Australian and Pacific Plates. This geographic setting produces a relatively youthful and complex landscape subject to volcanism, glaciation, and tectonism. This latter characteristic of New Zealand has facilitated the preservation of marine terrace sequences spanning the Pleistocene and into the Holocene, which serve as the primary source of last interglacial paleo sea level records.

35 The impetus behind much of the marine terrace research completed in New Zealand (Pillans, 1999 and references therein; this review) has been to constrain long-term uplift rates using the many preserved sequences; an approach typically used on active margins (Creveling et al., 2015; Simms et al., 2016). However, recent advances in sea-level studies have highlighted the need for increased spatio- and temporal density of relative sea-level (RSL) indicators, analyzed using standard definitions and methods, to assist in constraining paleo ice sheet limits and to improve models of glacio- and hydro- isostatic adjustment (GIA)



40 and future predictions of ice-sheet and sea-level responses to regional warming (Dutton et al., 2015; Rovere et al., 2016; Austermann et al., 2017; Barlow et al., 2018; Capron et al., 2019; Golledge, 2020).

In this paper we use the standardized framework provided by the World Atlas of Last Interglacial Shorelines Database (WALIS, <https://warmcoasts.eu/world-atlas.html>) to build a database of previously published RSL indicators: their descriptions, geochronological constraint, and associated metadata. The database is available open-access at this link: <http://doi.org/10.5281/zenodo.4056376> (Ryan et al., 2020a). Database field descriptors are available at this link: <https://doi.org/10.5281/zenodo.3961543> (Rovere et al., 2020). In total, we reviewed over 120 publications and identified 77 unique RSL indicators of varying quality. Previous reviews of New Zealand sea-level indicators (Gage, 1953; Pillans, 1990a) have recommended, “more accurate height data, careful attention to shoreline datums, and greater use of dating techniques yielding numerical ages” (Pillans, 1990a, p. 227). This compilation summarizes the current knowledge of last interglacial (LIG) sea level in New Zealand by identifying RSL indicators within the literature, provides an overview of the standardized treatment of LIG RSL indicators using WALIS, identifies problems, and makes recommendations for the improvement of the NZ LIG record. Below we first provide an overview of the historical development of sea-level studies within New Zealand to provide context for the outcomes of this compilation. The following sections describe the types of RLS indicators identified in this work (Section 2) and how geographic location, elevation, and associated uncertainty were assigned to each indicator (Section 3). The bulk of this publication (Section 4) discusses the current state of the last interglacial sea-level record within New Zealand and provides detailed descriptions of the RSL indicators identified, dividing the discussion into sub-regions within the North and South Islands. In the last sections of the paper, we address the main sources of uncertainty (Section 5), and provide further details on the Pleistocene and Holocene sea-level record (Section 6), and suggest future research directions (Section 7).

60 1.1 Overview of literature with reference to last interglacial sea level

The legacy of New Zealand sea-level change studies in many ways reflects the evolution of eustatic sea-level and plate tectonic science. By the latter half of the 19th Century, “raised beaches” were well-recognized within the New Zealand landscape and investigations were subject to review and synthesis as early as 1885 (Hutton). Hutton (1885) attributed the differing altitude of the marine terraces to slow retreat of a maximum sea level exceeding 800 ft [~244 m], whereas Henderson (1924) argued for uniform changes in RSL around New Zealand as the result of either retreat of the sea or uplift of the entire archipelago. Subsequent studies revealed regional differences in marine terrace elevations attributed to differential vertical land movement (Jobberns, 1928; King, 1930; 1932). Although the cause for larger, regional variation was not fully understood, correlation with overseas marine terraces to determine age was considered “extremely hazardous” (Gage, 1953, p. 27). However, awareness of regional warping did not dissuade many authors from attempts to determine terrace age by ‘matching’ the New Zealand terraces with the well-recognized, yet far-field, terrace sequences in the Mediterranean (the Monastirian, Tyrrhenian, Milazzian, Sicilian sequence) (e.g. Brothers, 1954; Cotton, 1957; Leamy, 1958; Suggate, 1965) and on the Huon Peninsula, Papua New Guinea (Chappell, 1975; Bishop, 1985; Bull and Cooper, 1986). The former correlation also promoted the widespread use of Mediterranean sea-level nomenclature as a framework for New Zealand marine terraces for decades. These correlations were often attempted by ‘counting backwards’, with increasing altitude or distance from the modern shoreline. Prior to the development and common use of other geochronological methodologies (e.g. luminescence dating, amino acid racemization), this approach or the similarities between terrace cover beds were commonly applied to correlate distant terrace sequences within New Zealand (Kear and Waterhouse, 1961; Fleming and Suggate, 1964; Chappell, 1970; 1975; Nathan and Moar, 1975; Heine, 1979; 1982; Ota et al., 1984; 1996; Bishop, 1985; Richardson, 1985).

With the recognition of regional tectonics playing a large role in the altitude and distribution of the marine terraces, researchers shifted focus, using marine sequences primarily to assess regional long-term vertical land movements (but also to address other



research questions such as paleoclimate) and away from attempts to resolve sea-level fluctuations or eustatic sea-level history (e.g. Ghani, 1978; Hesp and Shepherd, 1978; Pillans, 1983; 1986; Bull and Copper, 1986; Ward, 1988a; Suggate, 1992; Berryman, 1993; Ota et al., 1996; Rees-Jones et al., 2000; Begg et al., 2004; Kim and Sutherland, 2004; Litchfield and Lian, 2004; Alloway et al., 2005; Cooper and Kostro, 2006; Wilson et al., 2007; Claessens et al., 2009; Clark et al., 2010; Oakley et al., 2018). However, this heralded the practice of correlating a presumed last interglacial terrace with a generic age (e.g. 120,000 ka) and height of the highstand (e.g. 5 m), usually derived from the marine oxygen-isotope records, with little effort to provide precise locations, elevations, or stratigraphic descriptions of sea-level indicators. While the development of geochronological methods assisted in constraining the age of marine terraces to the appropriate marine oxygen-isotope stage, common practice to estimate sea level continued to be the use of a eustatic sea level-height determined from various marine oxygen-isotope records. Another unfortunate outcome of previous studies is the disparate nomenclature for marine terraces and their sediments across physical space and time; the latter referring to both terrace age and research publication date (Table 1; Section 4).

In what may be considered a benefit from the prevalence of uplift along the New Zealand coast, numerous terrace sequences have been identified and correlated to the sea-level peak of MIS 5e, and subsequent interstadials MIS 5c and MIS 5a, or a combination of two of the three. Such sequences can be, and have been, used within New Zealand to study sea-level and climate fluctuation within MIS 5 *sensu lato* (e.g. Brothers, 1954; McGlone et al., 1984; Bussell, 1990; 1992; Berryman, 1992; Shulmeister et al., 1999); however, the majority of these studies were palynological with a focus on resolving climatic fluctuations and not sea level. Two regions of New Zealand have historically been considered tectonically stable (Gage, 1953; Pillans, 1990a; Beavan and Litchfield, 2012): the Northland Region of the North Island and the *Canterbury* Region of the South Island. Although these regions provide the best opportunity to identify a New Zealand sea-level record that can contribute to the discussion of MIS 5e sea level in a global context, only a few RSL indicators have been identified in these areas (Section 4).

This review of the LIG sea-level indicators in New Zealand, like the preceding reviews by Gage (1953) and Pillans (1990a), found considerable lack of detail in the published literature; including that published post-1990. Many locations are referred to broadly, with the only indication of place a large-scale location map from which a field site cannot be precisely determined. Marine terrace descriptions are often limited to a range of altitude heights in meters without a defined sea level datum - commonly the reader is left to assume above mean sea level. In many instances where a stratigraphic description is provided, it is logged as meters depth of burial and the height above sea level cannot be determined. The use of altitude to determine age, whether above an undefined sea level datum or between successive terraces, is prevalent. On the North Island, tephrochronology is commonly used both as a regional stratigraphic marker and to assist in constraining marine terrace age (e.g. Pullar and Grant-Mackie, 1972; Chappell, 1975; Pain, 1976; Iso et al., 1982; Ota et al., 1989; Berryman, 1993; Wilson et al., 2007; Claessens et al., 2009). For the purposes of determining last interglacial age the tephtras most commonly referred to are the Rotoehu Tephra at ~47 ka (Danišik et al., 2012; Flude and Storey, 2016), providing a minimum age, and the Hamilton Ash at ~340 ka (Pillans et al., 1996; Lowe et al., 2001) providing a maximum age (Section 5.3.2). The most commonly applied geochronological methods for determining numerical age are calibrated amino acid racemization and various techniques of luminescence dating, both of which have complicated histories in New Zealand (Section 5.3). In summary, in New Zealand there are few well-described and constrained RLS indicators in proportion to the sizable quantity of literature published concerning the physical record of last interglacial (*sensu lato*) sea level.

2 Types of sea-level indicators



The majority of LIG RSL indicators identified within New Zealand are associated with a marine terrace. The marine terrace terminology used here follows that established by Pillans (1990a; b) (Figure 1). Briefly described, a marine terrace is a gently sloping, sub-planar landform, consisting of a basal, sub-horizontal shore platform (a *wave-cut* surface) overlain by cover beds




Table 1: Changes in nomenclature and age correlation of interglacial sequences in North and South Islands of New Zealand						
	North Island	MIS 7	MIS 5e	MIS 5c	MIS 5a	Post MIS 5
Auckland - Northland	Brothers, 1954	Shelly Beach Formation (33.5 - 39.6 m)	Waioneke Formation (13.7 - 22.9 m)	Waioneke Formation (4.5- 7.6 m)		
	Chappell, 1975		Shelly Beach Formation (40 m)	Waioneke Formation		
	Richardson, 1985	South Head Formation	Shelly Beach Formation/Terrace (40 m)	Undifferentiated Hawera sediments (24 to 8 m)		
	**Nichol, 2002		One Tree Point (prograding beach face 5 +/- 4 m)			
Whanganui - Taranaki - Waikato	**Chappell, 1970	Waiau A Formation (marine sediments, max. height 39.6 m)	Waiau B Formation (isolated littoral beds, max. height of 12.9 m)	Waiau B Formation (isolated littoral beds, max. height of 12.9 m)		
	Chappell, 1975		Waiau A Formation (marine sediments, max. height 39.6 m)	NT2 (littoral and estuarine beds to 20 m)		
	Chappell, 1975		NT3 (shore platform 40-42 m)			
	Pain, 1976			Te Akeake Sands Member (Waiau B Formation)		
	Alloway et al, 2005	NT3	NT2		NT1	
	Dickson et al, 1974	Brunswick Formation/Terrace	Ngarino Formation/Terrace	Rapanui Formation / Terrace		
	Chappell, 1975		Ngarino Terrace	Rapanui Terrace		
	**Pillans, 1983	Ngarino Terrace (76-128 m)	Rapanui Terrace (<30-70 m)	Inaha Terrace (24-40 m)	Hauriri Terrace (16 m)	
Gisborne - Bay of Plenty	Chapman-Smith & Grant-Mackie, 1971	Otamaroa Terrace; or <i>younger</i> (63 m)	Te Papa Terrace; or <i>younger</i> (33 m)			
	Pullar & Grant-Mackie, 1972			Otamaroa Terrace	Te Papa Terrace	
	Chappell 1975	BOP3 (shore platform >50 m)	BOP2 (shore platform >20 m)			
	Yoshikawa et al, 1980	Matakaoa Terrace (20-130 m)	Otamaroa Terrace (10-300m)		Te Papa Terrace (10-215 m)	
	Ota et al, 1989		Otamaroa Terrace (terrace surface 68-287 m)	Raukokore Terrace (terrace surface minus 4 m for tephra, ≥214 m)	Te Papa Terrace (terrace surface minus 4 m for tephra, 28-92 m)	
	**Wilson et al, 2007		Otamaroa Terrace (62-279 m); Te Papa Terrace (62-279)			
Hawke's Bay	Cashman and Kelsey, 1990		Kidnappers Terrace (100-200 m)			
	**Berryman, 1993	Moemoto Terrace (88-230 m)	Mahia III (42-150 m)	Mahia II (51-150 m)	Mahia I (83-116 m)	Auroa 2 Terrace (63-64 m)
Wellington - Wairarapa	**Mildenhall, 1995		Biozone P6 of the Hutt Formation (depth below surface: 82.0-105.4 m decreasing landwards to 48-65 m below surface)			
	Begg et al, 2004		Willford Shellbed			
	Heine, 1982		MIS 5 terrace levels (40-42 m and 18-20 m)			
	Leamy, 1958	Tyrhenian (30.7 - 31.6 m)	Main Monastrian (15.5 - 16.5 m)	Late Monastrian (5.1 m amsl)		
	Cowie, 1961		Tokomaru Marine Terrace			
	**Fleming, 1972		Otaki Formation			
	**Palmer, 1988		Tokomaru Marine Terrace & Otaki Formation. Beach sand (~30 m amsl)			
	Sewell, 1991		Tokomaru Marine Terrace & Otaki Formation	Post Tokomaru Marine Terrace 1	Post Tokomaru Marine Terrace 2	
	South Island	MIS 7	MIS 5e	MIS 5c	MIS 5a	Post MIS 5
West Coast - Southland	Suggate, 1965	Karoro Formation		Awatuna Formation		
	Suggate, 1985	Karoro Formation	Rutherglen Formation	Awatuna Formation		
	**Nathan & Moar, 1975			Sardine-2 Terrace (24.4-32 m)		
	**Cooper & Kostro, 2006		Knights Point Terrace (113-110 m)			?Sardine-2 Terrace
	Bull & Cooper, 1986		MIS 5 Terrace(s) (686-899 m)			
	Bishop, 1985	h ₂ strandline (ave. 500 m)	h ₂ strandline (ave. 370 m)	h ₂ strandline (ave. 300 m)	h ₂ strandline (ave. 210 m)	
	Ward, 1988b	Terrace 5 (230 m)	Terrace 4 (140 m)	Terrace 3 (90 m)	Terrace 2 (60 m)	
Otago	Kim & Sutherland, 2004		MIS 5e strandline (ave. 65 m)			
	Cotton, 1957		40 ft terrace (marine sediments, max height 12.2 m)			
	**Rees-Jones et al. 2000				h ₂ (beach sand, ~9 m high sea level)	
	**Litchfield & Lian, 2004		h ₂ (beach sand, ~9 m high sea level)			
	**Litchfield & Lian, 2004		Blueskin Bay Terrace (4-8 m)			
Marlborough - Canterbury	**Kennedy et al, 2007				Shag Point shore platform (~7.4 m mean low water)	
	**Brown, 1988		Bromley Formation (79-5 m below msl)			
	**Shulmeister et al, 1999	Unit C (lagoon sediments, 61.5-48.2 m bmsl)		Unit J (beach deposit, 22.7-22.1 m bmsl)	Unit H (marine embayment, 34.1-24.6 m bmsl)	
	Suggate, 1965		Hurunui-Blythe Rivers Terrace (54.9 m)			
	Suggate, 1965	Bob's Flat (106.7-97.3 m)	Motunau Plain Lower Terrace (54.9-45.7 m)			
	Carr, 1970		Tiromoana Terrace (80-45 m)			
	**Oakley et al., 2017; 2018		Bob's Flat Terrace (128 m)	Tiromoana Terrace (63 m)	Motunau Terrace (marine fossil 70 m)	
	Suggate, 1965	Upper Terrace (?121.9-106.7 m)	Amuri Bluff Lower Terrace (48.8-45.7 m)			
	**Ota et al., 1984	Kemps Hill Terraces. Upper (240-220 m) Lower (170-150 m)	Amuri Bluff Terrace (100-40 m)			
	**Ota et al., 1996			Tarapuhi Terrace (165 m)	Kemps Upper (110 m)	Kemps Lower (90 m)
	**Oakley et al., 2017; 2018			Tarapuhi Terrace (173.1 m)	Kemps Hill Upper (98 m), Keps Hill Lower (90 m), Amuri Bluff	
	Suggate, 1965	200-170 ft terrace (60.9-51.8)				
	**Ota et al., 1996			Terrace 1 (105 m)	Terrace II (80 m)	Terrace III (60 m)
	Suggate, 1965	Parikawa Formation (152 m)				
	**Ota et al., 1996	UM Terrace (222-190 m)	MM Terrace (shore platform 143-105 m) = Parikawa Formation			
	**Ota et al., 1996		MM Terrace (inner margin 80-55 m) = Parikawa Formation			
	**indicates source of RSL indicator					

of marine and/or terrestrial sediments. Terrace risers, identified by their steeper slopes, are present at the inland and seaward margins of the terrace. The inland terrace riser is the surface expression of a fossil sea cliff, formed at the peak, or sustained height, of the marine transgression. This is also the location of the shoreline angle (or strandline), a wave erosion feature at the



125 intersection of the  platform and the base of the fossil sea cliff, which serves as the best indicator of the height and
 timing of peak sea-level, although it is seldom exposed. The height of the base of the inland terrace riser or the terrace surface
 is often used as an alternative; however, because the thickness of cover beds is often unknown these elevations are not
 necessarily a good indicator of sea level (Pillans, 1983). The seaward terrace riser identifies the sea cliff of a subsequent sea-
 level highstand. In this review, depending on whether the elevation for the marine terrace RSL indicator came from the basal
 130 platform, marine cover beds or terrestrial cover beds, the  indicator is identified as either a marine-limiting point, a direct
 indicator, or terrestrial-limiting point, respectively. Marine sediments for which the depositional environment hasn't been
 constrained and/or are located an undefined distance from the inland terrace riser, are considered marine-limiting. Marine
 sediments described as beach deposits are considered to have been deposited between the storm wave swash zone and breaking
 depth of waves; i.e. within the upper and lower limits of the indicative range of a marine terrace (Rovere et al., 2016).

135 New Zealand RSL indicators have also been identified at depths below modern sea level. These indicators, from within
 sediment and well cores, are described as sediment packages that vary by location. The depositional environment with which
 those sediment facies were correlated dictate how the datapoint is identified in this review; i.e. a marine- or terrestrial-limiting
 point or a sea-level indicator (*sensu stricto*).

Modern analog information, used to determine indicative meaning (position relative to sea level at the time of formation) of a
 140 direct RSL indicator, was provided in only one study from the Northland Region (Nichol, 2002). For all other indicators, the
 indicative meaning was quantified using the  tool (Lorscheid and Rovere, 2019).




Figure 1: Illustration of marine terrace terminology (adapted from Pillans, 1990b) with storm wave swash height and breaking depth marking the upper and lower limit of indicative range for a marine terrace landform (Rovere et al., 2016).

3 Location and elevation measurements

145 The latitude and longitude for over half of the RSL indicators was determined using Google Earth to match locations from a
 publication map. The location for each well core described by Brown et al. (1988) was acquired from the Canterbury Regional
 Council well database (www.ecan.govt.nz/data/well-search; Environment Canterbury, 2020), which also provides accuracy of
 well elevation measurement above mean sea level. This value was incorporated into the elevation uncertainty provided by
 Brown et al. (1988). The other primary method used in publications for assigning location information has been New Zealand



150 Map Grid coordinates. In such instances, New Zealand map grid coordinates were converted to WGS84 (G1762) via the Land Information New Zealand online tool for coordinate conversions (<https://www.geodesy.linz.govt.nz/concord/index.cgi>; LINZ, 2020a). Oakley et al. (2017) determined locations with handheld GPS. Although Wilson et al. (2007) measured elevation using real-time kinetic GPS, locations were provided as map grid coordinates.

Three publications (Rees-Jones et al., 2000; McGlone et al., 2004; Kennedy et al., 2007) referred elevation data to a tidal
 155 datum other than mean sea level (msl). All other publications either referred to present or mean sea level or did not define the sea-level datum and only referred to sea level, in which case mean sea level has been assumed. The mean sea level definition has not been differentiated in the below identification (Section 4) of RSL indicators, but is shown in the database (<http://doi.org/10.5281/zenodo.4056376> Ryan et al., 2020a). The most commonly used method of elevation measurement has been altimetry, followed closely by metered tape or rod. In almost all cases of altimetry, the accuracy of measurement is
 160 considered to be within <5 m. The accuracy of the metered method, most commonly applied to core sediments, ranges from within <0.1 m to <5 m (Brown et al., 1988; Shulmeister et al., 1999). Less common means of determining elevation include differential GPS (Wilson et al., 2007; Oakley et al., 2017), total station (Kennedy et al., 2007), and a combination of topographic maps and digital elevation models (Wilson et al., 2007). In one instance, depression of the sea horizon in relation to the location of measurement was used (Ghani, 1978).

165 Similar to the sea level datum, in many instances the elevation measurement method is not stated. The intended approach for assigning elevation uncertainty with entry of an RSL indicator into WALIS was to determine elevation uncertainty by calculating the square root of the sum of the sea level datum error and adding a percentage of uncertainty of the elevation measurement depending upon the precision of the method used. If the sea level datum error was not provided, then the elevation uncertainty would be determined as 20% of the elevation. However, these approaches were not possible due to the lack of
 170 stated sea level datum and elevation measurement method in many studies. Furthermore, given the variable rates of uplift along the New Zealand coast, applying 20% of the elevation would result in uncertainty in the range of tens of meters in some instances. Instead, if an uncertainty for a described elevation method was provided in the original publication, this was accepted and applied with entry of the RSL indicator into WALIS. All other instances were assessed on a case-by-case basis and uncertainty was defined dependent upon the quality of the indicator description provided within the original publication. None
 175 of the elevations provided here have been adjusted for the effects of uplift, GIA, or dynamic topography.

4 Relative sea-level indicators

The approach here for describing RSL indicators will continue the practice of dividing New Zealand between the North and South Islands, with further subdivision based roughly upon the government regions and the dominant tectonic regimes (Figure 2). This approach was chosen for clarity in discussion and in recognition that coastal deformation in New Zealand is driven by
 180 the tectonic regimes resulting from the position of the archipelago over the active boundary between the Australian and Pacific Plates. The obliquely convergent plate boundary can be subdivided into three distinct components from north to south: (1) the obliquely westward-subducting Hikurangi Margin east of the North Island, (2) a transitional zone from subduction to dextral transpression and oblique continental extension along the Marlborough Fault System and the Alpine Fault, and (3) the northeastward-subducting Fijian subduction system (see Nicol et al., 2017 for a detailed review of Quaternary tectonics of
 185 New Zealand, and Litchfield et al., 2014 for a detailed review of active faulting in New Zealand).

The WALIS database identifiers for RSL indicators (e.g. WID 43) will be used throughout this section to assist in correlation between this publication and the database (<http://doi.org/10.5281/zenodo.4056376> Ryan et al., 2020a), within which they are




Figure 2: The New Zealand North and South Islands illustrated with government regions and dominant tectonic regimes as discerned by color-shading. Also shown are the primary faults of the North Island Dextral Fault Belt, the Marlborough Fault System, and the Alpine Fault (Langridge et al., 2016). Diamonds indicate locations of RSL indicators within the WALIS database and described here. Hill shade data sourced from the LINZ Data Service and licensed for reuse under CC BY 4.0 (LINZ, 2020b).

found under the heading “WALIS RSL_ID”. Unique identifiers for amino acid racemization and luminescence samples are as provided in original publication, but can also be correlated with a unique WALIS identifier within the database. Figures 3 and 4 provide spatial context for RSL indicators and illustrate measured elevation with relation to sea level datum and age attribution.

4.1 North Island

The North Island features distinctive tectonic and geomorphic domains manifested by the continental collision processes associated with the subduction of the Pacific Plate along the Hikurangi Margin (Figure 2). The entire eastern North Island constitutes the Hikurangi Margin forearc that is compressing and uplifting (Nichol et al., 2017). The forearc structure from east to west is composed of an accretionary wedge (outer forearc, mostly located offshore), a forearc basin (inner forearc), and the Axial Ranges (frontal ridge) bisected by the North Island Dextral Fault Belt (NIDFB) (Berryman, 1988). The oblique subduction of the Pacific Plate is rotating the eastern North Island clockwise producing a backarc rift system and volcanic arc:



the Taupo Volcanic Zone (TVZ). South of the TVZ lies the Whanganui Basin, a proto-back-arc sedimentary basin infilled with 4-5 km of dominantly marine Plio-Pleistocene sediment (Anderton, 1981). The western and northern North Island is a zone of backarc extension that normal faults have segmented into reticular blocks of uplifted basement rock and sedimentary basins. The backarc region also features extinct volcanic arcs west of the TVZ left relict by the clockwise rotation of the Hikurangi subduction system (Wallace et al., 2004).

The North Island is subdivided into nine government regions (Figure 2). Much of the north and west coasts are found within the Northland, Auckland, and Waikato regions. The northeast and east coasts are found within the Bay of Plenty, Gisborne, and Hawke's Bay regions with the southernmost coast of the North Island located within the Wellington region. The government region of Whanganui-Manawatu extends to both the southern and eastern coastline of the North Island, with the eastern coastline colloquially referred to as the Wairarapa within New Zealand. This practice is applied here to avoid confusion over reference to geographic location within the review.

4.1.1 Northland and Auckland regions

Portions of the Northland and Auckland regions have been considered relatively tectonically stable through the late Quaternary (Gage, 1953; Gibb, 1986; Pillans, 1990a; Beavan and Litchfield, 2012). Arguably the best described and constrained RSL indicator in New Zealand is the regressive nearshore, beach, and foredune sequence exposed at One Tree Point on the southern shoreline of Whangarei Harbour (Figure D 34; Nichol, 2002). Nichol (2002) traced the contact between the beach and foredune facies (delineated by a heavy mineral sand interpreted as the hightide swash deposit) for a distance of 3.4 km. The contact elevation decreases seawards from a maximum height of +6 m to +3 m above mean sea level (amsl), reflecting relative sea-level fall. Ground penetrating radar across the sequence revealed the swash lamination of the beach face and supports the interpretation of a prograding barrier sequence. Thermoluminescence (TL) samples of overlying foredune sand indicate deposition through MIS 5 (substages 5e to 5a). Given the height distribution of the swash deposit, and under the assumption that no tectonic uplift has occurred, Nichol (2002) argued for deposition during late MIS 5e. The height of the swash deposit associated with the TL sample providing MIS 5e age (Sample OTP1, 115 ± 19 ka) is 4.6 ± 0.92 m amsl.

The One Tree Point RSL indicator is the only one identified in the Northland and Auckland regions. Other remnants of MIS 5 sea level have been recorded, but lack sufficient detail to be used as RSL indicators. A series of Master's theses investigated the Quaternary geology and geomorphology of the Aupouri and Karikari Peninsulas in Northland, where possible MIS 5 marine terraces and estuarine sediments were identified (Goldie, 1975; Ricketts, 1975; Hicks, 1975). However, the correlations were considered tentative in appreciation for a lack of dating methods at the time and the dangers of distant terrace correlation or altimetry to determine age.

The west coast of Northland and Auckland has been a location of extensive sand deposition throughout most of the Quaternary and large volumes of these sands are subject to movement by nearshore currents and longshore drift with considerable erosion and deposition documented in historical record (Ballance and Williams, 1992; Blue and Kench, 2017). It is possible that remnants of the MIS 5 highstand may not exist outside of the more sheltered harbours. On the South Kaipara Peninsula, within the Kaipara Harbour and protected from the high energy coastline, Brothers (1954) identified and described the estuarine Shelly Beach and Waioneke Formations, the two youngest members of the Kaihu Group. The Shelly Beach Formation occurs as a marine terrace, with the surface recorded at 33 to 40 m amsl on the northern end and eastern side of the South Kaipara Peninsula. The Waioneke Formation is comprised mainly of pumiceous silts and localized sandy facies that infill valleys cut into the Shelly Beach and older formations. The terraces surfaces formed upon Waioneke Formation are typically found at 14 m to 23 m and 4.5 m to 7.5 m amsl. Brothers (1954) initially correlated the Shelly Beach Formation to the mid-Pleistocene and the younger Waioneke Formation to the late Pleistocene. Later assignment of the Shelly Beach Formation to MIS 5 was




Figure 3: Position, elevation, and age correlation of the North Island RSL indicators within the WALIS database and described here. Each RSL indicator is identified with their unique WALIS database identifiers (e.g. WID 34) and is correlated in text with their original publication. Symbols of indicators that would stack extend offshore for clarity. Marine and terrestrial limiting points are indicated with arrows. Locations and landscape features mentioned in text are also shown with symbols (e.g. towns and cities) indicated by stars. Hill shade data sourced from the LINZ Data Service and licensed for reuse under CC BY 4.0 (LINZ, 2020b).

made by Chappell (1975; Table 1). Richardson (1985) identified the Shelly Beach Formation on the southern shoreline of the Pouto Peninsula (on the northern side of the harbour), often occurring with a 40 m terrace. Chappell (1975) and Richardson (1985) determined the MIS 5 age of the Shelly Beach Formation by correlating it with the Ngarino Terrace in the Whanganui Basin to the south (Section 4.1.2). However, the Ngarino Terrace was later correlated with MIS 7 (Pillans, 1983; 1990b). Further south, Claessens et al. (2009) claim to have identified a MIS 5 terrace at 41.5 m amsl within the Waitakere Ranges, on the coastline north of the Manukau Harbour entrance; however, the stratigraphic description and geochronological constraint are poor and the publication was rejected for inclusion into the database.

4.1.2 Waikato, Taranaki and the Whanganui Basin

Chappell (1970) redefined the Kaihu Group, extending its framework to encompass the Waikato region, and correlated the Waiau A and Waiau B Formations in the Waikato with the northern Shelly Beach and Waioneke Formations, respectively, on the basis of elevation and position within the Kaihu Group. The sediments of both Waiau A and Waiau B are described as forming terraces overlying irregular erosion surfaces within coastal harbours and valleys. Waiau A is composed of transgressive marine sediments forming a terrace with a maximum surface elevation of ~40 m amsl; however, a detailed stratigraphic description of the Waiau A beds at a defined location is not provided and an RSL indicator could not be identified.



The Waiau A beds are made distinct from the overlying Waiau B aeolian and isolated littoral beds by an erosion surface. The surface of the Waiau B littoral beds is not found above 13 m amsl elevation; however, associated aeolian dunes exceed 106 m amsl. At the Waiau B representative section, aeolian beds are drawn to directly overlie the soil-mantled erosion surface of Waiau A at ~16.7 m amsl (Figure 7 Chappell, 1970), providing a terrestrial-limiting point (Figure 3 WDD 35). Remnants of these terraces, identified as North Taranaki 3 and North Taranaki 2 (Waiau A and Waiau B, respectively) in the northern Taranaki region, are used to correlate the west coast terraces to the marine terraces of the Whanganui Basin further south (Chappell, 1975; Table 1).

The south Taranaki coast, located on the western margin of the Whanganui Basin, is experiencing ongoing, gentle uplift due to crustal flexure as the basin depocenter migrates south (Anderton, 1981). The movement has allowed preservation of a globally significant shallow marine sequence, spanning the entire Quaternary, that has been mapped and described by multiple authors (Dickson et al., 1974; Chappell, 1975), but most extensively by Pillans (1983; 1990b; 2017). Marine terraces, formed over the past ~700 ka, provide a robust framework of Late Quaternary sea-level fluctuations. The marine terraces consist of a basal wide shore platform overlain by up to 15 m of marine sediments, grading upwards into non-marine sediments. The total thickness of terrace cover beds generally increases with terrace age and in the westward direction towards the Taranaki volcanoes due to greater tephra and lahar cover bed thickness; however, there can be considerable local variation due to sand dunes. The MIS 5 terraces are designated as the Rapanui Terrace (120 ka, MIS 5e), the Inaha Terrace (100 ka, MIS 5c), and the Hauriri Terrace (80 ka, MIS 5a). The uplift, which allowed preservation of the terraces, has also resulted in shore-parallel deformation of the terraces. The inferred strandline elevations (Rapanui Terrace <30-70 m, Inaha Terrace 24-40 m, Hauriri Terrace 16 m) must be viewed with additional caution due to the unknown thickness of the cover beds.

Age constraint for the Whanganui Basin sequence was provided by Pillans (1983; 1990b) using a combination of tephrochronology, specifically the Rangitawa Tephra, and amino acid racemization of wood samples (see Section 5.3.3 for discussion of AAR). RSL indicator elevations for each of the MIS 5 terraces were derived from type sections exposed on the modern coast. Most type sections are kilometers seaward of the inland terrace riser and indicate only minimum sea-level estimates for their respective highstands. The shore platform of the Rapanui Terrace (MIS 5e) is found within a cliff section at Castlecliff at 29 ± 0.3 m amsl and is directly overlain by ~5.5 m of marine cover beds (Bussell et al., 1992; Figure 3 WDD 38). An additional 7.5 m of terrestrial cover bed includes dune sand correlated to MIS 5c. It is unclear which depositional environment the marine sediments should be correlated with; therefore, the shore platform is considered as a marine-limiting point. The Rapanui Terrace is again exposed 7.5 km west of Hawera, with the shore platform measured to 12 ± 0.3 m above the high-water mark (HWM; McGlone et al., 1984; Figure 3 WDD 37). The type section for the Inaha Terrace (MIS 5c) is found 10 km west of Hawera, where the shore platform, at ~2 m above the HWM is overlain by near-shore marine and beach sands at 2.3 m to 4.17 m above HWM (McGlone et al., 1984; Figure 3 WDD 36). Given the proximity of this site to the inland terrace riser (within 2 km), the beach sands are considered a direct RSL indicator. The shore platform of the Inaha Terrace can be traced eastwards within the modern coastal cliff to a height of ~8 m above HWM. The shore platform of the Hauriri Terrace (MIS 5a) is exposed at 2 ± 2 m amsl near Waipara and overlain by ~2 m of fossiliferous marine sand with a basal conglomerate (Pillans, 1990b; Figure 3 WDD 245). Again, it is unclear which depositional environment the marine sediments should be correlated with and the shore platform is considered a marine-limiting point.

Chappell (1970; 1975) correlated the marine terrace sequences from the Whanganui Basin to northern Taranaki, Waikato, and Auckland Regions prior to any numerical age constraint and largely depended upon terrace elevation and tephrochronology. Correlation of the Waiau A and Waiau B Formations to the Whanganui sequence was reinforced through use of the Miocene-age Kaawa Shellbed as a chronostratigraphic marker and similarities in climatic and sea-level change signals perceived within the Whanganui sequence and the Kaihu Group (Chappell, 1970). In summary, Chappell (1975) correlated, from north to south: the Shelly Beach Formation, Waiau A, North Taranaki (NT) 3, and the Ngarino Terrace to MIS 5e and implied the Waioneke



305 Formation, Waiau B, North Taranaki (NT) 2, and the Rapanui Terrace as correlatives formed during a later sea-level highstand
 (Table 1). To determine age Chappell (1975) used the current heights of the Ngarino and Rapanui Terrace surfaces and
 estimates of uplift rates to correlate the terraces with the Huon Peninsula (Papua New Guinea) sea-level curve and assign an
 age of ~120 000 years for MIS 5e (Chappell, 1974; Bloom et al., 1974). Pillans (1983; 1990b) later correlated the Ngarino
 Terrace to MIS 7 and the Rapanui Terrace to MIS 5e. The accuracy of Chappell's terrace correlation across most of the west
 310 coast of the North Island otherwise remains untested and has been used for geochronological constraint in subsequent
 publications, e.g. Alloway et al. (2005).

4.1.3 Bay of Plenty and Gisborne regions

The central portion of the Bay of Plenty is defined by a subsiding backarc rift occupied by the Taupo Volcanic Zone (Figure
 2). The coastline marginal to the rift is experiencing moderated uplift. To the east, the Raukumara Peninsula, a northwards
 315 projection of the Axial Ranges, is subject to steady aseismic uplift, with some intermittent coseismic uplift events (Chapman
 et al., 2010).





Chappell (1975) attempted to correlate the marine terraces of the west coast of the North Island with new and previously
 described terrace remnants across the Bay of Plenty and Raukumara Peninsula, identified as Bay of Plenty 2 (BOP2) and Bay
 of Plenty 3 (BOP3; Table 1; Kear and Waterhouse, 1961; Selby et al., 1971; Chapman-Smith and Grant-Mackie, 1971; Pullar
 and Grant-Mackie, 1972). Briggs et al. (1996; 2006) showed that the Pleistocene terraces identified in the western Bay of Plenty, with
 320 which Chappell (1975) attempted correlation, are non-marine in origin. The marine terraces on the Raukumara Peninsula were
 first identified as the Otamaroa and Te Papa Members of the Rukuhanga Formation, described at a location near Cape Runaway
 (Chapman-Smith and Grant-Mackie, 1971). The terrace cover beds are comprised of marine sandstones and conglomerates
 overlain by tephra. After subtracting 3 m for tephra cover bed thickness, the inner margins of the terraces were measured to
 325 60 m amsl (Otamaroa Terrace) and 30 m amsl (Te Papa Terrace) (Pullar and Grant-Mackie, 1972). The tephra overlying the
 Otamaroa Terrace was tentatively correlated to the Hamilton Tephra and the tephra overlying the Te Papa Terrace was
 identified as Rotoehu Tephra. Both tephra were used to argue for the formation of the terraces during interstadials equivalent
 to the European Brörup (MIS 5c) and later Gottweig interstadials, respectively. Chappell (1975) incorporated the Te Papa
 Terrace into his Bay of Plenty 2 Terrace, which was correlated with west coast MIS 5 terraces in the Auckland and Waikato
 330 regions. The Otamaroa Terrace was incorporated in the Bay of Plenty 3 Terrace and correlated to MIS 7. Per Wilson et al.
 (2007), the Otamaroa and Te Papa Terraces were later correlated by Yoshikawa et al. (1980) to MIS 5c and MIS 5a,
 respectively; however, the publication is in Japanese and we were not able to confirm the correlation.

Wilson et al. (2007) extensively mapped the Te Papa and Otamaroa Terraces from Whitianga Bay, eastward around the
 Raukumara Peninsula to Te Araroa, providing section descriptions and elevations measured to heights above local mean sea
 335 level. The position of the described sections within the terrace, in relation to the inner and outer margins, is not provided. The
 cover beds overlying the shore platform are consistently comprised of marine sediments of sand and/or gravel, never exceeding
 6 m thickness, overlain by terrestrial sediments, typically less than 4 m thick, with the Rotoehu Tephra serving as a
 chronostratigraphic marker. Shore platform elevation measurements from this work provides one marine-limiting point for the
 Te Papa Terrace at Waihu Bay (22.3 ± 0.55 m amsl; Figure 4 ID 46) and four marine-limiting points for the Otamaroa
 340 Terrace: Omaio (11.5 ± 0.50 m; Figure 3 WID 43), Waihu Bay (61.9 ± 0.50 m; Figure 4 ID 45), Hicks Bay (129.9 ± 1.25
 m; Figure 4 ID 42), and Te Araroa (276.8 ± 10.0 m; Figure 4 ID 44). Wilson et al. (2007) also collected samples for
 infrared stimulated luminescence (IRSL) dating from terrace sand and loess cover beds to resolve the age of the terraces. The
 IRSL ages suggest correlation of the Otamaroa Terrace with MIS 5a and the Te Papa Terrace with MIS 3, implying MIS 5e
 and 5c terraces are not present along the northern Raukumara Peninsula. Wilson et al. (2007) found these results unsatisfactory.
 345 The MIS 5a and MIS 3 correlation was considered unlikely given the higher relative sea levels during the earlier highstands



and the consistent uplift of the peninsula. They also argue development of the terraces within MIS 5 is consistent with regional loess chronology and geomorphological characteristics of Pleistocene marine terraces. It is noted that, IRSL methods have developed significantly since this publication and it is likely the methods used for this work produced underestimated ages (Section 5.3.4).

350 4.1.4 Hawke's Bay region

Mahia Peninsula, in the Hawke's Bay region, is subject to coastal deformation and uplift due to its position within the accretionary wedge of the Hikurangi Margin forearc and its proximity to the Lachlan Anticline, the axis of which is directly offshore (Figure 2). Berryman (1993) provided a comprehensive description of the peninsula and its  marine terraces, of which three were correlated to MIS 5: Mahia III (5e), Mahia II (5c), and Mahia I (5a). All three terraces are identified on the
 355 northeast part of the peninsula. Mahia III is also found along the southwest coastline and across Portland Island directly to the south. The terraces are composed of shore platforms cut into underlying bedrock with overlying marine sand cover beds. The marine sands grade upward into aeolian dune sands, in turn overlain by sequences of tephra and loess. The cover beds are described as varying in thickness from approximately 4 m for the lower terraces to 20 m for the higher terraces, but which terraces are considered lower and higher is not stated. Berryman (1993) correlated Mahia III to the MIS 5e peak at 124 ka,
 360 citing an amino acid minimum age for a wood sample (Pillans, 1990a) and palynological evidence (personal communication with Matt McGlone). The stratigraphy of the tephra-loess cover beds was used to determine the age of the remaining terraces. Mahia I, II, and III were differentiated from the five other marine terraces on the peninsula by the composition of their cover beds, which include three loess units with intervening deposits of Kawakawa Tephra and Rotoehu Tephra. Berryman (1993) mapped numerous spot altitudes of shoreline angle positions, which were determined by subtracting a mean, undefined,
 365 thickness for terrestrial cover beds. Due to the deformation of the peninsula, the elevation measurements from each terrace vary significantly: Mahia I (83-116 m amsl), Mahia II (51-150 m amsl), northwest Mahia III (130-150 m amsl), southwest Mahia III (42-132 m amsl). The shoreline angle elevations used by Berryman (1993) to calculate uplift rates, chosen for their position approximately parallel to the strike of the Lachlan Anticline, are considered direct RSL indicators: Mahia I (92 ± 10 m amsl; Figure  39), Mahia II (124 ± 10 m amsl; Figure  40), and Mahia III (147 ± 10 m amsl; Figure  41).

370 The uplifting Cape Kidnappers, on the southern margin of Hawke's Bay and opposite of the Mahia Peninsula, is also within the accretionary wedge of the Hikurangi Margin forearc and is bisected by the Kidnappers Anticline. Cashman and Kelsey (1990) reference the Kidnappers Terrace, described as ~10 m of marine cover beds overlying a wave-cut shore platform and correlated to MIS 5. The elevation of the terrace surface varies from 200 m amsl at the axis of the Kidnappers Anticline, to 100 m amsl on either side. Additional details are not available; of the references referred to by Cahsman and Kelsey (1990),
 375 King (1932) provided insufficient information to define a RSL indicator, the remaining Kingma (1971) and unpublished Masters Theses (Kamp, 1978; Hull, 1985) were not made available in the course of this work.

4.1.5 Wairarapa and Wellington regions

380 The Wairarapa coast, south of Hawke's Bay, is subject to uplift and deformation through folding and faulting as it forms the southern portion of the accretionary wedge of the Hikurangi Margin forearc (Figure 2). Further south, deformation of the Wellington coast is driven by the northeast-trending strike-slip faults of the North Island Dextral Fault Belt that have uplifted basement blocks to form the Axial Ranges with intervening basins developed through differential subsidence (e.g. Wellington Harbour-Port Nicholson and the Hutt Valley) (Begg and Johnston, 2000; Lee et al., 2002). Uplifted marine terraces and subsurface marine sediments identified in well cores have been correlated to MIS 5 in multiple locations; however, few have been sufficiently described to qualify as an RSL indicator.



385 Ghani (1978) mapped four marine terraces in four locations between Riversdale and Wellington City, referring to the sequence
 as the Eparaima Marine Benches, subdivided from lowest to highest as EA, EB, EC, and ED. The best-preserved sequence is
 between Riversdale and Flat Point on the southeast coast. The remaining three locations, Cape Palliser, the lower Ruamahanga
 Valley, and the upper Ruamahanga Valley, are on the south coast and retain varying extents of the terraces, with EC the best
 preserved. The terraces are described as consisting of a wave-cut surface with overlying beach deposits and loess of undefined
 390 thickness. The age of each terrace is constrained by the number of “soil-units”: two soil-units are found on the lower terraces
 (EA and EB, 80 ka and 84 ka, respectively), three on EC (100 ka), and four on ED (125 ka); however, the two oldest soil-units
 are not consistently present and are limited in use for spatial correlation of the terraces. Mapping of the terraces was done from
 ground-truthing of features identified in aerial photographs. Primary spot heights were measured by determining the depression
 of the sea horizon. Secondary spot heights were determined from vertical angles and distances from 20-chains-to-an-inch maps
 395 [1 in = ¼ mile]; overall spot height uncertainty was determined by Ghani (1978) to be ± 3 m. The spot heights were then used
 to map the stranded shorelines and contour the outcrop pattern of each terrace from which vertical crustal movements were
 derived. None of the spot heights were added to the database due to concerns for precision of location and elevation, unknown
 thickness of cover beds, method of age constraint, and the extensive deformation of the coastline - numerous faults, anticlines
 and synclines run parallel and perpendicular to the coast. Between the four sequence locations, terrace surface elevation for
 400 each terrace varied in the tens of meters; however, the only surface to exceed 200 m amsl is ED.

Mildenhall (1995) analyzed the pollen profile of coastal sediment sequences retrieved from two drillholes (Petone and
 Seaview) in the Lower Hutt Valley and identified nine Biozones. Biozone P6 was tentatively correlated to the Last Interglacial
 as the taxa present are currently restricted to the north of the North Island and indicate a period of maximum warmth.
 Postglacial radiocarbon ages from the overlying Biozone P9 and the glacial climates indicated from the intervening Biozones
 405 P7 and P8 are used to strengthen the correlation to MIS 5. The sediment lithology from both drillholes, inclusive of marine
 shell, indicates a marine environment; however, proximity to the paleo shoreline is unknown. Biozone P6 is found within the
 Petone drillhole at 82.0 m to 105.4 m depth of burial from the land surface, which is ~ 1 m amsl. The biozone may extend
 higher within the sequence as there is a 6 m sampling gap between this biozone and the overlying Biozone P5. The sediments
 of Biozone P6 thin and increase in elevation to the east where they are found within the Seaview drillhole at 48 m to 65 m
 410 depth of burial from the land surface, ~ 1 m amsl. Biozone P6 was later referred to as the Wilford Shellbed, assigned an age of
 128 ka to 71 ka, and used in estimates of subsidence rates within the Lower Hutt Valley (Begg et al., 2004). Due to the
 uncertainty in depositional environment, Biozone P6 is considered marine-limiting (Figure 568 and 569).

Heine (1974; 1979; 1982) identified numerous terrace surfaces in and around Wellington City. In the latter two studies, Heine
 used a method described as, “a micro-survey of the topography to identify any systematic pattern in the elevations” (Heine,
 415 1979 p. 379), to deduce any relationship between coastal terraces and local tectonics. Over 350 elevation surfaces above mean
 sea level were identified within the city of Wellington using altimetry and from topographic maps with the only criteria that
 the surface was “near to level”. The spot heights were then subdivided into 56 discrete ‘levels’ within which ten ‘major’ terrace
 levels were identified and tentative correlation to other New Zealand marine terraces were made on the basis of terrace altitude.
 Heine (1982) expanded the spatial extent of this approach to include the entire Wellington Peninsula and suggested levels up
 420 to 200 m to be marine in origin with correlations with MIS 5 (18-20 m and 40-42 m), MIS 7 (70 m), and MIS 9 (107 m)
 highstands. The intermediate levels were attributed to sea-level fluctuations. Although spot heights are provided in publication
 maps, terrace descriptions as basement rock overlain by Quaternary terrestrial sediments of undefined thickness, and the
 tentative age correlation precluded the identification of a RSL indicator.

Leamy (1958) described a number of terraces within Porirua Harbour, an embayment in the coast north of Wellington City.
 425 Two terraces, described as of “probable” marine origin underlain by gravels, were correlated by height to European Main
 Monastirian and Late Monastirian, recognized as two stages of the Last Interglacial (*sensu lato*; Table 1). Aneroid barometer



measurements of the terrace surfaces placed the Main Monastirian terrace surface at 16.5 m to 15.5 m amsl. The Late Monastirian terrace surface was measured to 5.1 m amsl. These datapoints were not added to the database due to the uncertainty of their origin (fluvial or marine) and age. Webby (1964) described additional sections around the harbour identifying “red-weathering horizons”, of which some were correlated with the Last Interglacial. Estuarine sediments underlying a 60 ft [18.3 m] marine terrace correlated with the Last Interglacial are identified in section; however, the stratigraphic column is a composite with the elevation of the sequence surface described as varying between 3.0 and 26.5 m and the location was not included as a RSL indicator. Offshore and slightly to the south of the Porirua Harbour mouth, multiple terraces on Mana Island have been described as marine in origin (Williams, 1978). Although they are attributed to Pleistocene high sea levels, no other age constraint has been provided.

4.1.6 Southern Manawatu-Whanganui and northwestern Wellington region

West of the Axial Ranges, the onshore eastern margin of the Whanganui Basin begins as a narrow coastal plain at Paekakariki and broadens northward into the southern Manawatu-Whanganui region to form the Horowhenua lowlands (Begg and Johnston, 2000). Spatially variable deformation within the region is associated with faulting of numerous basement faults of the North Island Dextral Fault Belt and overall regional uplift of the ranges (Sewell, 1991; Begg and Johnston, 2000). Early studies of the region led to the identification two deposits of interest: the Otaki Formation (Oliver, 1948) and the Tokomaru Marine Terrace (Cowie, 1961), which were not explicitly correlated until 1988 (Palmer et al.).

Oliver (1948) formalized the Otaki Formation (originally Otaki Sandstone) designation for Late Pleistocene marine, beach, and dune sands deposited in the coastal region between Paekakariki and Palmerston North (Figure 3). The flat surface overlying the Otaki Formation was attributed to post-depositional erosion and the surface does not appear to be recognized as a marine terrace until Palmer et al. (1988). An exposure within the outer, seaward terrace riser allowed for a detailed lithological description of the Otaki Formation by Te Punga (1962); however, with limited paleoenvironmental interpretation and no effort for lateral correlation of the sequence with other deposits in the region. Fleming (1972) summarized the sequence as consisting of a wave-cut shore platform upon which is emplaced a sequence of transgressive marine beach gravel and sands (basal Otaki) that transition to beach-derived, micaceous dune sands (upper Otaki) deposited as sea level fell. Awatea Lignite was deposited by swamps ponded within the dunes. The Otaki Formation was correlated to the Last Interglacial at ~80,000 to 120,000 years based upon the minimum radiocarbon age of the lignite (>45 ka; N.Z. ¹⁴C, No. 65) and regional geology (Te Punga, 1962; Fleming, 1972). As described, it is difficult to identify the transition of marine to terrestrial sediments; however, a sponge spicule bearing unit (‘d’) is considered marine-limiting point at 22.6 ± 2 m amsl (Figure 4 ID 413).

The Tokomaru Marine Terrace has been described as extending north from Levin to Palmerston North and identified within the intervening Manawatu Valley (Cowie, 1961; Hesp and Shepherd, 1978). Hesp and Shepherd (1978), described the terrace sediments as partly marine and mapped the terrace as cutting into the Rapanui Formation. The terrace was correlated with MIS 5 on the basis that the “upper beds” had been tentatively correlated with the Oturian (MIS 5) Stage by Fleming (1971). Although Hesp and Shepherd (1978) make no mention of the Otaki Formation, it is near certain that the correlation referred to is that of the Otaki Formation section described by Te Punga (1962) and interpreted by Fleming (1971; 1972), who does not relate the Otaki Formation to the Tokomaru Terrace.

The correlation of the Tokomaru Terrace in the Manawatu Valley with MIS 5 prompted Palmer et al (1988) to identify a marine terrace surface at Otaki as the Tokomaru Terrace underlain by Otaki Formation. The terrace surface is described as being near continuous from Otaki through to Whanganui allowing confident correlation of the Tokomaru Terrace and the Rapanui Terrace. The identification of the terraces as equivalents and the correlation of the Rapanui Terrace with MIS 5e (Pillans, 1983; 1990b) implies the mapping by Hesp and Shepherd (1978) of the Tokomaru and Rapanui Terraces within the Manawatu Valley is incorrect. Although the shore platform is not visible, Palmer et al. (1988) identify both Otaki beach sand



and Otaki dune sand to the north and south of Otaki, inclusive of the terrace riser and sediments described by Te Punga (1962). A cross-section of the inner terrace riser, exposed by the Otaki River, provides the location for the fossil marine cliff, which would have been cut by peak sea level during MIS 5e; however, the shoreline angle is obscured by Otaki dune sand. Height constraint for the Otaki beach sand is not provided other than described as ~30 m amsl for the area and therefore, the location of the fossil marine cliff is identified as direct sea-level indicator but assigned a large uncertainty, 30 ± 5 m amsl (Figure WID 571).



Detailed stratigraphic and lithological analysis of the Otaki Formation accompanied extensive mapping in the region between Otaki and Tokomaru (Sewell, 1991). Sewell (1991) retained use of the Tokomaru Marine Terrace designation for the wave-cut platform underlying the sediments of the Otaki Formation and the fossil marine cliff cut at the peak of MIS 5e sea level. The extensive mapping confirmed the presence of two later terraces, designated Post Tokomaru Marine Terrace (PTMT) 1 and PTMT 2, correlated with MIS 5c and MIS 5a respectively (Palmer et al., 1988; Sewell, 1991). The age of PTMT 1 and PTMT 2 was determined from their relationship with the Tokomaru Marine Terrace (an extension of the MIS 5e Rapanui Terrace), the similarity of the terrace succession to the Rapanui, Inaha, and Hauriri Terrace succession to northwest, and comparative loess stratigraphy overlying PTMT 1 – it is overlain by four loess units similar to other marine terraces correlated with MIS 5c on south North Island. Sewell (1991) referred to the marine and dune sediments of PTMT 1 and PTMT 2 as Otaki Formation. Although the detailed stratigraphic and lithological descriptions allowed for a comprehensive paleoenvironmental interpretation of the region, the described sections are not assigned a height in relation to modern sea level and RSL indicators cannot be derived.

4.2 South Island

The South Island is comprised of seven government regions, listed in a clockwise order from the northernmost: Nelson, Tasman, Marlborough, Canterbury, Otago, Southland, and West Coast Regions (Figure 2). The majority of RSL indicators have been identified in the Marlborough and Canterbury Regions (Figure 4), which will be reviewed last. No RSL indicators have been correlated to MIS 5 in the Nelson or Tasman Region. Speleothems within a cave correlated with a 60 m amsl terrace within the Tasman region were dated to MIS 5e; however, it was concluded that the cave (and terrace) predated the Last Interglacial (Williams, 1988).



Terrace sequences of the South Island are less continuous and more fragmented than the North Island and attempts at broad correlation of terraces within South Island and New Zealand are fewer. Identification of paleo-shorelines and features on the west and south coast of the South Island is made difficult by proximity to the Alpine Fault (driving coastal deformation) and by dense vegetation (Wellman and Wilson, 1964; Suggate, 1992). These regions also remain remote and difficult to access. A large portion of the Canterbury coast is subject to subsidence and the northeast coastline is positioned within the Marlborough Fault System (Rattenbury et al., 2006).

4.2.1 West Coast and Southland

Extensive mapping of the northern West Coast Region in the mid-20th Century identified numerous marine terraces and associated formations (Suggate, 1965; 1992). In one of the earliest studies Suggate (1965) described terrace sequences for three separate segments of the coastline (Hokitika to Greymouth, Greymouth to Fox River, and Charleston to Westport) and designated MIS 5 marine sediments the Awatuna Formation. The best-preserved terrace sequence of the northern West Coast is found between Charleston and Westport, where four marine terrace surfaces are identified; two of which Suggate (1965) correlated to MIS 5. Nathan (1975) mapped six distinct terraces north of Charleston, correlating the marine sands and gravels underlying two of the terraces with the Early and Late Oturi Interglacial (MIS 5): the Virgin Flat Formation (6 m to 15 m amsl) and Waites Formation (also 6 m to 15 m amsl), respectively. In later reassessment of the northern West Coast, Suggate (1992)




Figure 4: Position, elevation, and age correlation of the South Island RSL indicators within the WALIS database and described here. Each RSL indicator is identified with their unique WALIS database identifiers (e.g. WID 34) and is correlated in text with their original publication. Symbols of indicators that would stack extend offshore for clarity. Marine- and terrestrial limiting points are indicated with arrows. The sea-level datum of WID 51 and WID 768 (Otago Region) are mean low tide and an higher high water, respectively. Locations and landscape features mentioned in text are also shown with built-up areas (e.g. towns and cities) indicated by stars. Hill shade data sourced from the LINZ Data Service and licensed for reuse under CC BY 4.0 (LINZ, 2020b).

estimated the age of the terraces from each of the three coastal segments previously described using the stratigraphic relationship of the paleo-shorelines to glacial deposits, previously published radiocarbon ages (the majority of which provide minimum ages), and individually calculated uplift rates for each sequence (with consideration for differential uplift along the coast) to correlate terraces with paleo-sea levels derived from the Huon Peninsula sea-level curve (Chappell and Shackleton, 1986). The Awatuna Formation was subdivided and correlated with MIS 5e and MIS 5c; the younger deposits retaining the Awatuna designation and the older MIS 5e deposits renamed Rutherglen Formation (Table 1; Suggate, 1985; 1992). The Virgin Flat and Waites Formations (Nathan, 1975) were correlated with the Rutherglen Formation (Suggate, 1985); however, insufficient information is present in either publication for an RSL indicator north of Charleston. The type sections for the Awatuna Formation and the Rutherglen Formation are identified as direct indicators (Figure 1; WIDs 824 and 825; Suggate, 1965; 1985; 1992), both within 20 km proximity to Greymouth. The formations consist of rhyolite cemented marine sands and gravels located near to former marine cliffs. The surface elevations of the Awatuna and Rutherglen Formations are 52 m and 67 m amsl, respectively, with surface elevation decreasing to the south. Preusser et al (2005) applied multiple luminescence dating techniques to polymineral fine-grain and K-rich feldspar sediment samples taken from within the Awatuna Formation



section and correlated the formation with MIS 5. Due to concerns with method expressed by Preusser et al (2005) and discussed below (Section 5.3.4), both the Awatuna and Rutherglen Formations are correlated in WALIS with MIS 5, *sensu lato*.

530 To the south, between the Moeraki and Haast Rivers, Nathan and Moar (1975) identified and briefly described three terraces: Sardine-2, Sardine-1, and Knights Point. The Sardine-1 Terrace is interpreted as fluvial and its surface is found at an intervening height (56.4–59.4 m amsl) to the marine Sardine-2 (24.4–32 m) and Knights Point (140–147 m) Terraces. The results of radiocarbon analysis (mostly of wood) were considered ambiguous with a mixture of ages implying contamination by either older or younger carbon; therefore, age was determined primarily by correlation with the glacial-interglacial
 535 sequences described in the north by Suggate (1965). Sardine-2 was correlated broadly to MIS 5, Sardine-1 to MIS 7 or possibly MIS 6 due to the harsh climatic conditions indicated by pollen within the sediments, and Knights Point Terrace was considered to pre-date MIS 7. The Sardine-2 Terrace forms an ~400 m-wide strip on the north side of Ship Creek and dips gently to the south. The marine sands are described as ilmenite-rich, typical of beach and near-shore sand found along the modern coast to the north, and are considered a direct RSL indicator (Figure ID 50); however, elevation constraint is poor and the
 540 uncertainty on the actual elevation of these sediments is large, 27.4 ± 7.87 m amsl.

Due to the rate of vegetative growth, the sections described by Nathan and Moar (1975) were quickly overgrown, but reanalysis of the Knights Point Terrace was made possible by road-widening early in the new century (Cooper and Kostro, 2006). Cooper and Kostro (2006) provided a detailed stratigraphic description and interpretation with IRSL age constraint. Seven lithofacies indicate a transgressive system within a high-energy coastal environment, possibly inclusive of storm event deposits. In section
 545 drawings, the shore platform varies between 104.2 m and 105.5 m amsl. The surface elevation of the uppermost facies, Facies 7, reaches a maximum height of 113 m amsl. Facies 7 is interpreted to have been deposited within the breaker or swash zone of a coastal beach and is considered a direct RSL indicator (Figure ID 52). The lower elevation of Facies 7 is not provided and is defined within WALIS as existing between 113 and 110 m amsl because the unit is drawn as exceeding 3 m thickness in the publication stratigraphic sections (Figure 3, Cooper and Kostro, 2006). Two OSL samples (KP-01-TL and KP-03-TL)
 550 were collected, both from Facies 7. The resulting age from KP-01-TL of 123 ± 7 ka was accepted. KP-03-TL had an older age (146 ± 8.4 ka) but the sample was reported to be in radioactive disequilibrium and was considered likely to be an overestimate of age. However, based on the SAR measurement procedure used at the time, it is possible that these ages suffer from ‘anomalous fading’ (Wintle, 1973) and should be viewed as minimum ages only. Although, correlation to other marine terraces in New Zealand is discussed, the ramifications of the MIS 5e age of Knights Point Terrace for the ages of the Sardine-1 and
 555 Sardine-2 Terraces is not (Cooper and Kostro, 2006). Given the uncertainty of the Knights Point ages, the original interpretation by Nathan and Moar (1975) with MIS 5 at ~ 24 m to 32 m amsl, retains validity.

Bull and Cooper (1986) claimed to have identified the remnants of numerous marine terraces in the Southern Alps on either side of the Alpine Fault, inland of the West Coast locations described immediately above. Terrace age was determined by correlation to the Huon Peninsula terrace sequence in Papua New Guinea (Chappell, 1974). Terraces correlated with MIS 5
 560 were identified by morphology: notched spur ridges associated with sea cliffs and shore platform remnants overlain by well-rounded quartz pebbles and cobbles interpreted to have been formed within a beach environment. The average altitude of these terraces is between 686 m and 899 m amsl. This work was refuted by Ward (1988b) on the basis of terrace morphology, the origin of the quartz pebbles, and the terrace altitudes. Ward (1988b) argued terrace morphology is not preserved as described, the pebbles could be moa gizzard stones, and cites concerns regarding correlation to the distant Papua New Guinea sequence
 565 and the implications of the inferred uplift rates, suggesting that any semi-regular sequence of terrace altitudes could be correlated with the Huon Peninsula – points refuted by Bull and Cooper (1988). Pillans (1990a) also considered the evidence for a marine origin of the terraces ambiguous and demonstrated, not only that the concerns of Ward (1988b) regarding the altitude correlation with Huon Peninsula were justified, but also that the terraces are as likely to be ridge crest notches



controlled by drainage density, slope, and uplift rate. It is the opinion of these authors that the argument remains unresolved and further research would be necessary to conclude whether the features are marine terraces.

Fiordland, forming the southwest corner of Southland and the South Island, is extremely rugged and largely inaccessible. Although numerous marine terraces have been identified, they have only been marginally described and studied (Wellman and Wilson, 1964; Bishop, 1985; Ward, 1988a, Kim and Sutherland, 2004). Wellman and Wilson (1964) identified a marine bench at 500 ft [152 m] cut across a penultimate glaciation moraine within Hollyford Valley and continuing to the coast where it forms a notch in the headland on the south side of Martins Bay. Bishop (1985) identified eleven levels of marine “surfaces”, inclusive of the present-day intertidal reef to the highest at ~1000 m amsl, between Preservation Inlet and Knife and Steel Harbour from which he inferred uplift rates. The terrace sediments are poorly exposed and not described. The age of the Pleistocene terraces was determined by comparing the terrace sequence morphology to that of the terrace sequence at Taranaki on the North Island (Pillans, 1983). The terrace surface correlated with 120 ka, designated h_6 , is extensive with an average altitude of 370 m amsl at the strandline. Terraces h_5 (300 m) and h_4 (210 m) are correlated with MIS 5c and 5a respectively. There are localized outcrops of rounded pebbles and cobbles and fossil islands and stacks rise above the terrace surface south of Preservation Inlet. However, no precise localities or descriptions are provided in the publication and a suitable RSL indicator was not identified.

Ward (1988a), working between the outlet of Big River and Te Waewae Bay, slightly overlapping the study area of Bishop, described a sequence of at least 13 marine terraces reaching up to 1000 m altitude. Finding a lack of suitable material for any geochronological method, terraces were matched to late Quaternary oxygen isotope stratigraphy of deep-sea cores using a “simple uplift model”. The terraces are described in general terms: the lower terraces, inclusive of the last interglacial, consist of loess overlying several meters (up to 20 m locally) of marine gravel and sand on bedrock. Terraces 2, 3, and 4 are correlated to MIS 5a, 5c, and 5e respectively. The height of the inner margin of each terrace above mean sea level was estimated from a combination of 1:63,360 topographic maps (NZMS1 S173 & 174), oblique air and ground photographs, and limited ground truthing with an altimeter: Terrace 2 at 60 m, Terrace 3 at 90 m, Terrace 4 at 140 m. These heights are distinctly different from the height of the terraces correlated to MIS 5 (*sensu lato*) by Bishop (1985) for the adjacent coastline to the west (Table 1). Ward (1988b) correlates the 370 m terrace in his study to MIS 9. Although the regions described by Bishop (1985) and Ward (1988a) are bisected by the Hauroko Fault, both authors recognize that there is no observable displacement of the terrace surfaces due to the fault and the difference in terrace heights is due to different interpretations of terrace age within each study.

Kim and Sutherland (2004) applied two methods of cosmogenic nuclide surface exposure dating (^{10}Be and ^{26}Al) to date shore platform erosion surfaces in bedrock located to the west of the Bishop (1985) and Ward (1988a) studies, between Newton River and Dusky Sound. Although three marine limiting points (Figure 53, 54, 55) were derived from this study, they are not of high quality for a number of reasons. Sample locations are briefly described as “uplifted beach region” and “uplifted sea stacks and reefs”, providing little context. Strandline elevations are generalized, determined from air photographs, a map of undefined scale, and field interpretations. The terrace strandline correlated with MIS 5e is described as 65 m amsl even though sample elevations are reported to have been collected from 51 m to 72 m amsl. The terrace ages cluster between 102.1 ± 4.6 ka to 118 ± 5.9 ka. The ages derived from ^{10}Be and ^{26}Al are consistent with each other, but of the three samples documented in WALIS (because the sample location and elevation could be constrained), an ^{26}Al age was derived from only one (WC02_15). The designation by the authors of the 65 m terrace to MIS 5e provides support to the terrace ages to the east derived by Ward (1988b) although, in the east the terrace described at 60 m is assigned to MIS 5a. The cluster of ages reported by Kim and Sutherland (2004) are better correlated to MIS 5c with a peak sea level at ~109 ka, which would result in better agreement between the studies. However, later work (Putnam et al., 2010; Kaplan et al., 2011) to calibrate *in situ* cosmogenic ^{10}Be production rates indicate lower production in the Southern Hemisphere and cosmogenic exposure ages published prior to 2010 are likely to be too young, probably by at least 12% (Williams et al., 2015). Due to these discrepancies, the terrace is



assigned to MIS 5 (*sensu lato*) within WALIS. Kim and Sutherland dated an additional surface between 92 m and 136 m amsl, which is at similar elevation to Terraces 3 (90 m) and 4 (140 m) correlated by Ward (1988a) with MIS 5c and 5e respectively, but all results were considered anomalously low and no age determination for that surface was made.

4.2.2 Otago Region

615 The Otago Region is the only region within New Zealand, other than Northland, that has historically been considered
 tectonically stable or to have had tectonic movement (Gage, 1953; Gibb, 1986; Pillans, 1990a, Beavan and Litchfield,
 2012); however, relatively few studies of shorelines have been completed there since the early review of Gage (1953). Ongley
 (1939) mapped an intermittently preserved 50 ft to 60 ft [15.2-18.3 m] terrace surface and underlying shore platform at 20 ft
 [6.1 m] northwards from the Clutha [then Molyneux] River mouth to Brighton. Cotton (1957), although referencing Ongley
 620 (1939), refers to this terrace as the 40 ft [12.2 m] terrace and discusses the possibility that it and the underlying shore platform
 are of Monastirian age; i.e. Last Interglacial. No specific locations are provided for the terrace, but it is described to extend
 from the Clutha River north to Brighton and north from Shag River into Canterbury.

Later work at Taieri Beach, in the region south of Brighton, was completed by Rees-Jones et al. (2000) and Litchfield and Lian
 (2004). The coastline south of Taieri Mouth (Waipori River) is on the upthrown (east) side of the reverse Akatore Fault. Rees-
 625 Jones et al. (2000) revisited a raised beach deposit identified by Bishop (1994) as 'h₂' and sampled sediments for IRSL
 determination. The terrace surface is described as occurring at 8-11 m above "high sea level" with cover beds consisting of
 two sand units upon a wave-cut platform capped by loess. The lower sand unit, ~1 m thick, is a coarse, well-sorted, reddish-
 orange sand and gravel, and is considered a direct RSL indicator of a beach deposit at 5.4 ± 2.2 m above "high sea level"
 (Figure 47). The upper sand, ~1.5 m thick, is fine and lightly weathered yellow sands of aeolian origin. The overlying
 630 loess varies between 1 to 4 m in thickness. Samples were taken from all three units and were dated using IRSL. The IRSL
 signal measured from K-feldspar inclusions inside of quartz sand grains coming from the beach sand (sample NZ10) returned
 an age of 71 ± 14 ka. The deposition of this unit, based upon this age estimate, was correlated with MIS 5a. Litchfield and
 Lian (2004), citing concerns for the luminescence technique used, resampled the beach sand (5.5 ± 2.5 m amsl Figure 48) and
 analyzed it using both thermoluminescence (TL) of quartz (Sample W2857) and IRSL of polymineral silicates
 635 (Sample TBE1). The resulting ages of 117 ± 13 ka (TL) and 117 ± 12 ka (IRSL) prompted reassignment of the beach sand to
 MIS 5e. However, the uncertainties provided are at 1-sigma deviation and the ages are consistent with any time over a span of
 ~50 ka. Furthermore, anomalous fading was observed in the IRSL sample (TBE1) indicating a potential underestimated age,
 but because age was consistent with the TL sample (W2857), the fading was considered insignificant and no correction was
 made. There are valid concerns regarding the luminescence methods at the time both studies were completed (Section 5.3.4)
 640 and the ages must be considered with skepticism.

The movement along the Akatore Fault, which runs roughly sub-parallel to the coast for ~22.5 km before moving offshore
 near the mouth of the Tokamairiro River to the south, had been recognized in the earlier work of Cotton (1957). He cited the
 mapping of the 40 ft surface (Ongley, 1939) to argue that the movement appeared to be localized to the coastal regions
 immediately surrounding the fault (not fully recognized at the time) as they are the only locations where the terrace shows any
 645 offset in height. Trenching across the fault revealed at least three reverse fault ruptures since c. 13,300 yr BP, with single event
 displacements of 1.6 m to 2.5 m (Taylor-Silva et al., 2020). Ground penetrating radar profiles show that the displacement of
 the shore platform is ~3 m and the marine terrace has likely only been displaced by the relatively recent events, following a
 minimum 110 ka period of quiescence on the fault. However, this does imply that the RSL indicator at Taieri Beach, has been
 displaced by potentially up to 3 m. The distinction in age of the beach sand is important for calculating uplift rates and
 650 implications for stability along the Otago coastline (Litchfield and Lian, 2004). If the beach sand (5.5 ± 2.5 m amsl) is
 correlated to MIS 5e, then relatively little uplift along the coastline since the Last Interglacial until reactivation of the Akatore



Fault is inferred; however, if deposited during MIS 5a, then the elevation of the beach sand would indicate significant uplift has occurred.

Litchfield and Lian (2004) also analyzed a marine terrace location at Warrington, within Blueskin Bay north of Dunedin. The terrace surface reaches 4–8 m amsl and is slightly deformed by a tilt to the northeast and a displacement of 1 m by a small, unnamed northwest-striking fault. Although the terrace is located adjacent to the Waitati Fault, that fault is not considered responsible for any deformation as it displays no evidence of Quaternary activity. The marine terrace sediments, exposed in a modern marine cliff, consist of Quaternary volcanic boulder beach deposits overlain by quartzofeldspathic beach sand of undefined thicknesses. The beach sands are overlain by loess and the sequence variably overlies an older loess or shore platform cut into Miocene volcanics. The contact between the beach boulders and beach sands at 2.5 ± 1.5 m amsl is considered a direct RSL indicator of beach deposition (Figure 49). One meter of uncertainty was added to the measurement because it is unclear whether the sample site is on the upthrown or downthrown side of the unnamed fault. Samples for IRSL dating of K-feldspar were taken from the beach sand (WBE2) and the underlying loess (WBE1), providing ages of 97 ± 11 ka and 96 ± 5 ka, respectively. Although inverse, the ages are practically indistinguishable and both samples were considered to have suffered from anomalous fading and to be minimum ages. Litchfield and Lian (2004) argue for MIS 5e deposition of the beach sand due to the likelihood that the underlying loess was deposited during MIS 6.

Kennedy et al. (2007) surveyed the coast north of Warrington to Oamaru and described the height of the last interglacial shoreline as varying between 5 and 8 m above low water level. It is unclear how accurate the measured height reflects paleo sea level as the surface is described as a mixture of deposits consisting of eroded platforms, gravel beaches, and estuarine mud and beach sand and a detailed stratigraphic description is only provided for a section at Shag Point. On the southeastern end of Shag Point, imbricated boulders and marine sands are deposited upon a shore platform surface cut into carbonaceous mudstone. The boulders (≤ 2.5 m long *a*-axis, 0.2 m to 1.2 m thick *c*-axis) are only found on the seaward edge of the terrace. Directly landward of the boulders is the well-sorted and laminated marine sand, which continues landward and is visible on both sides of the point. The boulders and sand are overlain by two distinct loess units of ~ 2 m total thickness. The distance from the inner margin is not stated and an accurate thickness of the marine sand is not provided; therefore, the shore platform at 7.4 ± 0.25 m above mean low water level is considered marine limiting (Figure 51). The sand directly behind the boulders was dated by IRSL analysis of the Na-feldspar component (WLL181), providing an age of 81.9 ± 11.7 ka. The overlying loess deposits were also analyzed and indicate two younger phases of deposition, 78.6 ± 4.2 ka and 28.9 ± 4.4 ka. The apparent MIS 5a age of the marine sands is used to constrain the age of the boulders and argue for deposition by a tsunami wave not exceeding 3 m. However, given the likely location of sea level during MIS 5a (-10.5 ± 5.5 m; Creveling et al., 2017) and concerns for the optical dating of Na-feldspar (Section 5.2.3), this interpretation must be regarded with skepticism.

Sections of the MIS 5 marine Hillsgrove Formation have been described on the southern coastline of Cape Wanbrow, located immediately south of Oamaru (Grant-Mackie and Scarlett, 1973). The two sections consist of nearshore marine sediments, likely deposited within a few meters depth of water, as indicated by the marine and terrestrial mollusc and avian fossil assemblages. The section at the north end of the South Oamaru Beach is described for height above the base of the cliff backing the modern beach; constraint above modern sea level is not provided and a relative sea level indicator cannot be derived. The section to the east is constrained to height above the present shore platform with the inner margin designated '0 m'. The provided elevations are considered as heights above mean higher high water (MHHW) because the inner margins of modern shore platforms are typically identified at that height in relation to mean sea level (Rovere et al., 2016). The sediments are deposited upon a shore platform at 4 m above MHHW, formed on Upper Eocene volcanics and around an erosion-resistant stack of Upper Eocene limestone. A basal gravel (0.3 m to 1.0 m thick) grades to a consolidated weathered yellow sand and clay with some pebbles (1.5 m to 2.0 m thick) containing avifauna fossils of large moa, which grades upwards through two gravel bands and into overlying loess and paleosols. The good preservation of the fossil fauna, the height of the deposits above



sea level, and a minimum radiocarbon age (undescribed) are used to correlate the marine sediments to the Last Interglacial
 (sensu lato; Grant-Mackie and Scarlett, 1973). Only the basal gravel is considered marine following the later description of
 the overlying sands as dunes (Worth and Grant-Mackie, 2003) and is designated a direct RSL indicator (Figure ID 768)
 at 4.5 ± 1.1 m above MHHW.

4.2.3 Canterbury and Marlborough Regions

The Canterbury Plains is a coastal plain stretching 160 km from Timaru to the Waipara River. The plain is formed of a series
 of coalescing alluvial fans, emerging from the Southern Alps and exceeding 50 km in width, that has been subsiding under the
 weight of glacial advances and outwash. The last interglacial RSL record of the southern Canterbury Region is dominated by
 stratigraphic correlation of the Bromley Formation beneath the Canterbury Plains. Brown et al. (1988) correlated the subsurface
 strata described in water well cores on the basis of gravel aquifers and intervening fine-sediment strata. The type section for
 the last interglacial (sensu lato) Bromley Formation is derived from two wells within the Bromley suburb (M35/1875 and
 M35/1926; Figure IDs 74 and 75). The lithology consists of beach, lagoonal, dune, and coastal swamp sediments
 associated with rising, high, and declining sea level. The broad definition of the Bromley Formation has limited its use to a
 terrestrial-limiting point. Twenty-one indicators, with upper and lower elevations, were identified (Figure IDs 65 to 85).
 The maximum depth of the Bromley Formation is 79 m below mean sea level (bmsl) and the minimum depth is 5 m bmsl.
 Heights were also constrained for WALIS entry by use of the Canterbury Regional Council well database
 (www.ecan.govt.nz/data/well-search; Environment Canterbury, 2020).

The well cores investigated by Brown et al. (1988) are predominantly clustered within Christchurch. Outside of the city, the
 Bromley Formation was identified at 33 m to 56 m bmsl within a well (M36/1251, Figure 4, WID 72) on the shore of Lake
 Ellesmere, south of Banks Peninsula (Brown et al., 1988). An additional core from the nearby Gebbies Valley was analyzed
 by Shulmeister et al (1999). This 75 m long core retains a 200-ka record of marine transgressions and regressions. Sediments
 attributed to MIS 5e are not identified; however, aqueous deposits within Units G through K are correlated to MIS 5a and MIS
 5c (Figure IDs 103 to 105). Units G and H record a transgressive succession and marine embayment, of which Lake
 Ellesmere is considered a modern equivalent. Unit H (34.10 m to 24.67 m bmsl; WID 103) is marine-limiting. Unit I is
 correlated to a relative sea-level fall, probably during MIS 5b, before the next sea-level transgression represented by Units J
 and K. Unit J is interpreted as a beach or storm deposit, at or close to (within 5 m) of sea level, and is considered a direct
 indicator of RSL (22.74 m to 22.15 m bmsl; WID 104). Diatoms and phytoliths within Unit K (22.15 m to 14.12 m bmsl; WID
 105) indicate a transition from a marine embayment to a lake or lagoon environment in probably interglacial conditions. The
 unit is considered a terrestrial-limiting point of sea level. The nine TL results for samples from 12.52 m to 43.75 m bmsl
 concentrate around 90 ka to 130 ka. Age estimates from throughout the core were further refined with radiocarbon ages and
 paleoenvironmental data to calibrate the sedimentary units to the oxygen isotope record (Martinson et al., 1987). Lagoon
 sediments (Unit C) were identified at greater depth (61.50 to 48.25 m bmsl) and correlated to MIS 7. Although in relatively
 close proximity (<5.5 km) to the Lake Ellesmere core reported by Brown et al. (1988), comparison of the core interpretations
 is difficult without age constraint or detailed stratigraphic description from the latter.

North of the Canterbury Plains and into the Marlborough Region, last interglacial marine terraces are uplifted by the North
 Canterbury fold-and-thrust belt, with terraces most prominent on the limbs of actively growing anticlines (Oakley et al., 2017).
 This coast was subject to extensive description by both Jobberns (1928) and Suggate (1965). One distinct terrace and multiple
 smaller, higher and more dissected terraces, occur at varying elevations intermittently along the coast. The marine terrace
 surfaces are in many locations obscured by later, glacial-period gravel fans. Additional studies (Powers, 1962; Fleming and
 Suggate, 1964) indicate locations of possible LIG marine terraces within the northern Canterbury Region. Studies providing
 the necessary stratigraphic descriptions, age and elevation constraint to identify RSL indicators have been concentrated in five



735 coastal segments: the Waipara River to Motunau Beach, Haumuri Bluffs, Kaikoura Peninsula, Clarence River to Woodbank Stream, and Long Point to Boo Boo Stream. The latter coastal section is within the Marlborough Region.

North of the Waipara River, terrace remnants are continuous along a vertically-steep coast for a distance of ~6 km. The terraces are then absent for ~10 km until the coastline broadens at Motunau Beach (Jobberns, 1928; Jobberns and King, 1933; Suggate, 1965). Yousif (1989), using remote sensing, mapped the terrace surfaces following the nomenclature and age correlation of
 740 Jobberns (1928) and Carr (1970); although, these designations were often inclusive of multiple terrace surfaces at different altitudes within one interglacial. The most extensive terrace immediately north of the Waipara River is the Tiromoana Terrace (Carr, 1970); at Glenafric the surface tilts upwards to the northeast from ~45 m to ~80 m amsl (Oakley et al., 2017). At Motunau Beach the most extensive terrace has been alternatively identified as the Motunau Coastal Plain or Motunau Terrace and extends offshore to include Motunau Island (Jobberns, 1928; Jobberns and King, 1933; Suggate, 1965; Carr, 1970; Oakley et al., 2017). Jobberns and King (1933) provided early detailed lithology for the marine sediments of the Motunau Beach Terrace, including identification of marine and estuarine fossil mollusc found *in situ*. Representative stratigraphy drawn by Oakley et al. (2017) show both Tiromoana and Motunau Terraces to consist of a shore platform overlain in upwards succession by fossiliferous marine sand and gravel, alluvium, and loess. Oakley et al. (2017) sampled both terraces in multiple locations for IRSL of loess and marine sediments and AAR analyses of fossil mollusc shell. Precise sample locations are not provided, but
 750 are shown on a publication map. In general elevations are not provided in Oakley et al. (2017), but multiple elevations for the Tiromoana Terrace inner margin are estimated from shore platform elevations measured in transect in a later publication (Oakley et al., 2018); however, exact locations for the inner margin elevations are not provided. The marine-limiting points for Tiromoana (58.7 ± 2 m amsl; Figure D 108) and Motunau Beach (70 ± 4 m amsl; Figure D 109) marine sediments were derived from AAR sample (GA4 and MB5, respectively) locations (Oakley et al. 2017).

755 The Tiromoana Terrace is correlated with MIS 5c indicating that higher terrace remnants in close proximity are of MIS 5e and MIS 7 age. Oakley et al. (2018) correlate the Bob's Flat Terrace, a less extensive terrace located to the southwest and northeast of the Tiromoana Terrace, with MIS 5e. However, this age interpretation appears to be reliant on the higher elevation of the terrace as no AAR or IRSL samples or data for this terrace are provided in Oakley et al. (2017). The IRSL and AAR results suggest at least two phases of deposition at Motunau Beach, a possibility first proposed by Suggate (1965). The distribution in
 760 ages indicates partial reoccupation, or incision with deposition, of the seaward edge of an earlier terrace during the MIS 3 highstand. The earlier, more extensive portion of the terrace, is correlated to MIS 5a, with the likelihood that the more minor terrace remnants at higher elevations are of MIS 5e and MIS 7 age. Oakley et al. (2017) argue that wave erosion explains the minimal record older terraces in the area as well as the lack of a MIS 5c terrace above the Motunau Beach Terrace and MIS 5a terrace below the Tiromoana Terrace. The apparent higher elevation of the younger MIS 5a Motunau Beach Terrace to the
 765 older MIS 5c Tiromoana Terrace is not discussed by Oakley et al. (2018), but is likely due to the variable uplift rates along the coastline.

Further north the Haumuri Bluffs (alternatively spelled "Amuri") have been subject to many studies (Jobberns, 1928; Fleming and Suggate, 1964; Suggate, 1965; Ota et al. 1984; 1996; Oakley et al. 2017; 2018). Ota et al. (1984) has provided the most extensive description of the area. Four marine terraces are identified in the sequence; from highest elevation to lowest they
 770 are: Tarapuhi Terrace, Kamps Hill Upper and Lower Terraces, and Amuri Bluff Terrace. The terraces each consist of a shore platform overlain by marine sands, localized alluvial fan deposits, and loess. The cover beds of the Amuri Bluff Terrace include extensive fluvial deposits. The Kamps Hill Terraces are not present at Haumuri Bluffs but appear in the sequence approximately 2 km to the south. The dip of the terrace surfaces indicates northward tilting.

The Tarapuhi Terrace retains a diverse molluscan fossil assemblage and collections were described by both Fleming and
 775 Suggate (1964) and Ota et al. (1996). The fossil molluscs were used by Ota et al. (1996) for AAR analyses in the first attempt



to provide numerical age to the sequence and returned a broad age of 135 ± 35 ka. The cold-water environment indicated by the fossil assemblage was used to constrain deposition of the Tarapuhi Terrace to MIS 5c. The age of the Kemps Hill Upper, Kemps Hill Lower, and Amuri Bluff Terraces were determined by best fit to the Huon Peninsula sea-level curve (Chappell and Shackleton, 1986) constraining formation within MIS 5 and MIS 4. Additional AAR analysis of the fossil mollusc assemblages within the Tarapuhi and Amuri Bluff Terraces, as well as IRSL analysis of the surrounding marine sediments, strengthened correlation of the Tarapuhi Terrace to MIS 5c and indicated formation of the Amuri Bluff Terrace within MIS 5a (Oakley et al., 2017). Formation of the two Kemps Hill terraces is attributed to a double peak in sea level at the beginning of MIS 5a (90.6 ± 2.0 ka and 84.0 ± 1.6 ka) based upon the sea level curves derived from coral terraces (Lambeck and Chappell, 2001) and a $\delta^{18}\text{O}$ curve (Siddall et al., 2007; Oakley et al., 2018). RSL indicators are derived from the Tarapuhi Terrace inner margin (173.1 ± 2 m amsl; Figure 4 ID 107; Oakley et al., 2017; 2018) and an overlying fossiliferous marine unit (163 ± 2.2 m amsl; Figure 4 ID 102; Ota et al., 1996) and from the elevation of the Amuri Bluff Terrace inner margin (40.7 ± 5 m amsl; Figure 4 ID 100; Oakley et al., 2017; 2018) and the contact between the shore platform and overlying marine gravels (32 ± 3 m amsl and 50 ± 5 m amsl; Figure 4 IDs 63 and 64; Ota et al., 1984). The Kemps Hill terraces have not been subject to direct geochronological analysis and indicators for these terraces are not identified. Ota et al. (1984) includes over a dozen stratigraphic sections for the terraces; however, specific measurements above sea level are not provided and description in text can differ from the drawn section (e.g. thickness of the deposit and height above sea level) introducing uncertainty. Furthermore, the surface that underlies the terraces undulates by tens of meters. Only two marine-limiting points (WIDs 63 and 64) for the Amuri Bluff Terrace are included from this publication.

Approximately 20 km to the northeast of Haumuri Bluff, the Kaikoura Peninsula forms a prominent headland. Suggate (1965) identified four distinct surfaces, of which the third highest and most extensive surface (170 ft to 200 ft [~ 52 m to 61 m] altitude), could be traced on the mainland to the west. Ota et al. (1996) identified five marine terraces, in addition to Holocene surfaces, deformed by numerous folds and faults, which in general cut perpendicular across the peninsula, resulting in a down-tilt to the northwest, parallel to the long axis of the peninsula. The terraces are labelled in decreasing elevation and age as I, II, III, IV, and V. Terrace III is the extensive terrace identified by Suggate (1965). Representative sections of each terrace were drawn from auger cores and AAR analyses of fossil mollusc from Terrace I provided geochronological constraint. A total of seven direct RSL indicators are derived from Terrace I, II, and III, where rounded pebbles and gravel are interpreted as beach deposits, providing direct indicators except for one marine-limiting point from Terrace III at 58.8 ± 2 m amsl (Figure 4 ID 101). The beach deposits for each terrace were measured: for Terrace I at 89.5 ± 2 m amsl (Figure 4 ID 100); for Terrace II at 61.3 ± 2.0 m amsl, 71.5 ± 2.1 m amsl, and 73 ± 2.1 m amsl (WIDs 97, 98, 99); and for Terrace III at 40 ± 2.1 m amsl and 51 ± 2.1 m amsl (WIDs 95 and 96). Ota et al. (1996) analyzed the same mollusc species for AAR from Terrace I as from the Tarapuhi Terrace at Haumuri Bluff, *Tawera spissa*, allowing for direct comparison of data. The age derived from Terrace I of 110 ± 20 ka and similar dominance of cool-water species within the fossil assemblage, prompted the correlation of Terrace I with MIS 5c and the Tarapuhi Terrace (Table 1). Terrace II was correlated with sea-level fall into MIS 5b and Terrace III, the most expansive terrace, with MIS 5a. The remaining Terraces IV and V were considered younger than MIS 5. The age estimate for the remaining terraces were determined by assuming constant rates of uplift and correlation with the Haumuri Bluff Terraces; however, the complexity of the tectonics in the intervening distance recommends against comparing the two locations (Duffy, 2020).

The two remaining marine terrace sequences on the northeast South Island coast are located between Clarence River and Woodbank Stream in north Canterbury, and Long Point and Boo Boo Stream in the Marlborough Region (Ota et al., 1996). The main marine terrace within these coastal sections is designated “MM”. Below MM, at lower elevation is a later Holocene terrace. Minor remnants of an upper terrace, “UM”, above MM, are present at both locations, but marine sediments were not identified within the Marlborough section and a marine origin for the UM Terrace there cannot be confirmed. Between the



Clarence River and Woodbank Stream, the MM Terrace is covered by thick (often >10 m) slope-wash deposits. The elevation of the underlying shore platform decreases from 143 m amsl in the south to 105 m amsl in the north along a distance of 5 km, indicating significant down-tilting to the north. Detailed stratigraphic descriptions were derived from five exposures of the terrace sediments (Ota et al., 1996), allowing for the identification of multiple direct sea level indicators at elevations, from south to north, of 144 ± 2.2 m amsl, 117 ± 3.6 m amsl, 112 ± 2.2 m amsl, 105 ± 2 m amsl, 107 ± 2.8 m amsl (Figure 94 to 90). Beach deposits are identified in each exposure, inclusive of bored boulders abutting a bedrock cliff at Location 10, indicating exact shoreline position. Four of the five exposures show the beach deposits directly overlying the shore platform, but at Location 7 (WID 91), the beach deposits are found overlying estuarine silt, indicating that the marine terrace sediments record a transgressive rise in sea level.

Between Long Point and Boo Boo Stream, the surface of the MM terrace is well defined with the best preservation to the north and increasing dissection by streams to the south (Ota et al., 1996). The elevation of the inner margin is measured between 55 and 80 m amsl, with the highest elevations at either end of the terrace. The marine sediments and terrestrial cover beds of the MM Terrace are exposed along the former sea cliff between it and the younger Holocene terrace and four direct RSL indicators were identified at elevations, from south to north, of 72.5 ± 2.5 m amsl, 45.5 ± 2.5 m amsl, 42.5 ± 3.2 m amsl, and 51.5 ± 3.2 m amsl (Figure 89 to 86). The well-stratified beach deposits, composed of well-rounded pebbles within a coarse, sandy matrix, are typically about 3 m thick overlying the shore platform. The surface elevation of the beach sediments varies between 45 m to 74 m amsl. The non-marine cover beds vary in thickness between 4 m and 11 m and are inclusive of multiple paleosols within loess. At the most representative section, Location 3 (WID 88), 11 m of non-marine cover beds includes at least 4 loess units with three paleosols.

Ota et al. (1996) correlated the MM Terrace from both regions with MIS 5e based upon the size of the terrace, the well-defined beach sediments, and position above earlier Holocene terraces. To the south, between the Clarence River and Woodbank Stream, a fluvial terrace merges with the MM Terrace. A TL age from loess overlying the fluvial terrace provides minimum age constraint in support of the MIS 5e designation. Ota et al. (1996) also argue that the number of loess layers (typically 3 to 4) overlying the MM terrace is consistent with loess stratigraphy overlying other last interglacial deposits in the northern South Island and southern North Island. The UM Terrace is correlated to MIS 7 due to its higher elevation.

4.3 Summary of New Zealand RSL indicators

This work identified and, using the WALIS database, standardized 77 unique RSL indicators (direct, marine- or terrestrial-limiting points) along the coastline of the North and South Islands of New Zealand (<http://doi.org/10.5281/zenodo.4056376> Ryan et al., 2020a). The slim majority (26) are direct indicators of sea level, with the remaining almost evenly split between terrestrial-limiting (23) and marine-limiting (24) points. Direct indicators of sea level were identified from elevation measurements of beach deposits or marine terrace shoreline angles. All but one of the terrestrial-limiting points are derived from the below surface Bromley Formation record described by Brown et al. (1988) from well cores within the Canterbury Region. Marine-limiting points are derived from shore platform elevations or poorly described marine sediments from which proximity to the coastline could not be determined. The majority (41) of RSL indicators are correlated broadly with MIS 5 (*sensu lato*). The remainder are correlated with the interglacial peak MIS 5e (12), warm interstadials MIS 5c (6) and MIS 5a (11), and indicators from one marine terrace on the Kaikoura Peninsula were correlated with sea-level fall from MIS 5a into MIS 5b (3). The most common methods for age determination are terrace correlation, luminescence methods, and amino acid racemization.

The North Island contains eighteen RSL indicators, of which the majority are marine-limiting points (12) correlating broadly with MIS 5 (10). The only indicator not likely to have been displaced by VLM is the One Tree Point direct sea-level indicator in the Northland Region with most other indicators identified in regions of well-recognized tectonic deformation associated



with the Hikurangi Margin forearc and Whanganui Basin (Figure 2). The Otamaroa and Te Papa Terraces on the Raukumara Peninsula are most likely MIS 5e and MIS 5a in age, in contrast to the luminescence results within Wilson et al (2007). We agree with the reasoning of Wilson et al. (2007) that although the peninsula is uplifting, it is unlikely that the MIS 5a and MIS 3 record would be preserved and the MIS 5e would not. The IRSL luminescence method at the time was not fully developed and the results most likely represent minimum ages (Section 5.3.4). On the west coast, the Shelly Beach Formation, younger undifferentiated Hawera sediments, and correlated Waiau A and B Formations and northern Taranaki marine terraces need reassessment. These formations and terraces would likely be sources of multiple RSL indicators, but have not been described in detail since Chappell (1975) and have not been analyzed by any absolute geochronological method. Reassessment of the age of these features is important because the use of terrace correlation by Chappell (1975) to determine age introduces uncertainty in the chronology of the region.

Of the twelve of the fifty-nine South Island RSL indicators are located outside of the Canterbury-Marlborough Regions. Except for those located within the Otago Region, all indicators have been identified in locations subject to significant coastal deformation. Of the four Otago locations, at least two (Taieri Beach and Blueskin Bay) appear to have been subjected to tectonic displacement since deposition. Either the section of coastline within Otago is not as stable as previously presumed, or care must be taken to ensure future field locations are not located adjacent to minor faults. The indicators from the West Coast (4) and Southland (3) regions are broadly correlated with MIS 5; although most of the West Coast and Southland indicators have been analyzed by either luminescence or stable isotope methods, there are concerns for the validity of the results (Section 5.3.4). Similarly, other than the recent reassessment by Oakley et al. (2017), age constraint for sediments and marine terraces in the Canterbury and Marlborough Regions is poor, relying predominantly upon stratigraphic succession or terrace correlations. The sharp transition from the subsiding landscape of the Canterbury Plains to an uplifting one in the north is distinctly marked by the elevation of RSL indicators. The minimal preservation of the MIS 5e terrace and variable preservation of MIS 5c and MIS 5a in the coastal sections of the northern Canterbury Region is attributed to wave erosion and fluvial dissection by Oakley et al. (2017).

In summary, in agreement with Gage (1953) and Pillans (1990a), the New Zealand record of last interglacial sea level lacks quality description, measurement, and age constraint even amongst those records produced since 1990. The following section will provide in greater detail the sources of uncertainty. However, the literature indicates robust relative sea-level indicators are present, not only for MIS 5e, but also the following MIS 5c and 5a interstadials. Better constraint on these records would assist in the understanding of sea-level fluctuation throughout MIS 5 (*sensu lato*). Furthermore, there is potential for the development of a regionally-specific New Zealand sea-level record for MIS 5e from the Northland and Otago Regions; although care must be taken within the latter region to ensure no influence from minor tectonic movement.

5 Sources of uncertainty

This section provides a summary of the issues contributing the greatest amount of uncertainty to the New Zealand marine terrace records: inconsistent terrace terminology and nomenclature, coastal deformation, and methods of geochronological constraint.

5.1 Terrace terminology and nomenclature

The disparate application of terminology and nomenclature used when discussing marine terraces in New Zealand contributed to uncertainty in interpreting the meaning of potential sea-level indicators within our database. The example used by Pillans (1990a, pg. 221) in his argument for the use of standardized terminology for marine terraces was the prevalence for authors to refer to a terrace by altitude, e.g. the “100 metre terrace”. The questions being: which feature of the terrace is at 100 metre and at what position within the terrace? The work of numerous authors over the decades has also contributed to an inconsistency



in terrace nomenclature. Perhaps the most glaring example is found in the Horowhenua lowlands where the Tokomaru Marine
 Terrace designation refers to the shore platform, the inner margin fossil marine cliff, and the terrace surface but the cover beds
 are identified as the Otaki Formation (Section 4.1.6).

Uncertainty is also introduced through age correlation (in absence of a numerical age) to either distant marine terrace sequences
 or local stage names. The early 20th Century practice of correlating marine terrace sequences to Mediterranean stages
 (Monastirian, Tyrrhenian, Milazzian, Sicilian), which at the time also lacked numerical age and have since been redefined,
 precludes the correlation of any potential RSL indicator to MIS 5 with certainty. The youngest formally accepted Pleistocene
 stage within New Zealand is the Haweran Stage, which encompasses the past 0.340 Ma (Raine et al., 2015). The stage boundary
 is identified at the base of the Rangitawa Tephra - the oldest bed of the Hamilton Ash Formation (Section 5.3.2). Local stage
 names for interglacial/glacial periods within the Pleistocene have been proposed (e.g. Suggate, 1965; 1985), but these were
 often developed from local sediment sequences without numerical age constraint and have not been formally adopted.

Within our database we have endeavored to adapt any inconsistent marine terrace terminology to that recommended by Pillans
 (1990a; Figure 1) and provide clarity for terrace nomenclature. To avoid confusion and to assist in understanding the global
 context of the New Zealand sea-level record, we have chosen not to use local New Zealand stage names for the Pleistocene
 and have given preference for marine oxygen-isotope stages (MIS) designations, an internationally recognized scale.

5.2 Coastal deformation and GIA

The position of the New Zealand archipelago straddling the active boundary of the Australian and Pacific plates has produced
 a coastline subject to variable rates of vertical land movement (VLM) due to complex tectonics and displacement associated
 with earthquakes. Although displacement by an earthquake can have dramatic effect, inter-⁹⁰⁵ seismic deformation (deformation
 between earthquake events) is more likely to influence long-term trends in VLM (Beavan and Litchfield, 2012). Estimates of
 long-term VLM in New Zealand have been consistently derived from marine terraces or below-surface marine deposits (e.g.
 Chappell, 1975; Pillans, 1983; Bishop, 1985; Suggate, 1992; Berryman, 1993; Begg et al., 2004; Wilson et al., 2007; Oakley
 et al., 2018).

While the marine terrace sequences have proven useful for estimating long-term (beyond Holocene) rates of uplift or
 subsidence (e.g. Beavan and Litchfield, 2012), the significance of Holocene ice-volume change (outside and within New
 Zealand) on glacial and hydro-isostatic adjustment (GIA) signals on coastal deformation and relative sea level have not been
 quantified for New Zealand (King et al., 2020). GIA encompasses all deformational, as well as gravitational, and rotational-
 induced changes to relative sea level in response to the buildup and retreat of ice sheets with residual and variable affect along
 coastal sections depending upon their proximity to former glaciers, ice caps and sheets (Arctic and Antarctic, Simms et al.,
 2016). In other words, the magnitude and wavelength of the solid Earth response to ice-and water-load history varies with time
 (in relation to glacial maxima) and geographical location producing a gradient in relative sea level that is modulated by mantle
 rheology. As a result, sea-level change around New Zealand is not uniform. For example, the Holocene highstand peaked in
 the North Island at ~2.65 m amsl between 8.1 to 7.2 cal ka BP, whereas in the South Island, the highstand peaked at no more
 than ~2 m amsl between 7.0 to 6.4 cal ka BP (Clement et al., 2016).

GIA models for the late Pliocene (Grant et al., 2019) suggest that interglacial sea-level high-stands in New Zealand might
 deviate significantly from the global mean (eustatic) when the largest meltwater contributors are from the northern hemisphere
 (larger distance) and/or when a significant imbalance between West and East Antarctic Ice Sheets (WAIS and EAIS,
 respectively) melting exists. In particular, a larger contribution from EAIS would shift the eustatic band northward (above the
 northern island), causing a lower-than-eustatic local sea-level high-stand. On the other hand, a larger contribution from WAIS
 would result in a larger-than-eustatic high-stand in New Zealand. A uniform melting of WAIS and EAIS would result in a



RSL change that is very close to the eustatic, thus making New Zealand an excellent location for the constraining Antarctic Ice Sheet volume variations. Accordingly, New Zealand is an interesting location, both in terms of fingerprinting and ice-volume constraints; although it should be combined with other relevant locations such as South America (most sensitive to WAIS), southern Africa, and the northern hemisphere. Finally, the lack of consideration for GIA in the assessment of VLM rates in New Zealand is concerning. Neglecting GIA on active coastlines has been shown to lead to overestimated uplift rates at an average of 40%, but also up to 72% (Simms et al., 2016).

5.3 Geochronological constraint

Marine terrace ages have been constrained in New Zealand using multiple different approaches. Correlated-age determination of a terrace at a known elevation, either with distant terrace sequences or a marine oxygen-isotope sea-level curve, is the most consistently applied method to determine age. Three additional age constraint methods are core to New Zealand marine terrace chronology: tephrochronology for terraces proximal to volcanic centers in the North Island; amino acid racemization (AAR); and luminescence techniques. The limitations of these methods and the consequent impacts on age constraint and terrace correlation are briefly discussed here.

5.3.1 Marine terrace correlation

Muhs (2000) summarized three common global correlation methods that have been used in New Zealand to determine marine terrace age (and subsequently uplift rates): 1) assumption of a constant uplift rate on a shore-normal terrace sequence; 2) relation diagrams for shore-parallel terrace sequences; and 3) unique altitudinal spacing of terraces assuming a constant uplift rate. Methods 2 and 3 were developed within New Zealand, the former was used by Pillans (1983) to assist with age determination of the Whanganui Basin sequence, and the latter by Bull (1985) to determine the age of marine terraces along the Alpine Fault. Major assumptions of each of these methods are that the uplift rate has been constant over time and/or that there has been no gradient of uplift normal to the coastline; although shore-parallel variation in uplift rate is expected in method 2 (Muhs, 2000). All of these methods also suffer from the circular problem of assigning a glacio-eustatic sea level to a terrace to estimate its uplift rate, which is then used to determine age of additional terraces.

Global correlation methods largely rely upon a chosen distant terrace sequence (e.g. Huon Peninsula) for paleo sea level and/or a marine oxygen-isotope curve to assist in determining age (Table 2). However, these records are unlikely to accurately reflect the timing and height of sea-level highstands that formed New Zealand marine terraces. Distant terrace sequences are expected to have a local peak sea level differing in both height and timing due to their own GIA signal and other localized processes (e.g. steric effects, Creveling et al., 2015). Marine oxygen-isotope records represent a convoluted signal of ocean mass and sea temperature variation resulting in a paleo global mean sea-level estimate with considerable uncertainty (Rovere et al., 2016) and can also include their own tectonic uplift correction (Simms et al., 2016).

The importance of chosen sea level record was recently stressed by Duffy (2020) when determining slip rates on the Kaikoura Peninsula of New Zealand. Concerned with the use of a LIG sea-level highstand estimate from a region without the same GIA characteristics, Duffy (2020) expressed preference for an MIS 5e highstand estimate of $+2.1 \pm 0.5$ m that was derived from RSL indicators identified on the tectonically stable, far-field Gawler Craton in South Australia (Murray-Wallace, 2002; Murray-Wallace et al., 2016). This record was preferred because southern Australia and the South Island lie within 5° of latitude, a comparable distance from the Antarctic ice cap peripheral bulge, and because Australia and New Zealand display similar records of the mid-Holocene highstand and subsequent sea-level fall (Sloss et al., 2007; Clement et al., 2016; Duffy, 2020). However, the Australian record can be expected to differ from that of New Zealand because the North Island extends into much lower latitudes and due to South Island ice volume changes (e.g. Golledge et al., 2012; James et al., 2019; Carrivick et al., 2020); the role of Holocene glacial mass on GIA requires further investigation (King et al., 2020). The difference in the



Table 2: Numerical estimates of the height and timing of MIS 5 sea-level highstands used in age correlations of RSL indicators and/or to generate uplift rates

RSL Indicator Primary Reference	Last Interglacial		
	Height (m)	Age (ka)	Reference(s)
MIS 5			
Te Punga 1962; Fleming, 1972	-	120-80	-
Mildenhall, 1995	-	125-70	Pillans, 1991
MIS 5e			
Chappell, 1970; 1975	5 ± 3	120	Chappell, 1974; Bloom et al., 1974
Pillans, 1983; 1990b	5	120	Chappell & Veeh, 1978
Palmer, 1988	-	c. 120	Pillans, 1985
Berryman, 1993	6 ± 5	124 ± 5	Chappell & Shackleton, 1986
Kim and Sutherland, 2004	3 ± 2	120	Lambeck & Chappell, 2001
Litchfield and Lian, 2004	-	128-113	Chappell et al., 1996
Cooper and Kostro, 2006	5 ± 2	c. 125	Veeh & Chappell, 1970; Harmon et al., 1983; Chappell et al., 1996; Stirling et al., 1996
Wilson et al 2007	0 ± 5	125 ± 5	Pillans et al., 1998
Oakley et al, 2017; 2018	5 ± 2	124.5 ± 5.5	Lambeck & Chappell, 2001; Siddall et al., 2007
MIS 5c			
Chappell, 1970; 1975	-14	104	Chappell, 1974; Bloom et al., 1974
Berryman, 1993	-19 ± 5	106 ± 5	Chappell & Shackleton, 1986
Ota et al., 1996	-9 ± 3	100 ± 5	Chappell & Shackleton, 1996
Wilson et al 2007	-28 ± 5	105 ± 5	Pillans et al., 1998
Oakley et al, 2017; 2018	-23 ± 11	106.9 ± 3.0	Lambeck & Chappell, 2001; Siddall et al., 2007
MIS 5b			
Ota et al., 1996	-24 ± 5	96 ± 5	Chappell & Shackleton, 1996
MIS 5a			
Chappell, 1970; 1975	-15	83	Chappell, 1974; Bloom et al., 1974
Berryman, 1993	-19 ± 5	81 ± 5	Chappell & Shackleton, 1986
Ota et al., 1996	-19 ± 5	81 ± 5	Chappell & Shackleton, 1996
Rees-Jones et al 2000	-20 to 0	c. 80	Pillans, 1983; Chappell & Shackleton, 1996; Ludwig et al., 1996
Wilson et al 2007	-24 ± 5	80 ± 5	Pillans et al., 1998
Oakley et al, 2017; 2018	-50 ± 10	71.3 ± 2.0	Lambeck & Chappell, 2001; Siddall et al., 2007

980 timing and amplitude of the Holocene highstand between South and North Island (Clements et al., 2016) makes it is reasonable to expect a similar difference during the Last Interglacial; although, given the current resolution of available geochronological methods, it is unlikely that difference can be discerned within the LIG or deeper time.

An additional concern is the correlation of paleo-coastlines to the correct MIS. As pointed out by Litchfield and Lian (2004), the correlation of the Taieri Beach to either MIS 5e or MIS 5a has significant implications for derived uplift rates. Furthermore, 985 estimates of the height and timing for each sea-level highstand within MIS 5 (*sensu lato*) has changed, particularly for the interstadials, significantly through time (Table 2).

5.3.2 Tephrochronology

On the North Island, two tephtras have been used consistently as stratigraphic markers to constrain and identify MIS 5 terraces and associated sediments: the Rotoehu Ash and the Hamilton Ash (Pullar and Grant-Mackie, 1972; Chappell, 1975; Pain, 990 1976; Iso et al., 1982; Ota et al., 1989; Berryman, 1993; Wilson et al., 2007; Claessens et al., 2009). The Rotoehu Ash, the basal member of the Rotoiti Tephra Formation, erupted from the Taupo Volcanic Zone and serves as an important regional stratigraphic marker in the North Island. Numerous age estimates, ranging from c. 61 ka to 45 ka, have been published (see comprehensive summary by Flude and Storey, 2016), but the most recently published age estimates for the Rotoehu Ash (derived from a combination of ¹⁴C-accelerator mass spectrometry, (U-Th)/He, and ⁴⁰Ar/³⁹Ar geochronological methods) place 995 deposition at 47.5 ± 2.1 ka (Daniššik et al., 2012; Flude and Storey, 2016). The age of the Rotoehu Ash, although well-constrained, does only provide minimum age constraint to Pleistocene marine terraces.



The Hamilton Ash Formation has been described as a sequence of time transgressive ash beds (H1 to H7) (Ward, 1967; Vucetich and Pullar, 1982; Iso et al., 1982; Lowe et al., 2001). Recently (Lowe, 2019), it has been proposed that the sequence is more accurately described as a, “composite set of clayey, welded paleosols very probably developed by upbuilding pedogenesis from MIS 10 to 5” (p. 23 of 24). Alternatively, Briggs et al. (2006) found the designation as a Formation inappropriate given that some of the tephra beds could be found preserved as individual layers in the Maketu area – near to the locus of Mid-Pleistocene TVZ activity, the likely source of the tephra. Where both are present, the Hamilton Ash sequence is overlain by the Rotoehu Ash (Ward, 1967; Vucetich and Pullar, 1982; Iso et al., 1982; Lowe et al., 2001). The Hamilton Ash has been used to constrain marine terrace development or associated dune sand deposition with the understanding that the youngest tephra bed dated to ~MIS 6/5 (Pullar and Grant-Mackie, 1972; Chappell, 1975; Pain, 1976; Iso et al., 1982; Ota et al., 1989). Only the basal ash bed, the Rangitawa Tephra (H1), has been directly dated using radiometric methods to c. 400 ka to 340 ka (Kohn et al., 1992; Pillans et al., 1996; Lowe et al., 2001). Age constraint for the overlying ashes is based upon the physical characteristics, clay mineral assemblages, and climatostratigraphic associations of the deposits and their paleosols (Lowe, 2019; Lowe et al., 2001). The uppermost Hamilton Ash, the Tikotiko Ash (H6/H7), was likely deposited within MIS 6 or MIS 5e and subject to pedogenic alteration throughout MIS 5. A minimum age for the surface has been estimated at c. 74 ka; however, the only other chronostratigraphic constraint for the H2 through H7 ash beds is that they are older than the Rotoehu Ash (Lowe et al., 2001; Lowe, 2019). The possible 74 ka age for the H6/H7 beds does have implication for marine terrace chronology, where the beds have been used to argue a minimum MIS 5e age. Until the age of the H6/H7 bed is better constrained, it is not recommended for use as a chronostratigraphic marker to discern MIS 5e age.

It should be noted that loess cover bed stratigraphy has also been used to assign relative age to marine terrace sequences in New Zealand, e.g. the Mahia Peninsula (Berryman, 1993). However, loess stratigraphy, as described in the publications reviewed, is generally localized and lacking in regional (or greater) spatial correlations. Furthermore, without numerical age constraint to the sedimentation history of the location, the resolution of such an approach is questionable (Muhs, 2000).

5.3.3 Amino Acid Racemization (AAR)

Amino acid racemization has been applied to both wood and marine mollusc shell to produce numerical ages for marine terraces in New Zealand (<http://doi.org/10.5281/zenodo.4056376> Ryan et al., 2020a). The first numerical age constraint was derived from alloisoleucine-isoleucine D/L values of fossil wood fragments overlying marine terraces of the Whanganui Basin terrace sequence analyzed on a modified Technicon amino acid auto-analyser. The fossil wood D/L values were calibrated with the Rangitawa Tephra, which overlies numerous mid-Pleistocene marine terraces, the youngest of which is the Ararata Terrace (Pillans, 1983; Pillans, 1990b; Pillans and Kohn, 1981; Pillans et al., 1996). The age of the tephra, at c. 370 ka, was determined by fission-track dating of both zircon and glass components, and used to constrain the age of the Ararata Terrace to c. 400 ka, MIS 11. Calibration of the fossil wood retrieved from the Ararata Terrace lignite cover bed, closely underlying the Rangitawa Tephra, allowed numerical ages to be derived from the D/L values of fossil wood samples from the younger Rapanui Terrace using the integrated rate equation.

The minimum age derived for the Rapanui Terrace was 110 ka and the terrace was correlated with the MIS 5e transgression cycle, culminating at c. 120 ka with peak sea level between 5 and 8 m aspl (Chappell and Veeh, 1978; Pillans, 1983). An uplift model, developed using the Ararata and Rapanui Terraces as anchors, was used to calculate the ages for the other marine terraces, which were correlated with every interglacial (odd MIS) from MIS 17 to MIS 3. The only marine isotope stage to have multiple sea-level peaks represented is MIS 5, the Rapanui Terrace (MIS 5e, 120 ka), the Inaha Terrace (MIS 5c, 100 ka), and the Hauriri Terrace (MIS 5a, 80 ka) (Section 4.1.2). This geochronological framework has not only served as a basis for determining regional uplift rates of the marginal Whanganui Basin in the mid-late Quaternary, but also underpins many of the marine terrace age correlations within New Zealand since (Bishop, 1985; Ward, 1988a; Bussell, 1990; Pillans, 1990a; Ota



et al., 1996). Ota et al. (1996) used unpublished AAR alloisoleucine-isoleucine D/L values of *Tawera* shell from the Hauriri Terrace (80 ka) to calculate numerical ages from alloisoleucine-isoleucine D/L values (determined using an automated amino acid analyzer) of *Tawera spissa* (*T. spissa*) shell collected on the Kaikoura Peninsula and Haumuri Bluffs (Section 4.2.3). The numerical ages were calculated using the integrated rate equation and the Arrhenius equation to allow for the difference in long-term temperature history between the regions.

The aminostratigraphy of the Whanganui sequence was redefined by Bowen et al. (1998) using shell of the marine bivalves *T. spissa* and *Austrovenus stutchburyi*, analyzed by ion-exchange high-performance liquid chromatography (HPLC) to provide alloisoleucine-isoleucine D/L values. The D/L values were constrained by the previous fission-track ages and also biostratigraphy, magnetostratigraphy, and correlated with marine $\delta^{18}\text{O}$ isotope stages (Beu and Edwards, 1984; Beu et al., 1987; Shackleton et al., 1990; Bassinot et al., 1994; Pillans et al., 1994). The reassessment by Bowen et al. (1998), although not resulting in any change in MIS correlation, provided more direct constraint for the marine terraces as the previous analysis by Pillans (1983) was of wood fragments of unknown genera in lignite beds overlying marine sediments. Furthermore, there are methodological concerns regarding the application of AAR to wood that have never been resolved; namely the kinetics of racemization within wood and the presence of internal sugars (e.g. arabinose), which interact with amino acids to form melanoidin polymers (Zumberge et al., 1980; Blunt et al., 1987; Rutter and Vlahos, 1988). Additional causes for concern are wood degradation and the seemingly increased sensitivity of wood to internal and external environmental factors.

The aminostratigraphy developed by Bowen et al (1998) does not appear to have as broad application to determining marine terrace age as the former framework developed by Pillans (1983). However, recently, Oakley et al. (2017) collected additional shell samples from sites used by Bowen et al. (1998), as well as other sites of known age (MISs 1, 5a, 5e, 7, 11, 17), to provide multiple calibration points for AAR analyses of *T. spissa* shell from marine terraces in the North Canterbury region. The different pretreatment methods and machine types used in each study (ion-exchange HPLC by Bowen et al (1998) vs. reverse-phase HPLC by Oakley et al. (2017)) to determine D/L values, and preference for different amino acids in developing chronologies, meant that results from the studies were not directly comparable. Oakley et al. (2017) also utilized a new approach of Bayesian statistical methods (Allen et al., 2013) to determine the best fit AAR age equation and calculate uncertainties. The best fitting function for relating sample age to D/L value was determined to be the simple power-law kinetics where an exponent is applied to either the D/L value or t , time (Goodfriend et al., 1995), with aspartic acid D/L values used for calibration (Oakley et al., 2017). These results were used in combination with infrared stimulated luminescence to develop a new chronology for North Canterbury marine terrace development, resulting in significantly different chronologies in some locations than earlier studies where elevation or degree of fluvial dissection of the marine terrace had been used to assess age (e.g. Carr, 1970; Yousif, 1987).

AAR has also been applied to fossil shell of the warm-water estuarine bivalve *Anadara trapezia* found in multiple locations around North Island associated with interglacials ranging in age from MIS 11 through MIS 5e (Beu and Maxwell, 1990; Murray-Wallace et al., 2000). Currently extinct in New Zealand, almost all fossil *A. trapezia* correlated with MIS 5e and MIS 7 were restricted to the northeast, with exceptions at Gisborne and Mahia Peninsula dating to MIS 5e, and southern Hawke's Bay to MIS 7. The only known samples correlated with MIS 11 (on the basis of stratigraphic evidence) are found within the Whanganui Basin and are the only *A. trapezia* identified in the southwest. The geographic distribution of fossil *A. trapezia*, increasingly restricted to the northeast with successive interglacials, probably reflects the poor fossil record of estuarine fauna, and decreasing extent of warm-waters during interglacials as a consequence of geographic changes to the New Zealand landmass from progressive uplift (Murray-Wallace et al., 2000).

In summary, although early age constraint for the Whanganui Basin marine terraces was derived from AAR analysis of wood fragments (Pillans, 1983), an unreliable method, AAR of mollusc shell has proven useful in building an aminostratigraphic



framework in New Zealand and for discerning shell of different interglacial age; e.g. MIS 5 vs MIS 7 (Bowen et al., 1988; Ota et al., 1996; Murray-Wallace et al., 2000; Oakley et al., 2017). It has been shown to be complimentary to luminescence methods, providing more certainty where results converge, with the ability to assist in resolving discrepant results in the latter method (Oakley et al., 2017). The relatively new (Allen et al., 2013) Bayesian method to determine numerical age from D/L values has been proven successful when applied to a New Zealand Pleistocene dataset (Oakley et al., 2017). The applicability of AAR to mollusc and foraminifer tests for resolving relative age at Pleistocene timescales, identifying reworked contributions to deposits, and developing useful aminostratigraphic frameworks, as well as its complimentary nature to luminescence techniques, has been proven in numerous locations globally (Hearty et al., 1992; 2004; Murray-Wallace and Belperio, 1994; Murray-Wallace et al., 2010; Wehmiller et al., 1995; Wehmiller, 2013; Kaufman et al., 2013; Ryan et al., 2020b). These capabilities of AAR, and its relatively low cost (in comparison to other geochronological methods), recommend it for continued and future use resolving geochronology in New Zealand.

5.3.4 Luminescence

Both thermoluminescence (TL) and infrared stimulated luminescence (IRSL) have been used in the dating of New Zealand last interglacial sediments; predominantly of multi-grain aliquots of fine polymineral silts (4-11 μm), and rarely, coarser sand-size grains of quartz and feldspars (<http://doi.org/10.5281/zenodo.4056376> Ryan et al., 2020a). The application of luminescence dating to New Zealand quartz grains has been found to be unsuitable because the quartz grains generally have a very dim optically stimulated luminescence (OSL) signal intensity, are adversely affected by thermal transfer of charge between single-aliquot regenerative-dose (SAR) measurement cycles, and display unpredictable sensitivity changes across these cycles (Preusser et al., 2005; 2006). These characteristics are, at least in part, attributable to the young sedimentary history of the quartz grains, but may also be a characteristic of the geological provenance of these grains influencing their intrinsic brightness (Preusser et al., 2006). Even if these hurdles could be overcome, the relatively high dose rates in New Zealand sediments (e.g. in comparison to Australia) would more than likely lead to the saturation of the quartz OSL signal well within 100 ka (Roberts et al., 2015), meaning OSL analysis of quartz sediments dating to the last interglacial would more than likely result in minimum ages only. These numerous issues effectively remove quartz OSL from the chronological toolkit for dating Last Interglacial sediments in New Zealand.

Greater success in dating LIG sediments may be gained using K-feldspar grains; although these too have impediments. The primary concern is ‘anomalous fading’ (Wintle, 1973), which is the leak of electrons from unstable traps in the grains over the period of burial resulting in underestimated ages. Significant developments in the procedures used for measuring K-feldspar grains allow for this fading to be either reduced to negligible levels or accounted for (Thiel et al., 2011; Buylaert et al., 2012; Rui et al., 2019). However, the stability of the non-fading component of the IRSL signal results in a much slower IRSL decay rate (Buylaert et al., 2012; Li et al., 2013; Smedley et al., 2015). Thus, a longer period of sunlight exposure is required to fully reset the signal and/or a residual dose measurement must be made to account for this prior to age determination. Alternatively, the measurement of single-grains of K-feldspar (rather than multi-grain polyminerals) would enable the adequacy of the previous resetting to be tested, which the use of multi-grain aliquots cannot as reliably disentangle (Jacobs and Roberts, 2007). Finally, unlike quartz, the onset of saturation of K-feldspars occurs at much higher doses enabling the accurate dating of LIG and older sediments (Aitken, 1998; Huntley and Lamonthé, 2001; Li et al., 2014).

The lack of critical information regarding how the luminescence signal/s was measured in the studies reviewed here does not instill confidence in the reported ages. Given the methodological advances in luminescence measurement procedures since many of the studies reviewed here were completed, with the exception of the recent work by Oakley et al. (2017), the ages presented above represent, at best, a starting point. This uncertainty in absolute age of the associated marine terrace sequences and correlation with the appropriate sea-level peak within MIS 5 has significant implications for estimates of uplift rates in



1120 some locations (e.g. Taieri Beach, Section 4.2.2). Advances in method, particularly in using the elevated temperature post-
 infrared IRSL (pIRIR), have been shown to be effective in overcoming the fading issues in very young New Zealand coastal
 sediments (Madsen et al., 2011) and indicate the ability to produce ages with a higher degree of accuracy. The use of single
 feldspar grains, instead of polymineral aliquots, has the potential to assist in the identification and removal of grains that are
 ‘poorly behaved’ prior to age estimation, identify whether or not the deposits contain a reworked component, and the extent
 1125 to which the previous resetting of electron traps was completed (Jacobs and Roberts, 2007). The applicability of the single
 grain method depends upon the grains having a sufficiently bright signal for accurate equivalent dose measurement – New
 Zealand quartz is often reported as rather dim, K-feldspars as much brighter. We therefore recommend a complete and
 systematic reassessment of all previously reported ages. Furthermore, in future all luminescence data (e.g. individual dose rate
 components, fading factors, equivalent dose distribution patterns and associated statistics, and age model selection) needs to
 1130 be reported completely. Such transparency will enable future reviewers to make a much more informed judgement about the
 quality of the data.

6 Further details: Pleistocene and Holocene sea-level fluctuations and climatic change

The Whanganui Basin, in addition to the Rapanui (MIS 5e), Inaha (MIS 5c), and Hauriri (MIS 5a) terraces (Section 4.1.2),
 retains a shallow marine basin sequence recording all high sea-level marine oxygen-isotope stages of the last 2.6 Ma of which
 1135 the past 0.7 Ma are represented by marine terraces. This record is important not only for its completeness, but also because it
 offers a detailed paleoenvironmental record from an isolated part of the South Pacific (Pillans, 1991; see Pillans, 2017 and
 references therein) and has been subject to an extensive variety of geochronological and stratigraphical methods. The stratotype
 section and point of the four stages representing Quaternary New Zealand are defined by the fossiliferous marine sediments
 within the Whanganui Basin. A paleovegetation and paleoclimatic record spanning much of the Haweran Stage (0.340 Ma to
 1140 present) has been developed from the terrace cover beds. It also retains a long record of tephra (to c. 2.17 Ma) and loess (to c.
 0.50 Ma) deposition, which provide opportunity for regional correlation in the North Island.

Although the Whanganui Basin sequence is surely the longest and most complete record of sea-level fluctuation and climatic
 change within New Zealand, it is not the only one to extend beyond MIS 5. The vertical land movement along several sections
 of the New Zealand coastline has allowed for the preservation of paleo records of marine, coastal and terrestrial environments
 1145 extending not only to the present but also farther into the Pleistocene; e.g. Mahia Peninsula, North Island (Berryman, 1993)
 and the Firth of Clyde, South Island (Bishop, 1985; Ward, 1988a). The sediments preserved along the New Zealand coastline have
 proven valuable sources of proxy data (commonly in the form of fossil marine mollusc assemblages or fossil pollen) useful
 for biostratigraphy and paleoclimatic reconstructions (e.g. Grant-Mackie and Scarlett, 1973; Dickson et al., 1974; Moar, 1975;
 McGlone et al., 1984; Mildenhall, 1985; 1995; Moar and Mildenhall, 1988; Bussell, 1990; 1992; Berryman, 1992; 1993; Ota
 1150 et al., 1996; Shulmeister et al., 1999; Murray-Wallace et al., 2000). However, lacking the scrutiny of the Whanganui Basin,
 any sea-level record (or associated climatic record) derived from older Pleistocene marine terraces and sediments is likely to
 suffer the similar problems of insufficient description as the MIS 5 record, leading to large uncertainty in age and interpretation.

A recent study (Clement et al., 2016) found spatial and temporal variation in New Zealand’s Holocene relative sea-level change
 may be influenced by a number of different mechanisms. A north-south gradient in RSL may be a result of the position of the
 1155 archipelago within the intermediate field around Antarctica across broad degrees of latitude. Continental levering could have
 a significant effect on the timing and magnitude of sea-level change at a regional to local scale as driven by glacial meltwater
 loading and width of the adjacent continental shelf. Potentially significant drivers of relative sea-level change are regional and
 local effects of tectonic regime, wave climate, and sediment regime – all of which require further research to characterize
 (Clement et al., 2016).



1160 7 Future research directions

The last interglacial sea-level record within New Zealand continues to lack accurate height data, descriptions from which indicative meaning can be derived, and adequate numerical age control. Improvement of this record would have great benefit to both global studies of sea-level and ice-volume change and to New Zealand estimates of long-term vertical land movement and paleoenvironment. A New Zealand regional sea-level curve, derived from Northland and Otago regions, would provide a valuable record within the remote South Pacific, which could assist in understanding the eustatic sea-level response to ice mass change in Antarctica and would allow for better assessment of coastal deformation and improved estimates of long-term vertical land movement around New Zealand. In recognition that most RSL indicators are poorly described, not related to a defined sea-level datum, lack numerical age control, and that most existing luminescence-derived numerical ages were derived from outdated methods, greater accuracy would also require reassessment of nearly all RSL indicators identified to present.

The apparent extensive preservation of MIS 5e, 5c, and 5a marine and coastal sediments and the terrestrial sediments of intervening MIS 5d and MIS 5b substages provides opportunity for study of sea-level fluctuations and climatic changes throughout MIS 5 (*sensu lato*) and changes to long-term rates of vertical land movement.

8 Data Availability

The database is available open access and kept updated as necessary at the following link:
<http://doi.org/10.5281/zenodo.4056376> (Ryan et al., 2020a). The files at this link were exported from the WALIS database interface on 23 September 2020. Description of each data field in the database is contained at this link:
<https://doi.org/10.5281/zenodo.3961543> (Rovere et al., 2020), that is readily accessible and searchable here: <https://walis-help.readthedocs.io/en/latest/>. More information on the World Atlas of Last Interglacial Shorelines can be found here: <https://warmcoasts.eu/world-atlas.html>. Users of our database are encouraged to cite the original sources in addition to our database and this article.

9 Author Contributions

D.D.R. was primary author, responsible for all entries into WALIS, and the conceptualization, development and writing of manuscript, as well as providing expert review of amino acid racemization data. A.C. contributed to the structure and writing of the manuscript, and assisted D.D.R. in developing figures. N.R.J. provided expert review of luminescence data, contributing significantly to that section of the manuscript as well as more minor contributions elsewhere. P.S. contributed significantly to the discussion of coastal deformation and GIA in New Zealand.

10 Competing Interests

The authors declare that they have no conflict of interest.

11 Acknowledgements

The data presented in this publication were compiled in WALIS, a sea-level database interface, developed with funding from the ERC Starting Grant “WARMCOASTS” (ERC-StG-802414), in collaboration with PALSEA (PAGES/INQUA) Working Group. The database structure was designed by A. Rovere, D. Ryan, T. Lorscheid, A. Dutton, P. Chutcharavan, D. Brill, N. Jankowski, D. Mueller, M. Bartz, E. Gowan and K. Cohen.

D.D.R. wishes to thank David Lowe for clarifying aspects of New Zealand tephrochronology and providing useful publications towards that purpose. The authors also thank Alessio Rovere for constructive comments on various drafts of this manuscript.

12 References

- Aitken, M.J., An introduction to optical dating: the dating of Quaternary sediments by the use of photon-stimulated luminescence. Oxford University Press, Oxford. 280 pgs., 1998.
- Allen, A.P., Kosnik, M.A., Kaufman, D.S. Characterizing the dynamics of amino acid racemization using time-dependent reaction kinetics: A Bayesian approach to fitting age-calibration models. *Quaternary Geochronology*, 18, 63-77, 2013.



- Alloway, B., McComb, P., Neall, V., Vucetich, C., Gibb, J., Sherburn, S., Stirling, M. (2005). Stratigraphy, age, and correlation of voluminous debris-avalanche events from an ancestral Egmont Volcano: Implications for coastal plain construction and regional hazard assessment. *Journal of the Royal Society of New Zealand*, 35 (1-2), 229-267.
- 1205 Anderton, P.W. Structure and evolution of the South Wanganui Basin, New Zealand. *New Zealand Journal of Geology and Geophysics*, 24, 39-63, 1981.
- Austermann, J., Mitrovica, J.X., Huybers, P., Rovere, A. Detection of a dynamic topography signal in last interglacial sea-level records. *Science Advances*, 3 (7), e1700457, 2017.
- Ballance, P.F. and Williams, P.W. The geomorphology of Auckland and Northland. In: *Landforms of New Zealand*, second edition, edited by: Soons, J.M. and Selby, M.J., Longman Paul, Auckland, 210-232, 1992.
- 1210 Barlow, N.L.M., McClymont, E.L., Whitehouse, P.L., Stokes, C.R., Jamieson, S.S.R., Woodroffe, S.A., Bentley, M.J., Callard, S.L., Ó Cofaigh, C., Evans, D.J.A., Horrocks, J.R., Lloyd, J.M., Long, A.J., Margold, M., Roberts, D.H., Sanchez-Montes, M.L. Lack of evidence for a substantial sea-level fluctuation within the Last Interglacial. *Nature Geoscience*, 11, 627-634, 2018
- 1215 Bassinot, F.C., Labeyrie, L.D., Vincent, E., Quidelleur, X., Shackleton, N.J., Lancelot, Y. The astronomical theory of climate and the age of the Brunhes-Matuyama magnetic reversal. *Earth and Planetary Science Letters*, 126, 91-108, 1994
- Beavan, R.J. and Litchfield, N.J. Vertical land movement around the New Zealand coastline: implications for sea-level rise. *GNS Science Report*, 2012/29, 41 pgs., 2012
- Begg, J.G. and Johnston, M.R. *Geology of the Wellington area, scale 1:250 000*. Institute of Geological & Nuclear Sciences Geological Map 10. Lower Hutt, New Zealand, Institute of Geological & Nuclear Sciences Ltd., 64 pgs., 2000.
- 1220 Begg, J.G., Van Dissen, R.J., Rhoades, D.A. Subsidence in the Lower Hutt Valley and the interplay between Wellington and Wairarapa Fault earthquakes. 2004 NZSEE Conference, Paper Number 43, 2004.
- Berryman, K.R. Tectonic geomorphology at a plate boundary: a transect across Hawke Bay, New Zealand. *Zeitschrift für Geomorphologie Supplementband* 69, 69-86, 1988.
- 1225 Berryman, K.R. A stratigraphic age of Rotoehu Ash and late Pleistocene climate interpretation based on marine terrace chronology, Mahia Peninsula, North Island, New Zealand. *New Zealand Journal of Geology and Geophysics*, 35 (1), 1-7, 1992
- Berryman, K.R. Distribution, age, and deformation of late Pleistocene marine terraces at Mahia Peninsula, Hikurangi Subduction Margin, New Zealand. *Tectonics*, 12 (6), 1365-1379, 1993
- Beu, A.G. and Edwards, A.R. New Zealand Pleistocene and late Pliocene glacio-eustatic cycles. *Palaeogeography, Palaeoclimatology Palaeoecology*, 46, 119-142, 1984.
- 1230 Beu, A.G. and Maxwell, P.A. Cenozoic Mollusca of New Zealand. *New Zealand Geological Survey Paleontological Bulletin* 58, 518 pgs., 1990.
- Beu, A.G., Edwards, A.R., Pillans, B. A review of New Zealand Pleistocene stratigraphy, with emphasis on the marine rocks. In: *Proceedings of the First International Colloquium on Quaternary Stratigraphy of Asia and Pacific Area*, edited by: Ithara, M. and Kamei, T. (eds). Osaka, Japan, 250-269, 1987.
- 1235 Bishop, D.G. Inferred uplift rates from raised marine surfaces, southern Fiordland, New Zealand. *New Zealand Journal of Geology and Geophysics*, 28 (2), 243-251, 1985.
- Bishop, D.G. *Geology of the Milton area, scale 1:50 000*. Institute of Geological & Nuclear Sciences Geological Map 9. Lower Hutt, New Zealand, Institute of Geological & Nuclear Sciences Ltd, 32 pgs., 1994.
- 1240 Bloom, A.L., Broecker, W.S., Chappell, J.M.A., Matthews, R.K., Mesolella, K.J. Quaternary sea level fluctuations on a tectonic coast: new $^{230}\text{Th}/^{234}\text{U}$ dates from the Huon Peninsula, New Guinea. *Quaternary Research*, 4, 185-205, 1974.
- Blue, B. and Kench, P.S. Multi-decadal shoreline change and beach connectivity in a high-energy sand system. *New Zealand Journal of Marine and Freshwater Research*, 51 (3), 406-426, 2017.



- Blunt, D.J., Easterbrook, D.J., Rutter, N.W. Chronology of Pleistocene sediments in the Puget Lowland, Washington. Washington Division of Geology and Earth Resources, Bulletin 77, 321-353, 1987.
- 1245 Briggs, R.M., Hall, G.J., Harmsworth, G.R., Hollis, A.G., Houghton, B.F., Hughes, G.R., Morgan, M.D., Whitbread-Edwards, A.R. Geology of the Tauranga area, Sheet U14 1:50 000. Occasional Report. Hamilton, Department of Earth Sciences, University of Waikato, 72 pgs., 1996.
- Briggs, R.M., Lowe, D.J., Esler, W.R., Smith, R.T., Henry, M.A.C., Wehrmann, H., Manning, D.A. Geology of the Maketu area, Bay of Plenty, North Island, New Zealand, Sheet V14 1:50 000. Occasional Report No. 26. Hamilton, Department of Earth and Ocean Sciences, University of Waikato, 50 pgs., 2006.
- 1250 Brothers, R.N. The relative Pleistocene chronology of the South Kaipara District, New Zealand. Transactions of the Royal Society of New Zealand, 82, 677-694, 1954.
- Brown, L.J., Wilson, D.D., Moar, N.T., Mildenhall, D.C. Stratigraphy of the late Quaternary deposits of the northern Canterbury Plains, New Zealand. New Zealand Journal of Geology and Geophysics, 31 (3), 305-335, 1998.
- 1255 Bull, W.B. and Cooper, A.F. Uplifted marine terraces along the Alpine Fault. Science Reports, 234, 1225-1228, 1986.
- Bull, W.B. and Cooper, A.F. Response: New Zealand Marine Terraces: Uplift Rates. Science, 240 (4853), 804-805, 1988.
- Bussell, M.R. Palynology of Oxygen Isotope Stage 6 and Substage 5e from the cover beds of a marine terrace, Taranaki, New Zealand. Quaternary Research, 34, 86-100, 1990.
- Bussell, M.R. Late Pleistocene palynology of terrestrial cover beds at the type section of the Rapanui Terrace, Wanganui, New Zealand. Journal of the Royal Society of New Zealand, 22 (2), 77-90, 1992.
- 1260 Buylaert, J.-P., Jain, M., Murray, A.S., Thomsen, K.J., Thiel, C., Sohbati, R. A robust feldspar luminescence dating method for Middle and Late Pleistocene sediments. Boreas, 41, 435-451, 2012.
- Capron, E., Rovere, A., Austermann, J., Axford, Y., Barlow, N.L.M., Carlson, A.E., de Vernal, A., Dutton, A., Kopp, R.E., McManus, J.F., Meniel, L., Otto-Bliesner, B.L., Robinson, A., Shakun, J.D., Tzedakis, P.C., Wolff, E.W. Challenges and research priorities to understand interactions between climate, ice sheets and global mean sea level during past interglacials. Quaternary Science Reviews, 219, 308-311, 2019.
- 1265 Carr, M.J. The stratigraphy and chronology of the Hawera Series marginal marine succession of the North Canterbury Coast. Unpublished PhD Thesis, University of Canterbury, Christchurch, 303 pgs., 1970.
- Carrivick, J.L., James, W.H.M., Grimes, M., Sutherland, J.L., Lorrey, A.M. Ice thickness and volume changes across the Southern Alps, New Zealand, from the little ice age to present. Science Reports, 10, 13392, 2020.
- 1270 Cashman, S.M. and Kelsey, H.M. Forearc uplift and extension, southern Hawke's Bay, New Zealand: Mid-Pleistocene to Present. Tectonics, 9 (1), 24-44, 1990.
- Chapman-Smith, M. and Grant-Mackie, J.A. Geology of the Whangaparaoa area, eastern Bay of Plenty. New Zealand Journal of Geology and Geophysics, 14 (1), 3-38, 1971.
- 1275 Chappell, J. Quaternary geology of the south-west Auckland coastal region. Transactions of the Royal Society of New Zealand, 8 (10), 133-153, 1970.
- Chappell, J. Geology of coral terraces, Huon Peninsula, New Guinea: a study of Quaternary tectonic movements and sea-level changes. Geological Society of America Bulletin, 85 (4), 553-570, 1974.
- 1280 Chappell, J. Upper Quaternary warping and uplift rates in the Bay of Plenty and west coast, North Island, New Zealand. New Zealand Journal of Geology and Geophysics, 18 (1), 129-154, 1975.
- Chappell, J. and Veeh, H.H. Late Quaternary tectonic movements and sea-level changes at Timor and Atauro Island. Geological Society of America Bulletin, 89, 356-368, 1978.
- Chappell, J. and Shackleton, N.J. Oxygen isotopes and sea level. Nature, 324, 137-140, 1986.



- 1285 Claessens, L., Veldkamp, A., ten Broeke, E.M., Vloemans, H. A Quaternary uplift record for the Auckland region, North Island, New Zealand, based on marine and fluvial terraces. *Global and Planetary change*, 68, 383-394, 2009.
- Clark, K., Berryman, K., Litchfield, N., Cochran, U., Little, T. Evaluating the coastal deformation mechanisms of the Raukumara Peninsula, northern Hikurangi subduction margin, New Zealand and insights into forearc uplift processes. *New Zealand Journal of Geology and Geophysics*, 53 (4), 341-358, 2010.
- 1290 Clement, A.J.H., Whitehouse, P.L., Sloss, C.R. An examination of spatial variability in the timing and magnitude of Holocene relative sea-level changes in New Zealand archipelago *Quaternary Science Reviews*, 131, 73-101, 2016.
- Cooper, A.F. and Kostro, F. A tectonically uplifted marine shoreline deposit, Knights Point, Westland, New Zealand. *New Zealand Journal of Geology and Geophysics*, 49 (2), 203-216, 2006.
- Cotton, C.A. Pleistocene shorelines on the compound coast of Otago. *New Zealand Journal of Science and Technology*, B38, 750-762, 1957.
- 1295 Creveling, J.R., Mitrovica, J.X., Hay, C.C., Austermann, J., Kopp, R.E. Revisiting tectonic corrections applied to Pleistocene sea-level highstands. *Quaternary Science Reviews*, 111, 72-80, 2015.
- Creveling, J.R., Mitrovica, J.X., Clark, P.U., Waelbroeck, C., Pico, T. Predicted bounds on peak global mean sea level during marine isotope stages 5a and 5c. *Quaternary Science Reviews*, 163, 193-208, 2017.
- 1300 Danišák, M., Shane, P., Schmitt, A.K., Hogg, A., Santos, G.M., Storm, S., Evans, N.J., Fifield, L.K., Lindsay, J.M. Re-anchoring the late Pleistocene tephrochronology of New Zealand based on concordant radiocarbon ages and combined $^{238}\text{U}/^{230}\text{Th}$ disequilibrium and (U-Th)/He zircon ages. *Earth and Planetary Science Letters*, 240-250, 2012.
- Dickson, M., Fleming, C.A., Grant-Taylor, T.L. Ngarino Terrace: An addition to the late Pleistocene stranded sequence in the Wanganui-Taranaki district. *New Zealand Journal of Geology and Geophysics*, 17 (4), 789-798, 1974.
- 1305 Duffy, B. A geometric model to estimate slip rates from terrace rotation above an offshore, listric thrust fault, Kaikōura, New Zealand. *Tectonophysics*, 786, <https://doi.org/10.1016/j.tecto.2020.228460>, 2020.
- Dutton, A., Carlson, A.E., Long, A.J., Milne, G.A., Clark, P.U., DeConto, R., Horton, B.P., Rahmstorf, S., Raymo, M.E. Sea-level rise due to polar ice-sheet mass loss during past warm periods. *Science*, 349 (6244), aaa4019, 2015.
- Environment Canterbury Regional Council. Well search. <https://www.ecan.govt.nz/data/well-search/> 2020.
- 1310 Fleming, C.A. and Suggate, R.P. The 550ft raised beach at Amuri Bluff. *New Zealand Journal of Geology and Geophysics*, 7 (2), 353-358, 1964.
- Fleming, C.A. 14C Dates in the Quaternary Geology of the “Golden Coast”, Wellington. N.Z. Department of Scientific and Industrial Research, Wellington. Radiocarbon Users Conference, Wellington N.Z. pgs 139-148, 1971.
- Fleming, C.A. The contribution of ^{14}C dates to the Quaternary Geology of the ‘Golden Coast’, Western Wellington. *Tuatara*, 19 (2), 1972.
- 1315 Flude, S. and Storey, M. $^{40}\text{Ar}/^{39}\text{Ar}$ age of the Rotoiti Breccia and Rotoehu Ash, Okataina Volcanic Complex, New Zealand, and identification of heterogeneously distributed excess ^{40}Ar in supercooled crystals. *Quaternary Geochronology*, 33, 13-23, 2016.
- Gage, M. The study of Quaternary strand-lines in New Zealand. *Transactions of the Royal Society of New Zealand*, 81 (1), 27-34, 1953.
- 1320 Ghani, M.A. Late Cenozoic vertical crustal movements in the southern North Island, New Zealand. *New Zealand Journal of Geology and Geophysics*, 21 (1), 117-125, 1978.
- Gibb, J.G. A New Zealand regional Holocene eustatic sea-level curve and its application to determination of vertical tectonic movements. *Royal Society of New Zealand, Bulletin* 24, 377-395, 1986.



- 1325 Golledge, N.R., Mackintosh, A.N., Anderson, B.M., Buckley, K.M., Doughty, A.M., Barrell, D.J.A., Denton, G.H., Vandergoes, M.J., Andersen, B.G., Schaefer, J.M. Last Glacial Maximum climate in New Zealand inferred from a modelled Southern Alps icefield. *Quaternary Science Reviews* 46, 30-45, 2012.
- Grant, G.R., Naish, T.R., Dunbar, G.B., Stocchi, P., Kominz, M.A., Kamp, P.J.J., Tapia, C.A., McKay, R.M., Levy, R.H., Patterson, M.O. The amplitude and origin of sea-level variability during the Pliocene epoch. *Nature*, 574, 237-241.
- 1330 Langridge, R.M., Ries, W.F., Litchfield, N.J., Villamor, P., Van Dissen, R.J., Barrell, D.J.A., Rattenbury, M.S., Heron, D.W., Rock, S., Townsend, D.B., Lee, J.M., Berryman, K.R., Nicol, A., Cox, S.C., Stirling, M.W. The New Zealand Active Faults Database. *New Zealand Journal of Geology and Geophysics*, 59, 86-96, Doi:10.1080/00288306.2015.1112818, 2016.
- Goldie, P.J. The Quaternary geology of an area north of Houhora. Unpublished Masters Thesis, University of Auckland, 139 pgs., 1975.
- Golledge, N.R. Long-term projections of sea-level rise from ice sheets. *WIREs Climate change*, 11, e634, 2020.
- 1335 Goodfriend, G.A., Kashgarian, M., Harasewych, M.G. Use of aspartic acid racemization and post-bomb ^{14}C to reconstruct growth rate and longevity of the deep-water slit shell *Entemnotrochus adansonianus*. *Geochimica et Cosmochimica Acta*, 69, 1125-1129, 1995.
- Grant-Mackie, J.A. and Scarlett, R.J. Last interglacial sequence, Oamaru. Guidebook for Excursions A7 and C7 central and southern Canterbury. DK INQUA Congress, Christchurch, New Zealand, 87-99, 1973.
- 1340 Hearty, P.J., Vacher, H.L., Mitterer, R.M. Aminostratigraphy and ages of Pleistocene limestones of Bermuda. *Geological Society of America Bulletin*, 104 (4), 471-480, 1992.
- Hearty, P.J., O'Leary, M.J., Kaufman, D.S., Page, M.C., Bright, J. Amino acid geochronology of individual foraminifer (*Pulleniatina obliquiloculata*) tests, north Queensland margin, Australia: A new approach to correlating and dating Quaternary tropical marine sediment cores. *Paleoceanography*, 19, PA4022, 2004.
- 1345 Heine, R.W. Marine terraces at Cape Terawhiti and Tounge Point, Wellington. *Journal of the Royal Society of New Zealand*, 4 (4), 485-492, 1974.
- Heine, R.W. Microanalysis of planation surfaces in the Port Nicholson Basin (Wellington City), and relationship of tectonic features. *New Zealand Journal of Geology and Geophysics*, 22 (3), 379-390, 1979.
- 1350 Heine, R.W. A new interpretation of the geomorphology of Wellington. *Journal of the Royal Society of New Zealand*, 12 (2), 189-205, 1982.
- Henderson, J. The Post-Tertiary history of New Zealand. *Transactions of the New Zealand Institute*, 55, 580-599, 1924.
- Hesp, P.A. and Shepherd, M.J. Some aspects of the late Quaternary geomorphology of the lower Manawatu Valley, New Zealand. *New Zealand Journal of Geology and Geophysics*, 21 (3), 403-412, 1978.
- 1355 Hicks, D.L. Geomorphic development of the southern Aupouri and Karikari Peninsulas with special reference to sand dunes. Unpublished Masters Thesis, University of Auckland, New Zealand, 257 pgs., 1975.
- Hull, A.G. Late Quaternary geology of the Cape Kidnappers area, Hawke's Bay, New Zealand. Unpublished Masters Thesis, University of Wellington, New Zealand, 140 pgs., 1985.
- Huntley, D.J., Lamothe, M. Ubiquity of anomalous fading in K-feldspars and the measurement and correction for it in optical dating. *Canadian Journal of Earth Sciences*, 38, 1093-1106, 2001.
- 1360 Hutton, F.W. Sketch of the geology of New Zealand. *Quarterly Journal of the Geological Society*, 41, 191-220, 1885.
- Iso, N., Okada, A., Ota, Y., Yoshikawa, T. Fission-track ages of late Pleistocene tephra on the Bay of Plenty coast, North Island, New Zealand. *Journal of Geology of Geophysics*, 25 (3), 295-303, 1982.
- Jacobs, Z., Roberts, R.G. Advances in optically stimulated luminescence dating of individual grains of quartz from archeological deposits. *Evolutionary Anthropology*, 16, 210-223, 2007.



- 1365 James, W.H.M., Carrivick, J.L., Quincey, D.J., Glasser, N.F. A geomorphology based reconstruction of ice volume distribution at the Last Glacial Maximum across the Southern Alps of New Zealand. *Quaternary Science Reviews*, 219, 20-35, 2019.
- Jobberns, G. The raised beaches of the north east coast of the South Island of New Zealand. *Transactions of the New Zealand Institute*, 59 (3), 508-570, 1928.
- Jobberns, G. and King, L.C. The nature and mode of origin of the Motunau Plain, North Canterbury, New Zealand. *Transactions of the New Zealand Institute*, 63, 355-369, 1933.
- 1370 Kamp, P.J.J. Stratigraphy and sedimentology of conglomerates in the Kidnappers Group, Hawkes Bay. Unpublished Masters Thesis, University of Waikato, New Zealand, 276 pgs., 1978.
- Kaplan, M.R., Strelin, J.A., Schaefer, J.M., Denton, G.H., Finkel, R.C., Schwartz, R., Putnam, A.E., Vandergoes, M.J., Goehring, B.M., Travis, S.G. In-situ cosmogenic ^{10}Be production rate at Lago Argentino, Patagonia: implications for late-glacial climate chronology. *Earth and Planetary Science Letters*, 309, 21-32, 2011.
- 1375 Kaufman, D.S., Cooper, K., Behl, R., Billups, K., Bright, J., Gardner, K., Hearty, P., Jakobsson, M., Mendes, I., O'Leary, M., Polyak, L., Rasmussen, T., Rosa, F., Schmidt, M. Amino acid racemization in mono-specific foraminifera from Quaternary deep-sea sediments. *Quaternary Geochronology*, 16, 50-61, 2013.
- Kear, D. and Waterhouse, B.C. Quaternary surfaces and sediments at Waihi Beach. *New Zealand Journal of Geology and Geophysics*, 4 (4), 434-445, 1961.
- 1380 Kennedy, D.M., Tannock, K.L., Crozier, M.J., Rieser, U. Boulders of MIS 5 age deposited by a tsunami on the coast of Otago, New Zealand. *Sedimentary Geology*, 200, 222-23, 2007.
- Kim, K.J. and Sutherland, R. Uplift rate and landscape development in southwest Fiordland, New Zealand, determined using ^{10}Be and ^{26}Al exposure dating of marine terraces. *Geochimica et Cosmochimica Acta*, 68 (10), 2313-2319, 2004.
- 1385 King, L.C. Raised beaches and other features of the south-east coast on the North Island of New Zealand. *Transactions of the New Zealand Institute*, 61 (3-4), 498-523, 1930.
- King, L.C. Notes on the geology and geomorphology of the coast between Napier and Castlepoint. *Transactions of the New Zealand Institute*, 63 (1), 72-79, 1932.
- King, D.J., Newnham, R.M., Gehrels, W.R., Clark, K.J. Late Holocene sea-level changes and vertical land movements in New Zealand. *New Zealand Journal of Geology and Geophysics*, DOI: 10.1080/00288306/2020.1761839, 2020.
- 1390 Kingma, J.T. Geology of the Te Aute Subdivision. New Zealand Department of Scientific and Industrial Research, No. 70-71, 1971.
- Kohn, B.P., Pillans, B., McGlone, M.S. Zircon fission track age for middle Pleistocene Rangitawa Tephra, New Zealand: stratigraphic and paleoclimatic significance. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 95, 73-94, 1992.
- 1395 Land Information New Zealand. New Zealand Coordinate Conversions. Licensed by Land Information New Zealand (LINZ) for re-use under the Creative Commons Attribution 4.0 International license, <https://www.geodesy.linz.govt.nz/concord/index.cgi> 2020a.
- Land Information New Zealand. NZ Terrain Relief (Topo250). Licensed by Land Information New Zealand (LINZ) for re-use under the Creative Commons Attribution 4.0 International license, <https://data.linz.govt.nz/layer/50766-nz-terrain-relief-topo250/> 2020b.
- 1400 Leamy, M.L. Pleistocene shorelines at Porirua Harbour, near Wellington, New Zealand. *New Zealand Journal of Geology and Geophysics*, 1 (1), 95-102, 1958.
- Lee, J.M., Begg, J.G. Geology of the Wairarapa area. Institute of Geological & Nuclear Sciences 1:250 000 geological map 11. Lower Hutt, New Zealand. Institute of Geological and Nuclear Sciences Limited, 66 pgs., 2002.
- 1405 Li, B., Roberts, R.G., Jacobs, Z. On the dose dependency of the bleachable and non-bleachable components of IRSL from K-feldspar: Improved procedures for luminescence dating of Quaternary sediments. *Quaternary Geochronology*, 17, 1-13, 2013.



- Li, B., Jacobs, Z., Roberts, R.G., Li, S.-H. Review and assessment of the potential of post-IR IRSL dating methods to circumvent the problem of anomalous fading in feldspar luminescence. *Geochronometria*, 41, 178-201, 2014.
- 1410 Litchfield, N.J. and Lian, O.B. Luminescence age estimates of Pleistocene marine terrace and alluvial fan sediments associated with tectonic activity along coastal Otago, New Zealand. *New Zealand Journal of Geology and Geophysics*, 47 (1), 29-37, 2004.
- 1415 Litchfield, N.J., Van Dissen, R., Sutherland, R., Barnes, P.M., Cox, S.C., Norris, R., Beavan, R.J., Langridge, R., Villamor, P., Berryman, K., Stirling, M., Nicol, A., Nodder, S., Lamarche, G., Barrell, D.J.A., Pettinga, J.R., Little, T., Pondard, N., Mountjoy, J.J., Clark, K. A model of active faulting in New Zealand. *New Zealand Journal of Geology and Geophysics*, 57 (1), 32-56, 2014.
- Lorscheid, T. and Rovere, A. The indicative meaning calculator – quantification of paleo sea-level relationships by using global wave and tide datasets. *Open Geospatial Data, Software and Standards*, 4 (10), 2019.
- 1420 Lowe, D.J., Tippet, M., Kamp, P.J.J., Liddell, I.J., Briggs, R.M., Horrocks, J.L. Ages on weathered Plio-Pleistocene tephra sequences, western North Island, New Zealand. In: *Tephra: Chronology, Archaeology*, edited by: Juvigné, E.T., Raynal, J.-P. CDERAD éditeur, Goudet, Les Dossiers de l'Archéo-Logis, 1, 45-60, 2001.
- Madsen, A.T., Buylaert, J.P., Murray, A.S. Luminescence dating of young coastal deposits from New Zealand using feldspar. *Geochronometria*, 38 (4), 378-390, 2011.
- Martinson, D.G., Pisias, N.G., Hays, J.D., Moore, Jr., T.C., Shackleton, N.J. Age dating and orbital theory of ice ages: development of a high resolution 0 to 300,000-year chronostratigraphy. *Quaternary Research*, 27, 1-29, 1987.
- 1425 McGlone, M.S., Neall, V.E., Pillans, B.J. Inaha Terrace deposits: A late Quaternary terrestrial record in south Taranaki, New Zealand. *New Zealand Journal of Geology and Geophysics*, 27 (1), 35-49, 1984.
- Mildenhall, D.C. Appendix 1 of Quaternary geology of the North Kaipara Barrier, Northland, New Zealand by Richardson, R.J.H. *New Zealand Journal of Geology and Geophysics*, 28 (1), 121-127, 1985.
- 1430 Mildenhall, D.C. Pleistocene palynology of the Petone and Seaview drillholes, Petone, Lower Hutt Valley, North Island, New Zealand. *Journal of the Royal Society of New Zealand*, 25 (2), 207-262, 1995.
- Moar, N.T. Appendix of Late Quaternary terraces between Ship Creek and the Whakapohai River, south Westland, New Zealand by Nathan, S. and Moar, N.T. *Journal of the Royal Society of New Zealand*, 5 (3), 323-327, 1975.
- 1435 Moar, N.T. and Mildenhall, D.C. Appendix 1 of Stratigraphy of the late Quaternary deposits of the northern Canterbury Plains, New Zealand by Brown, L.J., Wilson, D.D., Moar, N.T., Mildenhall, D.C. *New Zealand Journal of Geology and Geophysics*, 31 (3), 331-335, 1988.
- Murray-Wallace, C.V. Pleistocene coastal stratigraphy, sea-level highstands and neotectonism of the southern Australian passive continental margin-a review. *Journal of Quaternary Science*, 17, 469-489, 2002.
- Murray-Wallace, C.V. and Belperio, A.P. Identification of remanié fossils using amino acid racemization. *Alcheringa: An Australian Journal of Palaeontology*, 18 (3), 219-227, 1994.
- 1440 Murray-Wallace, C.V., Beu, A.G., Kendrick, G.W., Brown, L.J., Belperio, A.P., Sherwood, J.E. Palaeoclimatic implications of the occurrence of the arcoid bivalve *Anadara trapezia* (Deshayes) in the Quaternary of Australasia. *Quaternary Science Reviews*, 19, 559-590, 2000.
- 1445 Murray-Wallace, C.V., Bourman, R.P., Prescott, J.R., Williams, F., Price, D.M., Belperio, A.P. Aminostratigraphy and thermoluminescence dating of coastal aeolianites and the later Quaternary history of a failed delta: The River Murray mouth region, South Australia. *Quaternary Geochronology*, 5, 28-49, 2010.
- Murray-Wallace, C.V., Belperio, A.P., Dosseto, A., Nicholas, W.A., Mitchell, C., Bourman, R.P., Eggins, S.M., Grün, R. Last interglacial (MIS 5e) sea-level determined from a tectonically stable, far-field location, Eyre Peninsula, southern Australia. *Australian Journal of Earth Sciences*, 63, 611-630, 2016.



- 1450 Nathan, S. Geological Map of New Zealand 1: 63 360, Sheets S23 and S30 Foulwind and Charleston. New Zealand Department of Science and Industrial Research, 1975.
- Nathan, S. and Moar, N.T. Late Quaternary terraces between Ship Creep and the Whakapohai River, south Westland, New Zealand. *Journal of the Royal Society of New Zealand*, 5 (3), 313-327, 1975.
- Nichol, S.L. Morphology, stratigraphy and origin of last interglacial Beach Ridges at Bream Bay, New Zealand. *Journal of Coastal Research*, 18 (1), 149-159, 2002.
- 1455 Nicol, A., Seebeck, H., Wallace, L. Quaternary Tectonics of New Zealand. In: *Landscape and Quaternary Environmental Change in New Zealand*, edited by: Shulmeister J. Atlantis Press, Paris, Atlantis Advances in Quaternary Science. 3, 1-34, 2017.
- Oakley, D.O.S., Kaufman, D.S., Gardner, T.W., Fisher, D.M., VanderLeest, R.A. Quaternary marine terrace chronology, North Canterbury, New Zealand, using amino acid racemization and infrared-stimulated luminescence. *Quaternary Research*, 87, 151-167, 2017.
- 1460 Oakley, D.O.S., Fisher, D.M., Gardner, T.W., Stewart, M.K. Uplift rates of marine terraces as a constraint on fault-propagation fold kinematics: Examples from the Hawkswood and Kate anticlines, North Canterbury, New Zealand. *Tectonophysics*, 724-725, 195-219, 2018.
- Oliver, R.L. The Otaki Sandstone and its Geological History. New Zealand Department of Scientific and Industrial Research, Geological Memoirs, No. 7, 51 pgs., 1948.
- 1465 Ongley, M. The Geology of the Kaitangata-Green Island subdivision, eastern and central Otago Divisions. New Zealand Department of Scientific and Industrial Research, Bulletin No. 38, 90 pgs., 1939.
- Ota, Y., Yoshikawa, T., Iso, N., Okada, A., Yonekura, N. Marine terraces of the Conway coast, South Island, New Zealand. *New Zealand Journal of Geology and Geophysics*, 27 (3), 313-325, 1984.
- 1470 Ota, Y., Omura, A., Iwata, H. ^{230}Th - ^{238}U age of Rotoehu Ash and its implications for marine terrace chronology of eastern Bay of Plenty, New Zealand. *New Zealand Journal of Geology and Geophysics*, 32 (3), 327-331, 1989.
- Ota, Y., Pillans, B., Berryman, K., Beu, A., Fujimori, T., Miyauchi, T., Berger, G., Beu, A.G., Climo, F.M. Pleistocene coastal terraces of Kaikoura Peninsula and the Marlborough coast, South Island, New Zealand. *New Zealand Journal of Geology and Geophysics*, 39 (1), 51-73, 1996.
- 1475 Pain, C.F. Late Quaternary dune sands and associated deposits near Aotea and Kawhia Harbours, North Island, New Zealand. *Journal of Geology and Geophysics*, 19 (2), 153-177, 1976.
- Palmer, A.S., Barnett, R., Pillans, B.J., Wilde Loess, river aggradation terraces and marine benches at Otaki, southern North Island, New Zealand. In: *Loess, its distribution, geology and soils*, edited by: Eden, D.N., Furkert, R.J., A.A. Balkema, Rotterdam, 163-174, 1988.
- 1480 Pillans, B. Upper Quaternary marine terrace chronology and deformation, south Taranaki, New Zealand. *Geology*, 11, 292-297, 1983.
- Pillans, B. Pleistocene marine terraces in New Zealand: A review. *New Zealand Journal of Geology and Geophysics*, 33 (2), 219-231, 1990a.
- Pillans, B. Late Quaternary Marine Terraces South Taranaki-Wanganui, Sheet Q22 and part sheets Q20, Q21, R21, & R22. New Zealand Geological Survey Miscellaneous Series Map 18, 1990b.
- 1485 Pillans, B. Quaternary Stratigraphy of Whanganui Basin – A Globally Significant Archive. In: *Landscape and Quaternary Environmental Change in New Zealand*, edited by: Shulmeister J. Atlantis Press, Paris, Atlantis Advances in Quaternary Science. 3, 141-170, 2017.
- Pillans, B.J. and Kohn, B.P. Rangitawa Pumice: a widespread (?) Quaternary marker bed in Taranaki-Wanganui. *Victoria University of Wellington Geology Department, Publication 20*, 94-104, 1981.
- 1490



- Pillans, B., Kohn, B.P., Berger, G., Froggatt, P., Duller, G., Alloway, B., Hesse, P. Multi-method dating comparison for mid-Pleistocene Rangitawa Tephra, New Zealand. *Quaternary Science Reviews*, 15, 641-653, 1996.
- Powers, W.E. Terraces of the Hurunui River, New Zealand. *New Zealand Journal of Geology and Geophysics*, 5 (1), 114-129, 1962.
- 1495 Preusser, F., Anderson, B.G., Denton, G.H., Schlüchter, C. Luminescence chronology of Late Pleistocene glacial deposits in North Westland, New Zealand. *Quaternary Science Reviews*, 24, 2207-2227, 2005.
- Preusser, F., Ramseyer, K., Schlüchter, C. Characterisation of low OSL intensity quartz from the New Zealand Alps. *Radiation Measurements*, 41, 871-877, 2006.
- Pullar, W.A. and Grant-Mackie, J.A. Correlation of the Te Papa and Otamaroa Terraces, Eastern Bay of Plenty. *New Zealand Journal of Geology and Geophysics*, 15 (1), 186-191, 1972.
- 1500 Putnam, A.E., Schaefer, J.M., Barrell, D.J.A., Vandergoes, M., Denton, G.H., Kaplan, M.R., Finkel, R.C., Schwartz, R., Goehring, B.M., Kelley, S.E. In situ cosmogenic ^{10}Be production-rate calibration from the Southern Alps, New Zealand. *Quaternary Geochronology*, 5, 392-409, 2010.
- Raine, J.I., Beu, A.G., Boyes, A.F., Campbell, H.J., Cooper, R.A., Crampton, J.S., Crundwell, M.P., Hollis, C.J., Morgans, H.E.G. Revised calibration of the New Zealand Geological Timescale: NZGT2015/1. *GNS Science Report 2012/39*, 53 pgs., 2015.
- 1505 Rattenbury, M.S., Townsend, D.B., Johnston, M.R. Geology of the Kaikoura area. Institute of Geological & Nuclear Sciences 1:250 000 geological map 13. Lower Hutt, New Zealand, GNS Science, 1 sheet + 70 p., 2006.
- Rees-Jones, J., Rink, W.J., Norris, R.J., Litchfield, N.J. Optical luminescence dating of uplifted marine terraces along the Akatore Fault near Dunedin, South Island, New Zealand. *New Zealand Journal of Geology and Geophysics*, 43 (3), 419-424, 2000.
- Richardson, R.J.H. Quaternary geology of the North Kaipara Barrier, Northland, New Zealand. *New Zealand Journal of Geology and Geophysics*, 28 (1), 111-127, 1985.
- 1515 Ricketts, B.D. Petrology and provenance of Pleistocene deposits in the south Parengarenga-Te Kao district, northern New Zealand. *New Zealand Journal of Geology and Geophysics*, 22 (1), 21-27, 1975.
- Roberts, R.G., Jacobs, Z., Li, B., Jankowski, N.R., Cunningham, A.C., Rosenfeld, A.B. Optical dating in archaeology: thirty years in retrospect and grand challenges for the future. *Journal of Archaeological Science*, 56, 41-60, 2015.
- Rovere, A., Raymo, M.E., Vacchi, M., Lorscheid, T., Stocchi, P., Gómez-Pujol, L., Harris, D.L., Casella, E., O'Leary, M.J., Hearty, P. The analysis of Last Interglacial (MIS 5e) relative sea-level indicators: Reconstructing sea-level in a warmer world. *Earth-Science Reviews*, 159 (404-427), 2016.
- 1520 Rovere, A., Ryan, D.D., Murray-Wallace, C.V., Simms, A., Vacchi, M., Dutton, A., Lorscheid, T., Chutcharavan, P., Brill, D., Bartz, M., Jankowski, N., Mueller, D., Cohen, K., Gowan, E. Descriptions of database fields for the World Atlas of Last Interglacial Shorelines (WALIS). doi:10.5281/zenodo.3961544, 2020.
- Rui, X., Guo, Y.J., Zhang, J.F., Hu, Y., Mei, H.J., Wang, Y.P., Xie, F., Li, B. Luminescence chronology of the Palaeolithic–Neolithic transition in the Yujiagou site at the Nihewan Basin, northern China. *Journal of Quaternary Science*, 34, 125-137, 2019.
- 1525 Rutter, N.W. and Vlahos, C.K. Amino acid racemization kinetics in wood; Applications to geochronology and geothermometry. *Geological Society of America, Special Paper 227*, 51-67, 1988.
- Ryan, D.D., Clement, A.J.H., Jankowski, N.R., Stocchi, P., Rovere, A. The last interglacial sea-level record of New Zealand (Aotearoa) – WALIS database of sea-level indicators [Data set]. Zenodo. <http://doi.org/10.5281/zenodo.4056376>
- 1530 Ryan, D.D., Lachlan, T.J., Murray-Wallace, C.V., Price, D.M. The utility of single foraminifera amino acid racemization analysis for the relative dating of Quaternary beach barriers and identification of reworked sediment. *Quaternary Geochronology*, 60, 101103, 2020b.



- 1535 Shackleton, N.J., Berger, A., Peltier, W.R. An alternative astronomical calibration of the lower Pleistocene timescale based on ODP Site 677. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 81, 251-261, 1990.
- Selby, M.J., Pullar, W.A., McCraw, J.D. The age of Quaternary surfaces at Waihi Beach. *Earth Science Journal*, 5 (2), 106-113, 1971.
- Sewell, A.H. Paleoenvironmental analysis of Quaternary strata in the Levin Area. Unpublished Masters Thesis, Massey University, New Zealand, 182 pgs., 1991.
- 1540 Shulmeister, J. Soons, J.M., Berger, G.W., Harper, M., Holt, S., Moar, N., Carter, J.A. Environmental and sea-level changes on Banks Peninsula (Canterbury, New Zealand) through three glaciation-interglaciation cycles. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 152, 101-127, 1999.
- Simms, A.R., Rouby, H., Lambeck, K. Marine terraces and rates of vertical tectonic motion: The importance of glacio-isostatic adjustment along the Pacific coast of central North America. *GSA Bulletin*, 128 (1/2), 81-93, 2016.
- 1545 Sloss, C.R., Murray-Wallace, C.V., Jones, B.G. Holocene sea-level change on the southeast coast of Australia: a review. *The Holocene*, 17, 999-1014, 2007.
- Smedley, R.K., Duller, G.A.T. and Roberts, H.M. Bleaching of the post-IR IRSL signal from individual grains of K-feldspar: Implications for single-grain dating. *Radiation Measurements*, 79, pp.33-42, 2015.
- 1550 Suggate, R.P. Late Pleistocene geology of the northern part of the South Island, New Zealand. *New Zealand Geological Society*, 77, 1965.
- Suggate, R.P. The glacial/interglacial sequence of North Westland, New Zealand. *New Zealand Geological Survey, Record* 7, 1985.
- Suggate, R.P. Differential uplift of middle and late Quaternary shorelines, northwest South Island, New Zealand. *Quaternary International*, 15/16, 47-59, 1992.
- 1555 Taylor-Silva, B.I., Stirling, M.W., Litchfield, N.J., Griffin, J.D., van den Berg, E.J., Wang, N. Paleoseismology of the Akatore Fault, Otago, New Zealand. *New Zealand Journal of Geology and Geophysics*, 63, 151-167, 2020.
- Te Punga, M.T. Some geological features of the Otaki-Waikanae district. *New Zealand Journal of Geology and Geophysics*, 5 (4), 517-530, 1962.
- 1560 Thiel, C., Buylaert, J.-P., Murray, A., Terhorst, B., Hofer, I., Tsukamoto, S., Frechen, M. Luminescence dating of the Stratzing loess profile (Austria) – Testing the potential of an elevated temperature post-IR IRSL protocol. *Quaternary International*, 234, 23-31, 2011.
- Wallace, L.M., Beacan, J., McCaffrey, R., Darby, D. Subduction zone coupling and tectonic block rotations in the North Island, New Zealand. *Journal of Geophysical Research: Solid Earth*, 109, B12406, 2004.
- 1565 Ward, C.M. Marine terraces of the Waitutu district and their relation to the late Cenozoic tectonics of the southern Fiordland region, New Zealand. *Journal of the Royal Society of New Zealand* 18 (1), 1-28, 1988a.
- Ward, C.M. New Zealand Marine Terraces: Uplift Rates. *Science*, 240 (4853), 803-804, 1988b.
- Webby, B.D. Quaternary deposits in the Porirua Area, New Zealand. *New Zealand Journal of Geology and Geophysics*, 7 (2), 371-388, 1964.
- 1570 Wehmiller, J.F., York, L.L., Bart, M.L. Amino acid racemization geochronology of reworked Quaternary mollusks on U.S. Atlantic coast beaches: implications for chronostratigraphy, taphonomy, and coastal sediment transport. *Marine Geology*, 124, 303-337, 1995.
- Wehmiller, J.F. United States Quaternary coastal sequences and molluscan racemization geochronology – What have they meant for each other over the past 45 years? *Quaternary Geochronology*, 16, 3-20, 2013.



- 1575 Wellman, H.W. and Wilson, A.T. Notes on the Geology and Archaeology of the Martins Bay District. *New Zealand Journal of Geology and Geophysics*, 7 (4), 702-721, 1964.
- Williams, D.N. Pliocene and Quaternary geology of Mana Island (Note). *New Zealand Journal of Geology and Geophysics*, 21 (5), 653-656, 1978.
- Williams, P.W. Speleothem dates, Quaternary terraces and uplift rates in New Zealand. *Nature*, 298, 257-260, 1982.
- 1580 Williams, P.W., McGlone, M., Neil, H., Zhao, J-X. A review of New Zealand palaeoclimate from the Last Interglacial to the global Last Glacial Maximum. *Quaternary Science Reviews*, 110, 92-105, 2015.
- Wilson, K., Litchfield, N., Berryman, K., Little, T. Distribution, age, and uplift patterns of Pleistocene marine terraces of the northern Raukumara Peninsula, North Island, New Zealand. *New Zealand Journal of Geology and Geophysics*, 50 (3), 181-191, 2007.
- Wintle, A.G. Anomalous fading of thermo-luminescence in mineral samples. *Nature*, 245, 143-144, 1973.
- 1585 Yoshikawa, T., Ota, Y., Yonekura, N., Okada, A., Iso, N. Marine terraces and their deformation on the northeast coast of the North Island, New Zealand. *Geographical Review of Japan*, 53, 238-262, 1980.
- Yousif, H.S. The applications of remote sensing to geomorphological neotectonic mapping in North Canterbury New Zealand. Unpublished PhD Thesis. University of Canterbury, 1989.
- 1590 Zumberge, J.E., Engel, M.H., Nagy, B. Amino acids in Bristlecone Pine: An evaluation of factors affecting racemization rates and paleothermometry. In: *Biogeochemistry of Amino Acids*, edited by: Hare, P.E., Hoering, T.C., and King, K. Wiley, New York, pp 503-525, 1980.