



## Late Cretaceous – Early Palaeogene inversion-related tectonic structures at the NE margin of the Bohemian Massif (SW Poland and northern Czechia)

Andrzej Głuszyński<sup>1</sup>, Paweł Aleksandrowski<sup>2,1</sup>

5 <sup>1</sup>Polish Geological Institute, Rakowiecka 4, PL 00-975 Warszawa, Poland

<sup>2</sup>Institute of Geological Sciences, University of Wrocław, Cybulskiego 32, PL 50-205 Wrocław, Poland

*Correspondence to:* Andrzej Głuszyński (andrzej.gluszynski@pgi.gov.pl)

**Abstract.** A brief, regional-scale review of the Late Cretaceous – Early Palaeogene inversion-related tectonic structures affecting the Sudetes and their foreland at the NE margin of the Bohemian Massif is presented and complemented with results of new seismic studies. The Sudetes expose Variscan-deformed basement, partly overlain by post-orogenic Permo-Mesozoic cover, containing a wide spectrum of tectonic structures, both brittle and ductile, in the past in this area referred to as young Saxonian or Laramide. We have used newly reprocessed legacy seismics to study these structures at the two main post-Variscan structural units of the area, the North-Sudetic and Intra-Sudetic synclinoria, and discuss the results together with regionally-distributed examples coming from quarries and underground mines as well as those from the literature. The Late Cretaceous – Early Palaeogene tectonic structures in consecutively reviewed Sudetic tectonic units, from the north to south, typically include gentle to moderate buckle folds of detachment type or fault-related, high-angle reverse and normal faults, as well as low-angle thrusts – often rooted in the crystalline basement. The structures hitherto described as grabens, are frequently believed to be bounded by reverse faults (hence we use the term ‘reverse grabens’) and typically reveal strongly synclinal pattern of their sedimentary fill. The crystalline basement top, as imaged by seismic sections in the North Sudetic Synclinorium below the detachment-folded cover, is synformally down-warped with a wavelength of up to 30 km, whereas on the elevated areas, where the basement top is exposed at the surface, it is up-warped (i.e. tectonically buckled). The reviewed compressional structures typically show an orientation fitting the regionally-known Late Cretaceous – Early Palaeogene tectonic shortening direction of NE-SW to NNE-SSW. The same applies to the regional joint pattern, typically comprising an orthogonal system of steep joints of c. NW-SE and NE-SW strikes. All the reviewed structures are considered as due to the Late Cretaceous – Early Palaeogene tectonic shortening episode, although some of the discussed faults with a strike-slip component of motion may have been modified, or even produced, by later, Late Cenozoic, tectonism.



## 1 Introduction

During Late Cretaceous to Early Palaeogene times, compressional tectonic structures developed over vast areas of western and north-central Europe (e.g. Kley and Voigt, 2008; Navabpour et al., 2017; Kley, 2018; Nádaskay et al., 2019; Malz et al., 2020; Voigt et al., 2021). They are currently interpreted as due to collisional interaction of the Iberian mass with continental Europe and the resultant propagation of far-field compressional stress (Kley & Voigt 2008), which brought about widespread inversion of Permo-Mesozoic basins. In Poland and easternmost Germany (e.g. Krzywiec, 2002, 2006; Mazur et al., 2005; Krzywiec and Stachowska, 2016; Kley 2018) the resultant map-scale tectonic structures are mostly NW-SE-trending, large-scale (crustal-scale?), gentle folds up to 1000 km long, 150-250 km wide and up to at least 3 km high (e.g. Pożaryski, 1979). The Mid-Polish Swell (or Anticlinorium), spanning the entire breadth of Poland's NW-SE diagonal from the Baltic Sea to westernmost Ukraine, is the most spectacular example of such structures. To the SW of the Mid-Polish Swell there is the Szczecin-Łódź Trough (Synclinorium), which occurs next to the successive elevated structural element, occurring at the NE margin of the Bohemian Massif, in the borderland between Poland and Czechia. This is the Sudetic area - the principal object to be presented in this paper, whose goal is to briefly overview the wide spectrum of structural effects produced by the Late Cretaceous to Early Cenozoic trans-European compressional event at the NE margin of the Bohemian Massif, in the Sudety Mts and in their northern foreland with a similar geology. An important part of the review is based on new data derived by us from recently reprocessed legacy seismic materials coming from the central areas of the Sudetic region. The other presented examples supply the wide regional context for the seismic results and come either from our own collection or from descriptions made by other authors, though sometimes interpreted by us in a different way. Although the tectonic structures that formed at the turn of Cretaceous and Palaeogene times obviously must have developed also in the Variscan basement, they are – in general - difficult to distinguish from the older ones. Therefore, the scope of this paper is mostly limited to the tectonic phenomena that occur in the post-orogenic Permo-Mesozoic strata, though, in places also in the late-orogenic Upper Carboniferous.

Below, following the sections on outline geology and methods, an overview of the Late Cretaceous – Early Palaeogene deformation structures in the Sudetes is presented, mostly in a regional context and order, in reference to particular structural units of the Sudetes, moving from the north to south across the area.

## 2 Outline geology

The Sudetes, low-topography mountains located on the northeastern margin of the Bohemian Massif, together with the southern part of the Silesian-Lusatian Plain share the same pre-Cenozoic geological composition and structure and – in the geological sense – are usually considered collectively as the Sudetic region, or simply, the Sudetes (e.g. Aleksandrowski and Mazur, 2002; Mazur et al., 2006, Kroner et al., 2008). The differences in the topography – mountainous versus lowland - are due to splitting of the Sudetic area into two tectonic blocks: the currently downthrown Fore-Sudetic block in the north and the elevated Sudetic one in the south. The Sudetes constitute the northeasternmost segment of the Central European Variscan



60 internides, exposing crystalline basement of strongly deformed and metamorphosed Late Neoproterozoic to Carboniferous rocks abundantly intruded by Carboniferous granites (e.g. Mazur et al., 2006, 2007, 2020). In the Sudetes, the main tectonostratigraphic domains of the European Variscides: the Moldanubian, Tepla-Barrandian and – particularly – Saxothuringian find their continuation (e.g. Martinez-Catalan et al., 2021).

The **Sudety Mountains** acquired their present-day mountainous relief due to Neogene uplift of the Sudetic Block in front of 65 the then actively growing Alps and Carpathians (e.g. Żelaźniewicz and Aleksandrowski, 2008; Jarosiński et al., 2009; Żelaźniewicz et al., 2011). This uplift affected the area mostly planated during the Palaeogene and early Miocene times, following the earlier, Late Cretaceous – Early Palaeogene more prominent uplift, which at that time had occurred over a broader area, including also the Fore-Sudetic Block to the north (Fig. 1), which at present is downthrown along the Sudetic Boundary Fault (e.g. Cloos, 1922; Teisseyre, 1957; Oberc 1972).

70 The Late Cretaceous – Early Palaeogene uplift exhumed the Variscan basement from below the post-Variscan, uppermost Carboniferous through Permo-Mesozoic cover and left the Sudetes tectonically elevated with respect to the adjoining depressed areas: the Fore-Sudetic Homocline to the north and the North Bohemian Cretaceous Basin to the south, which managed to preserve much more of their Permo-Mesozoic post-Variscan sedimentary fill. For over a century, the Sudetes have been considered a part of the classical area in Central Europe, in which the term “Saxonische Tektonik” (cf. Kley, 75 2013) was applied to deformation structures, mostly fault-related, observed in the Permo-Mesozoic sedimentary basins. In the Sudetes, in the older literature these contractional structures were most often ascribed specifically to the young Saxonian (Closs, 1922; Beyer, 1939; Oberc, 1972, 1977) or Laramide tectonism (Oberc 1972, 1977).

### 3 Data and methods

The concise regional, though not fully systematic, review of the tectonic structures of likely Late Cretaceous – Early 80 Palaeogene age presented in this paper is based (1) on authors’ own analysis and structural interpretation of the newly reprocessed reflection seismic data coming from the two key areas in the Sudetes with the most complete post-Variscan sedimentary record and (2) on structural field data of the present authors or (3) **coming** from critically assessed literature descriptions. The analysed seismic data were acquired by mining companies in the end 1970s (1976-1980) in the **North Sudetic Synclinorium** and in the early 1990s (1991-1993) in the **Intra-Sudetic Synclinorium** (Fig. 2). We received them as 85 raw seismic data (field data, partly pre-processed, reloaded to discs from original magnetic tapes) and ~~the~~ accompanying-paper prints. These seismic data are unique at the scale of the entire Sudetes as ~~the~~ source of valuable information on the structural geology of the Permo-Mesozoic succession in ~~the~~ both main Sudetic synclinal units.

The reprocessing of, in total, c. 650 km of the seismic profiles was carried out in 2019, using ~~an~~ up-to-date oil-industry software (Głuszynski and Smajdor, 2020). The new processing included post stack time migration (PostSTM), while to some 90 of the profiles also the pre stack time migration stage (PreSTM) procedure was applied. No time-depth conversion was, however, attempted because of too scarce coverage of the area with appropriate drillhole data. As complementary material



we used also an analogue/paper print version of a seismic profile from the vicinities of Broumov in Czechia and also some seismic profiles coming from the Fore-Sudetic Homocline. A significant part of the structurally interpreted seismic profiles are presented in this paper. Our structural interpretation of these profiles provides an entirely new material showing the tectonic style and geometry of the compressional tectonic structures affecting the post-Variscan sedimentary cover on the NE margin of the Bohemian massif and allows for inferences as to the mechanics of their formation. An equally important part of the paper, aimed at giving its readers an overall information on the distribution and genetic and geometrical diversity of the tectonic structures that formed or evolved under the Late Cretaceous – Early Palaeogene compressional regime, are depiction and short description of a selection of such structures exposed in natural outcrops and in active mines and quarries throughout the entire Sudetic area or its direct vicinities. This material is either our own or is based on critically evaluated ~~other authors'~~ accounts.

#### 4. Products of Late Cretaceous – Early Palaeogene tectonic shortening: examples from particular structural domains

##### 4.1. The north-eastern margin of the Sudetes: transition to the Fore-Sudetic Homocline

Directly to the northeast of the Sudetic Block, separated by the Middle Odra Fault Zone, the Fore-Sudetic Homocline extends (e.g. Kłapciński et al., 1984; Kroner et al., 2008; Żelaźniewicz et al. 2011). It is defined by **shallowly** (1-5°) NE-dipping Permian to Mesozoic strata (Figs. 2 and 3) on top of Variscan-folded Carboniferous (Mazur et al., 2006, 2010), representing the SW limb of the Szczecin-Miechów Synclinorium of the Polish-German Basin. The Permian-Mesozoic succession ~~constitutes~~ the fill of the extensive Polish-German Basin and unconformably overlies the Carboniferous fold-and-thrust flysch belt of the Variscan externalides (e.g. Mazur et al., 2010, 2020).

The Permo-Mesozoic strata in the southern part of the the Fore-Sudetic Homocline are very gently folded and in few places affected by steep to **low-angle reverse** faults (Fig. 4). In the close vicinity of the NW-SE-trending Middle Odra Fault system, however, in galleries of c. 0.5-1.2 km-deep mines of the Legnica-Głogów Copper District, a rich inventory of contractional and, also, extensional tectonic structures occur in the base Zechstein/top Rotliegend ore-bearing formation, containing clayey shale, carbonates, anhydrite and rock-salt, apart from sandstones. These structures typically include **thrust** faults, accompanied by bedding-parallel décollements, as well as detachment folds ~~of origin easily explainable by~~ NE-SW directed tectonic shortening. The meso-scale structures can be studied directly on the galleries' walls (Fig. 5) and were described for decades by e.g. Salski (1965, 1968), Oberc and Salski (1968), Dumicz and Don (1977, 1990), Markiewicz and Szarowski (1990), Żelaźniewicz and Markiewicz (1991). Larger structures appear on mining maps as a complex network of faults, whose pattern reflects reactivation of a few major and numerous minor NW-SE trending fractures propagating upwards from the Variscan basement, which are accompanied by a number of relatively large WSW-ENE en-echelon **events** (cf. Markiewicz, 2007), the latter most probably formed due to a Late Cretaceous – Early Palaeogene (or Late Cenozoic?) sinistral strike-slip activity of the major NW-SE faults.



## 125 4.2. The Fore-Sudetic Block

At the base Cenozoic level, the Fore-Sudetic Block (Figs. 2 and 3) **exposes** Variscan basement rocks of various metamorphic grades, varying in age from the Neoproterozoic to probable Devonian (Pożaryski, 1979; Kotański, 1997; Cwojdzński and Żelaźniewicz, 1995; Cymerman, 2010), and, variable igneous plutonic rocks, mostly of **basic** to ultrabasic Silurian/Early Devonian ophiolitic suite and Carboniferous to Early Permian, granitoids (e.g. Mazur et al., 2006, 2007; Kroner et al., 2008).

130 As the Fore-Sudetic Block is (except at its easternmost **periphery**) devoid of Permo-Mesozoic deposits and its crystalline basement is eroded deeper than that in the mountainous Sudetes, an idea was conceived long ago (Cloos, 1922; Teisseyre, 1957) that the **Block was** uplifted with respect to the present-day Sudetes across the Sudetic Boundary Fault, following the Late Cretaceous – Early Palaeogene inversion. This situation was subsequently reversed in the Late Miocene, when the Sudetes were uplifted and the Fore-Sudetic Block downthrown when the Carpathian-Alpine forebulge was formed (e.g.

135 Żelaźniewicz et al., 2011). The lack of Permo-Mesozoic sedimentary cover hampers the direct recognition of the effects of the Late Cretaceous – Early Palaeogene tectonism on the Fore-Sudetic Block, which, nevertheless, can be partly assessed thanks to the presence of the ~~so-called~~ Żary Pericline (Fig. 2) in the geological map intersection pattern. This pericline consists in map-view enveloping of the NW-projecting basement “peninsula” ~~with~~ the Permo-Triassic outcrop zone, which ~~continues to the south, onto the Block,~~ from the area of the Fore-Sudetic Homocline. This basement/cover intersection

140 pattern illustrates the likely up-warping (updoming) effect of the Late Cretaceous – Early Palaeogene compression, which seems to have undulated the roof surface of the crystalline basement, producing a very gentle NW-SE trending antiform. This antiform is analogous – but of reverse polarity - to the large, gentle synform that occurs to the south of it and is ~~legible~~ in seismic sections– in the floor of the North-Sudetic Synclinorium (see below). The uplift related to the formation of the antiform in question, must have been one of the important factors ~~facilitating~~ the erosion of the original Permo-Mesozoic

145 cover over the most part of the Fore-Sudetic Block.

## 4.3. The North Sudetic Synclinorium

The **North Sudetic Synclinorium** (Figs. 2, 3 and 6) is a fault-bounded, NE-SE elongated tectonic structure, c. 60 km long and up to 24 km wide (Beyer, 1939; Oberc, 1972, 1977; Solecki, 1994; Śliwiński et al., 2003; Kiersnowski *in* McCann et al., 2008a; Żelaźniewicz and Aleksandrowski, 2008; Żelaźniewicz et al., 2011). It is filled with post-Variscan, end

150 Carboniferous-Permian-Triassic and Upper Cretaceous (Cenomanian to Coniacian) continental and shallow marine sedimentary rocks, including also a Lower Permian volcanic member. The succession overlies a subsided top surface of the Variscan epimetamorphic basement **the Kaczawa Slate Belt** (e.g. Aleksandrowski and Mazur, 2002; Mazur et al., 2006; Żelaźniewicz and Aleksandrowski, 2008). The synclinorium achieved its quasi-basinal structure due to the Santonian-early Campanian to late Maastrichtian-Palaeocene (Walaszczyk *in* Voigt et al., 2008) tectonism. The North Sudetic Synclinorium

155 in its western and central parts defines, in general, a single synclinal structure, while at its eastern and southern extremities it



splits into several NW-SE elongated second-order synclinal elements, grabens and half-grabens, some of them separated by basement horsts (Figs. 2 and 7; Beyer, 1939; Śliwiński et al., 2003).

The end Mesozoic – early Cenozoic compressional deformation affected both the crystalline basement and post-Variscan cover. The cover must have once been relatively thick and must have extended over much wider areas of the Lower Silesia Block ~~at that time~~ than it does today (cf. e.g. Migoń and Danišik, 2012; Sobczyk et al., 2015, 2020). It was subsequently eroded from the most uplifted areas, but became preserved in places, where the basement/cover interface had undergone compression-driven down-warping. According to Beyer (1939), this compression in the North Sudetic basin brought about inversion of some of the earlier normal faults, which were then transformed into reverse ones (Fig. 7a). The Variscan basement at the upthrown sides of the former inverted faults was believed by this author to have acted like rigid jaws of a vice, which horizontally squeezed the Permo-Mesozoic succession in the process of a “frame-controlled folding”(German *Rahmenfaltung*, Fig. 7b).

The structural geometry of the Late Cretaceous – Early Palaeogene folds and related thrusts in the North Sudetic Synclinorium was known so far mostly from geological map intersection patterns combined with drilling results (Beyer, 1939; Oberc, 1972, 1977; Leśniak, 1979; Solecki, 1986, 1994; Cymerman, 1998) and from likely smaller-scale analogues occurring in active quarries (Fig. 8a-c). A reflection seismic survey made in 1976-1980 made it possible to better constrain the structure of the post-Variscan cover, which was attempted by Bałazinska and Bossowski (1979; their Fig. 3), applying an assumption of a dominance of nearly vertical faults.

A much closer insight into the geometry and structural style of the Late Cretaceous-deformed Permo-Mesozoic succession was recently enabled by the reprocessing of the same, legacy seismic raw data in 2019. The structural interpretation results of these data are presented in Figures 9 and 10 which depict the along-strike changing geometry of the Permo-Mesozoic succession of the North-Sudetic Synclinorium. This geometry, in general, seems to show products of **detachment** folding (cf. McClay, 1992) likely formed by buckling ~~mechanism~~ above the metamorphic basement. The structures are complicated by local thrusting/reverse faulting, but, nevertheless do not resemble typical fault-folds. The metamorphic basement seem **m** to have been shortened by down-warping that resulted in the formation of a single synform below the detached Permo-Mesozoic cover. The down-warping may have been associated with development of low-angle thrusts within the basement (not shown in Figs. 9 and 10), where some of the thrusts affecting the Permo-Mesozoic are likely rooted. The *Rahmenfaltung* at the synclinorium edges, invoked by Beyer (1939) and understood here as upthrowing the metamorphic basement on reverse (inverted?) faults, seems to also have its impact on the deformation process as the local source of horizontal shortening ~~forces~~. The folds interpreted from both the reprocessed seismic sections and from map intersection patterns, show, in general, near-parallel geometry, and shallow WNW-plunging (**Solecki, 1986, 1994, 2011**) axes. As already mentioned, they are locally affected by high- to medium-angle reverse/thrust faults, sometimes following their axial planes. Our analysis of newly reprocessed seismic profiles made possible a subsurface mapping of a number of high-angle fault zones. They are, as a rule, trending NW-SE and often continue ~~at~~ distances of tens of kilometres. Other identified major faults represent low-angle thrusts, continuing into the Variscan basement. The thrusting polarity in the Permo-Mesozoic



190 sucession of the North-Sudetic Synclinorium is bimodal and directed both to the NE and SW. ~~This seems typical of a~~  
detachment folding. The interpreted fault zones cut across the Permo-Mesozoic strata from the Zechstein at the bottom to the  
Upper Cretaceous at the top, which confirms their activity continued until at least the Late Cretaceous.

The structural analysis of the reprocessed seismics, made clear the necessity for reinterpretation of the hitherto widely held  
concept of the internal structure for the North-Sudetic Synclinorium, assuming the dominance of high-angle block tectonics.  
195 In a new concept, the significance of low-angle thrust faults, of compressional down-warping of the top basement surface,  
and of the well-developed detachment folding pattern ~~should be~~ taken into account.

A recent mapping and structural study by Kowalski (2020) of one of the already mentioned second-order synclinal/graben  
elements branching off from the SE rim of the North Sudetic Synclinorium, the Wleń Graben (Figs. 2 and 7) offers an  
opportunity to have a closer look on another example of a Late-Cretaceous – Early Palaeogene macro-scale tectonic structure  
200 in the NW part of the Sudetes. The Wleń Graben is a narrow structure, c. 17.5 km long and up to 3.5 km wide, downthrown  
into the low-grade metamorphics of the Kaczawa Slate Belt, filled with Permian, Triassic and Upper Cretaceous (up to lower  
Coniacian) mostly sedimentary, shallow marine or continental deposits and “bounded by steep, NW–SE-oriented, normal  
and reverse faults” (Kowalski, 2020; Fig. 11). The graben was earlier studied by Gorczyca-Skała (1977) and discussed by  
Solecki (1994, 2011), who stressed the particular significance of NNE-SSW to NE-SW directed compression in its  
205 formation. Kowalski (2020) advocates a multistage evolution of the graben, from Late Cretaceous times (post-Santonian?)  
onwards, including the first, extensional stage of its formation with active NW-SE striking normal faults, followed by  
compressional event of “probably latest Cretaceous to early Palaeogene(?)” age, re-activating and inverting the ~~latter~~  
discontinuities and producing new reverse faults and overthrusts, particularly in the central and southern parts of the graben.  
This was postdated by minor sinistral displacements of the graben fragments along cross-cutting faults of NE-SE trend, and  
210 the evolution is believed by this author to have ended with an extensional episode of limited significance, presumably  
already during the mid- to late Cenozoic.

The cross-sections of the Wleń Graben prepared by Kowalski (2020) and based on his mapping fieldwork and scarce  
borehole data (Fig. 11) show a rather regular syncline, only slightly modified by the graben’s boundary faults and locally  
complicated by faults striking obliquely to the syncline’s axis (Fig. 11). The metamorphic basement top is depicted as  
215 concordantly ~~draped around~~ the overlying, synclinally bent base Rotliegend surface. The latter solution may prove  
~~problematic~~ and the cited author suggests a seismic survey to be made in order to better understand the structural geometry  
of the graben.

#### 4.4. The Intra-Sudetic Synclinorium

The Intra-Sudetic Synclinorium is another, apart from the North Sudetic one, extensive Late Cretaceous – Early Palaeogene  
220 tectonic structure of NW-SE trend in the Sudetes (Figs. 2, 3 and 11). It has a comparable, though somewhat larger size of c.  
80 km in length and up to 30 km in width. It, similarly, affects the post-Variscan continental to shallow marine Permo-



Mesozoic succession, which represents the upper structural level of the Intra-Sudetic Basin (Żelaźniewicz and Aleksandrowski, 2008; Żelaźniewicz et al., 2011), unconformably deposited on top of a thick succession of intramontane Carboniferous syn- to late-orogenic clastics deformed by Variscan tectonism, corresponding to the lower structural level of the Intra-Sudetic Basin (see also: Nemeček et al., 1982; Dziedzić and Teisseyre, 1990; Bossowski et al., 1995; McCann, 2008b). The Carboniferous deposits of the Intra-Sudetic Basin rest on the Variscan-deformed, mostly crystalline basement of the Sudetic Block. To the SE, the axial zone of the Intra-Sudetic Synclinorium, composed of Late Cretaceous deposits, merges the Upper Nysa-Králiky Graben (Figs. 2 and 12). Another similarity of the Intra-Sudetic to the North Sudetic Synclinorium is its generally simple, open structure of a single syncline in its NW and central parts, which is replaced by a number of (partly reverse) synclinal grabens separated by basement horsts on its north-eastern flank (Figs. 2 and 12).

The Late Cretaceous to Early Palaeogene tectonic structures of the Intra-Sudetic Synclinorium studied by us on recently reprocessed legacy seismic sections, come only from the NE limb of the synclinorium, located between Mieroszów in the NW and Ścinawka in the SE. In that area it is the Permian rocks that are exposed at the surface, while the Cretaceous succession of the Intra-Sudetic Synclinorium axial zone occurs to the SW and is not covered by the seismics. The seismic sections (location in Fig. 12) are mostly trending SW-NE, that is roughly perpendicular to the synclinorium's axis. They are consecutively presented, from NW to SE, in figures 13 and 14, in order to illustrate in a systematic way the tectonic deformation structures affecting the Permian and Carboniferous strata.

In the NW part of the area covered with seismic data (Figs. 12 and 13), three consecutive profiles show a bunch of faults in Carboniferous clastics, which does not continue upwards into the Permian, although in some profiles (Figs. 13 b and c) these faults manifest themselves also in the Permian strata as gentle ductile folds above the faults. These folds are geometrically entirely consistent with fault-related folds in the Carboniferous. The most likely interpretation of this case seems to be a post-Early Permian reverse-slip activity of faults in the Carboniferous succession (probably representing a reactivated Carboniferous flower structure), associated with gentle folding that locally affected also the overlying Permian strata, though locally were fully accommodated still at the level below the Permian (Fig. 13a). The fault zone in the Carboniferous succession can be followed on seismics at a distance of 7 km, but probably continues further on both to the NW and SE beyond the seismically explored area.

Further to the SE, near Broumov in Czechia (Figs. 12 and 14a,b), a low-angle, NE-vergent thrust zone (which may represent a much evolved continuation of the fault zone from Fig. 13) rooted in the crystalline basement, produces a reverse-slip displacement of the Carboniferous strata by at least 2 km with respect to the footwall, associated with the formation of fault-bend folds. The thrust seems to continue to the NE as a décollement within the Permian strata, overlain by a pair of anticline-syncline, which can be recognized in the surficial geological map outcrop pattern.

Moving further to the SE along the NE flank of the Intra-Sudetic Synclinorium axial zone, in the area of Radków, on the successive seismic profiles one can follow along-strike geometrical changes of a conspicuous low-angle, NE-vergent thrust fault, responsible for an up to 2.5 km magnitude of displacement of the crystalline basement and the overlying Carboniferous



255 and Permian over their footwall. The overthrust is related – ~~similarly as in~~ the case at Broumov – with an anticline-syncline pair of ~~the~~ fault-bend or fault-propagation(?) folds.

The conspicuous thrust fault identified at depth in the area of Radków – Ścinawka (Figs. 14 c-f) and – most probably – continuing into the vicinities of Broumov (Fig. 14 a,b), seems to correspond to the major NW-SE trending Ścinawka-Krosnowice Fault (Fig. 12) exposed at the surface. The latter fault has been, however, so far consistently interpreted as a NE-  
260 throwing normal (Grocholski and Augustyniak, 1971) or vertical fault (Bossowski and Ihnatowicz, 2006). Therefore, our interpretation of the reprocessed seismic data requires a thoroughgoing change in understanding the nature of this well-known mapped fault at least in its NW, Ścinawka segment, as a low angle Late Cretaceous – Early Palaeogene thrust. In the light of the geological and tectonic context, there can be little doubt about the ~~latter~~ timing of the tectonic structures identified by us on the seismic sections. ~~This~~ is corroborated by their occurrence near ~~to~~ the axial zone of the Intra-Sudetic  
265 Synclinorium with abundant presence of ~~the~~ preserved Upper Cretaceous deposits. ~~This~~ is also confirmed by the SE-ward surficial extension of the Ścinawka-Krosnowice Fault, which at its Krosnowice segment defines a tectonic boundary between the Permo-Mesozoic succession of the Intra-Sudetic Basin and Nysa-Králíky Graben on one side and the Śnieżnik – Krowiarki crystalline Massif on the other side.

An interesting Late Cretaceous – Early Palaeogene structure at the NE periphery of the Intra-Sudetic Synclinorium is the  
270 NW-SE trending Czerwieńczyce Reverse Graben (Figs. 12 and 15). According to Oberc (1972, 1977), it is bounded by high-angle reverse faults (and, hence, we apply to it the term “reverse graben”) and filled with Rotliegend and Upper Carboniferous sedimentary rocks (Oberc, 1972, 1977). On approaching the graben’s boundary faults, the strata are strongly dragged upwards, which renders a synclinal geometry to the graben’s fill. This ‘synclinal’ structural characteristics of the Czerwieńczyce Reverse Graben – if correct, being based on surficial geological mapping and involving much extrapolation  
275 at depth – is, nevertheless, quite symptomatic of a number of Sudetic grabens and to some degree it resembles cross-sections of e.g. the Wleń Graben by Kowalski (2020; Fig. 11) and – still more – ~~that~~ by Gorczyca-Skała (1977, not reproduced here), or of the Upper Nysa – Králíky Graben by e.g. Jerzykiewicz (1970) and Don and Gotowała (2008) – partly with reverse boundary faults.

#### 4.5. The Upper Nysa - Králíky Graben

280 The Upper Nysa Nysa – Králíky Graben (Fig. 16) is a distinctive tectonic and topographic, fault-bounded feature of ~~nearly~~ N-S trend, **c.45** km long and from 3 to 12 km wide, merging to the north with the Intra-Sudetic Synclinorium, and **filled with** shallow marine sedimentary rocks, including a time span from the early Cenomanian, through Turonian to Coniacian and Santonian (e.g. Don and Don, 1960; Jerzykiewicz, 1970, 1971; Radwański, 1975; Don and Gotowała, 2008; Badura and Rauch, 2014). Most of the earlier authors (except Jerzykiewicz, 1970, 1971; Radwański, 1975 and Oberc 1972) explained  
285 the origin of the Upper Nysa – Králíky Graben as a result of a Late Cretaceous rifting modified later by Cenozoic subsidence (e.g. Don and Don, 1960; Don, 1996; Wojewoda, 1997; Don and Gotowała, 2008). The early rifting was inferred to have



occurred on stratigraphic and sedimentological premises – not convincing in our opinion – and most often explained as due to compression- or extension-driven updoming of the Orlica-Śnieżnik Massif during the Cretaceous or, still, by pull-apart graben formation due to strike-slip displacements on NW-SE trending structural discontinuities in the crystalline basement.

290 In contrast, we, in this paper, interpret the mostly N-S trending Upper Nysa – Králiky Graben to represent a Late Cenozoic feature, whose boundary faults cut out and downthrow a strip of the Cretaceous (Cenomanian, Turonian, Coniacian through to Santonian) shallow marine succession previously much more widespread over the area. The Cretaceous is now preserved within the graben due to its being downthrown against the uplifted crystalline basement around. The uniqueness of the Cretaceous deposits in the Upper Nysa – Králiky Graben consists only in the preservation of the thick Coniacian to

295 Santonian succession, absent from the uplifted areas nearby, from where it must have clearly been eroded. The Upper Nysa – Králiky Graben is imposed on and cuts across earlier, Late Cretaceous – Early Palaeogene, tectonic structures of ~~stable~~ NW-SE strike, which are in most places oblique to and cross-cut by the graben's boundaries (Fig. 16). These structures are represented by differentially dip-slip displaced fault blocks, often bounded by reverse faults, and by gentle folds (Badura & Rauch, 2014; Don and Gotowała, 2008).

300 Locally, shallow-dipping thrust faults, most probably of Late Cretaceous–Early Palaeogene age were mapped, drilled or directly observed in outcrops within or in direct vicinities of the Upper Nysa – Králiky Graben. A relatively extensive, at least 25 km-long, NW-SE trending trace of the Zieleniec Thrust Fault was mapped by Cymerman (1990) and extrapolated on maps by other authors within the SW flank of the Upper Nysa – Králiky Graben. On this c. 40-50° SW-dipping fault a 500 Ma orthogneiss mass from the Góry Bystrzyckie Mts was overthrust, displaced by at least 5300 m toward the NE

305 (Cymerman, 1990) and emplaced on top of the Turonian clastics, equivalent to those from the lower part of the Upper Nysa - Králiky Graben fill (Fig. 17) and resting themselves on analogous orthogneiss basement. Cymerman (1990) ~~has~~ interpreted this thrust as produced by Late Alpine (Miocene) compression. The Zieleniec Thrust was later confirmed by drilling, as reported by Kozdrój (2014), who interpreted the thrusting as the result of a Cenozoic gigantic, gravity-driven landslide, at the same time referring to inversion of an original normal fault into a reverse one. One of most important arguments for this

310 ~~judgement~~ was finding a subhorizontal position of the Cretaceous/gneiss contact in the drillcore. In our opinion, taking into account its significant extent, ~~and~~ the involvement in the major NW-SE trending fault system (Fig. 12), termed the South-Sudetic Shear Zone by Wojewoda (e.g. in Wojewoda and Kowalski, 2016), ~~involving~~ the major Pořici-Hronov Fault (see below), there cannot be much doubt that the Zieleniec thrust represents a Late Cretaceous – Early Palaeogene thrust, similar to that of the Łużyce Thrust (see below), though of ~~an~~ opposite vergence.

315 To the NW the Zieleniec Thrust ~~merges~~ the ~~SE-extension~~ of the Pořici-Hronov major fault system (Fig. 12), partly representing the SW boundary to the Intra-Sudetic Synclinorium. It was active, as well, in the Late Cretaceous - Early Palaeogene as a high-angle reverse fault of complex structure, but of ~~the~~ opposite, SW, polarity (Prouza et al., 2014), which – according to some authors - is combined with the strike-slip component of motion (Wojewoda, 2007, 2009; Nováková, 2014) (Fig. 18a). To the SE, the Pořici-Hronov fault continues into ~~that of~~ Duszniki-Gorzanów/~~Krosnowice~~, at the northern

320 end of the Upper Nysa – Králiky Graben.



A presumable splay fault, branching off from the Zieleniec Thrust (Figs. 12 and 18b) was described directly in a c. 1 km-long chain of outcrops along a road cut between Duszniki Zdrój and Lewin Kłodzki near the pass of Polskie Wrota on the same SW flank of the Upper Nysa - Králiky Graben (Wojewoda et al., 2010; Aleksandrowski and Wojewoda, 2010). A c. 20° NW-dipping detachment can be seen to occur there roughly along the bedding in Middle Turonian mudstones and calcareous claystone. On a closer inspection, the detachment was found to represent a shear zone, attaining 7 to 25 cm in thickness and composed of interconnected, kinematically linked surfaces, confined between two distinct boundaries of mostly intact rock (Fig. 18b). The shear zone shows internal flaser bedding and is composed of cataclastic flow products of host-rock composition, which coexist with crush breccia and calcite veins. On evidence from the orientation of the accompanying joints, the detachment was then interpreted as SE-vergent and probably genetically related to a supposed strike slip motion on the nearby major NW-SE-trending Duszniki-Gorzanów Fault, being a SE continuation of the Poříčí-Hronov fault system (cf. Petrascheck, 1933; Prouza et al., 2014). It seems likely, however, that the joints can represent the youngest, Miocene, deformation and episode of motion on the detachment in question, whereas, ~~the~~ very origin of ~~it~~ and principal displacement ~~was~~ related to ~~that on~~ the Zieleniec Thrust.

#### 4.6. The south-western margin of the Sudetes: transition to the North Bohemian Cretaceous Basin

The NW segment of the Sudetic Block, represented by the Karkonosze-Izera Massif, to the SW adjoins the North Bohemian Cretaceous Basin (Fig. 2). At its Sudetic margin, the basin is composed of deltaic to shallow-marine and hemipelagic Cenomanian to Santonian clastic deposits, not exceeding 600 to 700 m in thickness that rest subhorizontally on scarce subcrops of Permian and Carboniferous, but mostly on the crystalline Variscan basement of the Bohemian Massif (e.g. Klein and Soukup, 1966; Malkovský, 1987; Uličný et al., 2009; Wilmsen et al., 2014). Its interior is only little affected by the Late Cretaceous – Early Palaeogene tectonism, however at its boundary with the Sudetic block the deformation is concentrated at the spectacular and at places well exposed Lusatian Thrust (Fig. 2 and 19 a-c), which continues NW up to the vicinities of Dresden (e.g. Wagenbreth, 1967), whereas to the east it continues to the western Karkonosze Piedmont area (Prouza et al., 2013). The Lusatian Thrust has been excellently described and analysed in detail by Coubal et al. (2014).

The Lusatian Thrust (e.g. Malkovský, 1976), similarly to other main NW-SE trending fault zones of the Sudetic area at the Bohemian NE margin (the Intra-Sudetic Fault, the Sudetic Boundary Fault, the Odra Fault Zone) derives its origin from a major Variscan fracture, rooted deeply in the basement and likely having a primary strike-slip characteristics (cf. Aleksandrowski et al., 1997). It must have been reactivated during the Late Cretaceous – Early Palaeogene compressional episode and propagated into the Permo-Mesozoic cover. The Lusatian Thrust emplaces the Variscan crystalline complexes of Lusatia and of the West Sudetes on top of the Mesozoic strata of the North Bohemian Cretaceous Basin. This fault zone reveals significant along-strike changes in the dip angle of its displacement surface, between low-angle or subhorizontal in the northwest (Figs. 2 and 19a) through medium-angle (Figs. 2 and 19b) and high-angle attitude (Figs. 2 and 19c). The angle made by the displacement zone/fault core with respect to the – supposedly generally subhorizontal - tectonic shortening



direction had serious influence on the structural style and complexity of the brittle structures developed around the fault core and on the width of the damage zone (Coubal et al., 2014). **The smaller the angle between the two,** the more smoothly the displacement occurred (Fig. 17a). An increase of the fault plane dip angle made the displacement more difficult to be achieved, which resulted in widening a zone of damage and of fault drag and favoured splitting the displacement into several slip surfaces (Fig. 19b and c).

### **5 Regional jointing pattern: a likely product of Late Cretaceous – Early Palaeogene deformation**

The regional pattern of tectonic joints over the area of the Sudetes or of SW Poland can be inferred from a number of local studies on jointing, most of which were conducted decades ago. Not attempting to make a systematic review of such studies, for the needs of a brief review made in this paper, several stereograms or rose diagrams of the dominant joint sets coming from various sources (Jerzykiewicz et al., 1974; Aleksandrowski, 1976; Solecki, 2011; Selerowicz et al., 2014) and various locations in the Polish Sudetes have been mounted on the geological map (Fig. 20). The data come mostly from the Permo-Mesozoic strata exposed in outcrops in the both Sudetic synclinoria, in underground galleries of the deep copper mines at the Fore-Sudetic Homocline and in the Opole Cretaceous Basin. Two stereograms, show, additionally the joint pattern in the crystalline basement rocks of the Karkonosze Massif.

All the presented diagrams show most steep joints to be concentrated in two mutually perpendicular maxima, in which the joint planes are approximately parallel and perpendicular, respectively, to the inferred Late Cretaceous-Early Palaeogene shortening direction (cf. e.g. Solecki, 2011, Nováková, 2014). As, moreover, they occur in the Permo-Mesozoic strata, a conclusion that they are genetically related to that shortening event seems reasonable. Actually, we believe that the formation of dominant jointing pattern in the post-Variscan (but also partly in the Variscan) Sudetes and the SW part of Poland resulted from the Late Cretaceous-Early Palaeogene compressional event.

### **6 Conclusion**

In our brief review of the Late Cretaceous – Early Palaeogene tectonic structures affecting the NE margin of the Bohemian Massif we have shown their common and widespread occurrence all over the region and, to some degree, shortly discussed their style and mechanisms of formation. These structures are easily recognizable in the Permo-Mesozoic post-Variscan cover as devoid of other, earlier formed contractional deformations, but, though present, they are much less obvious in the Variscan basement that was heavily tectonized before. Those developed in the post-Variscan cover, particularly the fault-displacement effects – and, among them, the strike-slip related ones - can still be difficult to be separated from the structures that might have resulted from the younger Cenozoic events.

Our structural analysis of newly reprocessed legacy seismic profiles, complemented with outcrop, drillhole data, have revealed an important contribution of thin-skinned fold-and-thrust type contractional deformation in the Permo-Mesozoic



(partly also in the Carboniferous) strata of the two main Sudetic synclinoria. Such deformation includes also gentle buckling of the basement/cover interface and decoupling/decollement of the Permo-Mesozoic cover along **weak, clayey horizons in Zechstein strata**. The low-angle thrust faults identified in the Permo-Mesozoic in the seismic record are often rooted in the top parts of the Variscan basement, so some elements of the thick-skinned style of shortening can be also involved (and, actually, are to be expected in the deep crystalline basement). It is, therefore, likely that also outside the areas covered with Permo-Mesozoic sedimentary sequences, over the crystalline basement exposure areas, some (even major) faults interpreted from surficial geological maps and outcrop relationships as vertical or high-angle structures, may have at depth a geometry of low-angle thrusts. This may have important practical implications on e.g. geothermal prospecting in the Sudetes, where the deep underground waters in the crystalline basement circulate, almost exclusively, along structural discontinuities, such as fault zones and fracture corridors (e.g. Dowgiałło, 2002).

The Late Cretaceous – Early Palaeogene compressional event seems to be also responsible for the formation of the regionally dominant tectonic joints pattern in both Permo-Mesozoic and older rocks of the Sudetic area, characterized by an **orthogonal** joint system with two sub-vertical sets of c. NW-SE and NE-SW strikes, as is shown by a comparison of results from a few local studies.

#### **Data availability**

The seismic data from Poland are available at the National Geological Archive and the seismic profile ~~from~~ Broumov area is available at the Czech Geological Survey Archive.

#### **Author contributions**

PA and AG wrote the text, prepared the figures, and compiled the paper.

#### **Competing interests**

The authors declare they have no conflict of interest.

#### **Acknowledgements.**

The consent of the Polish Oil and Gas Company (PGNiG) to publish their seismic data is gratefully acknowledged. Similarly, Janusz Badura, Aleksander Kowalski and Marta Rauch, together with the geological institutes of the Polish Academy of Sciences and of University of Wrocław as well as the Polish Geological Society and the Polish Geological Institute – National Research Institute are thanked for the permission to reproduce their illustrative materials.

#### **Financial support:**

This work was supported by Polish Geological Institute grant no 62.9012.2046.00.0. The reprocessing of the key reflection seismic data from the North- and Intrasudetic synclinoria was commissioned by the Polish Geological Institute as part of its



statutory duties, financed by the National Fund of Environmental Protection and Water Management in Warsaw under contract no 44/2017/Wn-07/FG-SM-DN/D and performed by GK Processing Sp. z o.o. in Cracow.

## References

- 420 Aleksandrowski P.: Geological mapping survey of the Karkonosze main range between Mt. Śnieżka and Mt. Skalny Stół. Unpubl. MSc thesis, University of Wrocław (in Polish), 1976.
- Aleksandrowski P., Kryza, R., Mazur, S. and Żaba J.: Kinematic data on major Variscan fault and shear zones in the Polish Sudetes, NE Bohemian Massif, *Geological Magazine*, 134, 727-39, 1997.
- Aleksandrowski P. and Mazur S.: Collage tectonics in the northeasternmost part of the Variscan Belt: the Sudetes, Bohemian Massif. In: Winchester J.A., Pharaoh T.C. and Verniers J. (eds): *Palaeozoic Amalgamation of Central Europe*, vol 201. Geological Society, Special Publications, London, 237–277. <https://doi.org/10.1144/GSL.SP.2002.201.01.12>, 2002.
- 425 Aleksandrowski, P. and Wojewoda, J.: Low-angle detachment related to strike-slip faulting in Late Cretaceous mudstones of the Table Mountains (SW Poland). In: 11th Czech-Polish Workshop On Recent Geodynamics of the Sudetes and Adjacent Areas, Třešť, November 4-6th, 2010, Academy of Sciences of the Czech Republic (Institute of Rock Mechanics) and Wrocław University of Environmental Sciences (Institute of Geodesy and Geoinformatics), Abstracts, 30-31, 2010.
- 430 Badura, J. and Rauch, M.: Tectonics of the Upper Nysa Kłodzka Graben. *Geologia Sudetica*, 42, 137-148, 2014.
- Bałażyńska, J. and Bossowski, A.: Wgłębna budowa geologiczna środkowej i zachodniej części synklinorium północnosudeckiego w świetle nowych danych (Deep geological structure of central and western parts of the North-Sudetic Synclinorium; some new data), *Kwartalnik Geologiczny*, 23, 2, 309-321, 1979.
- 435 Beyer, K.: Die nordsudetische Rahmenfaltung, *Abh. Naturforsch. Ges. Górlitz*, 32, 121-172, 1939.
- Bossowski, A. and Bałażyńska, J.: Ewolucja tektoniczno-strukturalna synklinorium północnosudeckiego (Tectonic-structural evolution of the North-Sudetic Synclinorium), *Biuletyn Instytutu Geologicznego*, 341, 163-167, 1982.
- Bossowski, A. and Ichnatowicz, A.: *Atlas geologiczny Dolnośląskiego Zagłębia Węglowego* (Geological atlas of the Lower Silesian coal Basin), 1 : 100 000, Państwowy Instytut Geologiczny, Warszawa, 2006.
- 440 Bossowski, A., Ichnatowicz, A., Mastalerz, K., Kurowski, L. and Nowak, G.: Lithostratigraphy and sedimentologic-paleogeographic development, Intra-Sudetic Depression. In: Zdanowski A. and Żakowa H. (eds), *The Carboniferous system in Poland*, *Prace Państwowego Instytutu Geologicznego* 148, 142-147, 1995.
- Brada, Z., Fejfar, M., Ibrmajer, I., Jihlavcová, R., Petřík, A.: Study of coal-bearing formations in the Bohemian Massif - Partial report on seismic reflection survey of the Broumov promontory. *Geofyzika Brno*, (Unpublished report in Czech language; Archive, Czech Geological Survey), 1982.
- 445 Cloos, H.: *Der Gebirgsbau Schlesiens und die Stellung seiner Bodenschätze*. Gebrüder Borntraeger, Berlin, 1-107, 1922.



- Coubal, M., Adamovič, J., Málek, J., and Prouza, V.: Architecture of thrust faults with alongstrike variations in fault-plane dip: anatomy of the Lusatian Fault, Bohemian Massif, *Journal of Geosciences*, 59, 183–208, DOI: 10.3190/jgeosci.174, 2014.
- 450 Coubal, M., Malek, J., Adamovič, J. and Štěpančíková, P.: Late Cretaceous and Cenozoic dynamics of the Bohemian Massif inferred from the paleostress history of the Lusatian Fault Belt, *Journal of Geodynamics*, 87. 10.1016/j.jog.2015.02.006, 2015.
- Cwojdziański, S. and Żelaźniewicz, A.: Podłoże krystaliczne bloku przedsudeckiego (Crystalline basement of the Fore-Sudetic block). In: *Annales Societatis Geologorum Poloniae, Special Volume: Przewodnik 66 Zjazdu Polskiego Towarzystwa Geologicznego (Guide to excursions, 66th Annual Meeting of the Polish Geological Society)*, Wrocław 21-24
- 455 September 1995, 11-28 (in Polish with English summary), 1995.
- Cymerman, Z.: Młodo-alpejskie nasunięcie Zieleniec w Górach Orlickich (Late-Alpine Zieleniec overthrust in the Orlickie Mts), *Przegląd Geologiczny*, 10, 422–427, 1990.
- Cymerman, Z.: Młodoalpejskie fałdy w depresji północnosudeckiej: przykłady z wapienia muszlowego z Raciborowic,
- 460 *Przegląd Geologiczny*, 4, 348-354, 1998.
- Cymerman, Z.: Alpejska transpresja w Sudetach (Alpine transpression in the Sudety Mts), *Przegląd Geologiczny*, 47, 942–945, 1999.
- Cymerman, Z.: Tectonic map of the Sudetes and the Fore-Sudetic Block. 1 : 200 000, 2nd Ed. Państwowy Instytut Geologiczny, Warszawa, 2010.
- 465 Don, J.: The Late Cretaceous Nysa Graben: implications for Mesozoic – Cenozoic fault – block tectonics of the Sudetes, *Zeitschrift für geologische Wissenschaften*, 24, 317–424, 1996.
- Don, J. and Gotowała, R.: Tectonic evolution of the late Cretaceous Nysa Kłodzka Graben, Sudetes, SW Poland, *Geologia Sudetica*, 40, 51–63, 2008.
- Don, B. and Don, J.: Geneza rowu Nysy na tle badań wykonanych w okolicach Idzikowa. [Notes on the origin of the Nysa
- 470 graben], *Acta Geologica Polonica*, 10, 71–106 (in Polish), 1960.
- Dowgiałło J.: The Sudetic geothermal region of Poland, *Geothermics*, 31: 343-359, 2002.
- Dumicz, M. and Don, J.: Analiza strukturalna monokliny przedsudeckiej w rejonie Polkowic (Structural analysis of the Fore-Sudetic Homocline in vicinities of Polkowice), *Acta Universitatis Wratislaviensis* 378, *Prace Geologiczno-Mineralogiczne*, 6, 279-297, 1977.
- 475 Dumicz, M. and Don, J., Próba odtworzenia następstwa zjawisk tektonicznych w osadach cechsztynu obszaru Polkowic na podstawie obserwacji drobnych struktur tektonicznych (An attempt at reconstruction of the sequence of tectonic phenomena in the Zechstein sediments at the Polskowice area on the basis of study on minor tectonic structures). In: *Materiały konferencji komisji tektoniki Komitetu Nauk Geologicznych PAN: Problemy tektoniki Legnicko-Głogowskiego Okręgu Miedziowego, część II – wycieczki geologiczne*, Lubin, Wydawnictwo CUPRUM, Wrocław, 39-45, (in Polish), 1990.



- 480 Dziedzic, K. and Teisseyre, A. K.,: The Hercynian molasse and younger deposits in the Intra-Sudetic Depression, SW Poland, *Neues Jahrb. Geol. Paläont. ABH*, 179, 285–305, 1990.
- Głuszyński, A., and Smajdor, Ł.: Analiza archiwalnych danych głębokiej sejsmiki poszukiwawczej na obszarze synklinoriów sudeckich (Analysis of legacy prospecting seismic data from the Sudetic synclinoria). In: Aleksandrowski P. (ed): Unpubl. report of Polish Geological Survey project *Młode strefy tektoniczne a warunki geotermalne w Sudetach w*  
485 *świecie badań geochronologicznych, strukturalnych i termometrycznych – etap II*, 259-269 (in Polish), National Geological Archive, Warszawa, 2020.
- Gorczyca-Skała, J.: Budowa geologiczna rowu Wlenia (Geological structure of the Wleń Graben), *Geologia Sudetica*, 12, 1, 71–100, 1977.
- Grocholski, A. and Augustyniak, K.: Atlas geologiczny Dolnośląskiego Zagłębia Węglowego (Geological atlas of the Lower  
490 Silesian coal Basin), 1 : 50 000. Pt. I, Instytut Geologiczny, Warszawa, 1971.
- Jarosiński, M., Poprawa, P., Ziegler, P. A.: Cenozoic dynamic evolution of the Polish Platform. *Geological Quarterly*, 53 (1), 3–26, 2009. Jerzykiewicz, T.: The Upper Cretaceous turbidite sequence in the Sudetes (South-western Poland), *Bulletin de l'Academie Polonaise des Sciences, Série des scis géol. et géogr.*, 18, 3, 149-159, 1970.
- Jerzykiewicz, T.: A flysch/lithoral succession in the Sudetic Upper Cretaceous. *Acta Geologia Polonica*, 21, 165–199, 1971.
- 495 Jerzykiewicz, T., Mierzejewski, M. and Żelaźniewicz, A.: Joint and Fracture Patterns in Basement and Sedimentary Rocks in the Sudetes Mountains, Proceedings of the First International Conference on the New Basement Tectonics (Utah Geol. Assoc. Publication no 5), 295-306, [https://archives.datapages.com/data/uga/data/091/091001/295\\_ugs0910295.htm](https://archives.datapages.com/data/uga/data/091/091001/295_ugs0910295.htm), 1974.
- Klein, V. and Soukup, J.: The Bohemian Cretaceous Basin. In: Svoboda J. et al. (Ed), *Regional Geology of Czechoslovakia, Pt 1, The Bohemian Massif*, Geological Survey of Czechoslovakia, Prague, 487-512, 1966.
- 500 Kley J.: Saxonische Tektonik im 21. Jahrhundert. *Z. Dt. Ges. Geowiss. (German J. Geosci.)*, 164 (2), 295–311, 2013.
- Kley, J.: Timing and spatial patterns of Cretaceous and Cenozoic inversion in the Southern Permian Basin, in: *Mesozoic Resource Potential of the Southern Permian Basin*, edited by: Kilhams, B., Kukla, P. A., Mazur, S., McKie, T., Munlieff, H. F., and van Ojik, K., Geological Society, London, Special Publications, 469, 19–31, <https://doi.org/10.1144/SP469.12>, 2018.
- Kley, J. and Voigt, T.: Late Cretaceous intraplate thrusting in central Europe: effect of Africa-Iberia-Europe convergence,  
505 not Alpine collision, *Geology*, 36, 839–842, 2008.
- Kotański, Z. (ed): *Geological Atlas of Poland, Geological maps of horizontal section, 1 : 750 000*, Państwowy Instytut Geologiczny, Warszawa, 1997.
- Kowalski, A.: Fault geometry and evidence of depocentre migration within a transtensional intra-basinal high : a case study from the Łączna Anticline (Intrasudetic Synclinorium, SW Poland) *Geological Quarterly*, 2017, 61 (4): 779–794, 2017.
- 510 Kowalski, A.: Multistage structural evolution of the end-Cretaceous–Cenozoic Wleń Graben (the Sudetes, NE Bohemian Massif) – a contribution to the post-Variscan tectonic history of SW Poland, *Annales Societatis Geologorum Poloniae*, 90, 10.14241/asgp.2020.21, 2020.



- Kozdrój, W.: Results of shallow scientific drillings in the Upper Nysa Kłodzka Graben and the Zieleniec area, Sudetes, *Geologia Sudetica*, 42, 149–159, 2014.
- 515 Kroner, U., Mansy, J.-L., Mazur, S., Aleksandrowski, P., Hann, H.P., Huckriede, F., Lacquement, F., Lamarche, J., Ledru, P., Pharaoh, T.C., Zedler, H., Zeh, A., Zulauf, G.: Variscan Tectonics. In: McCann, T. (ed.): *The Geology of Central Europe*. The Geological Society, London: 599-664, 2008.
- Krzywiec, P.: Mid-Polish Trough inversion – seismic examples, main mechanisms and its relationship to the Alpine-Carpathian collision, *Stephan Mueller Special Publication Series*, 1, 151–165, <https://doi.org/10.5194/smsps-1-151>, 2002.
- 520 Krzywiec, P. and Stachowska, A.: Late Cretaceous inversion of the NW segment of the Mid-Polish Trough – how marginal troughs were formed, and does it matter at all?, *Zeitschrift der deutschen geologischen Gesellschaft*, 167, 107–119, 2016.
- Krzywiec, P., Peryt, T., Kiersnowski, H., Pomianowski, P., Czapowski, G. and Kwolek, K.: Permo-Triassic evaporites of the Polish Basin and their bearing on the tectonic evolution and hydrocarbon system, an overview, Chapter 11 in Soto, J.I., Flinch, J., Tari, G. (Eds) - *Permo-Triassic salt provinces of Europe, North Africa and the Atlantic margins: tectonics and hydrocarbon potential*, 243-261, Elsevier, <http://dx.doi.org/10.1016/B978-0-12-809417-4.00012-4>, 2017.
- 525 Leśniak, T.: Tektonika obszaru między Raciborowicami a Łaziskami w północno-wschodniej części depresji północnosudeckiej, *Zeszyty Naukowe AGH, Geologia*, 5, 4, Kraków, 87-107, (in Polish), 1979.
- Malkovský, M.: Saxonische Tektonik der Böhmischen Masse, *Geologische Rundschau*, 65, 10.1007/BF01808459, 1976. Malkovský, M.: Important faults of the platform cover of the northern part of the Bohemian Massif. *Výzk Práce Ústí*
- 530 *Úst geol*, 14, 1–32, 1977.
- Malkovský, M.: The Mesozoic and Tertiary basins of the Bohemian Massif and their evolution, *Tectonophysics*, 137, 31-42, 10.1016/0040-1951(87)90311-8, 1987.
- Malz, A., Nachtweide, Ch., Emmerlich, S., and Schimpf, L.: Mesozoic intraplate deformation in the southern part of the Central European Basin - Results from large-scale 3D modelling, *Tectonophysics*, 776, 10.1016/j.tecto.2019.228315, 2020.
- 535 Markiewicz, A.: Tektonika obszaru złoża (The tectonics of the ore-deposit area), Chapter 2.11 in: Piętrzyński A., (ed.), *Monografia KGHM Polska Miedź S.A., KGHM CUPRUM, Wrocław*, 115-132, 2007.
- Markiewicz, A., Szarowski, W.: Zjawiska tektoniczne w południowej części kopalni Lubin, In: *Materiały konferencji komisji tektoniki Komitetu Nauk Geologicznych PAN: Problemy tektoniki Legnicko-Głogowskiego Okręgu Miedziowego, część II – wycieczki geologiczne*, Lubin, Wydawnictwo CUPRUM, Wrocław, 4-8, 1990.
- 540 Martínez Catalán J.R., Collett S., Schulmann K., Aleksandrowski P. and Mazur S.: Correlation of allochthonous terranes and major tectonostratigraphic domains between NW Iberia and the Bohemian Massif, European Variscan belt, *International Journal of Earth Sciences (Geol Rundsch)*, 109, 4, 1105-1131, <https://doi.org/10.1007/s00531-019-01800-z>, 2020.
- Mazur, S., Aleksandrowski, P., Kryza, R. and Oberc-Dziedzic, T.: The Variscan orogen in Poland, *Geological Quarterly*, 50, 89-118, 2006.
- 545 Mazur, S., Aleksandrowski, P., Turniak, K. and Awdankiewicz, M.: Geology, tectonic evolution and Late Palaeozoic magmatism of Sudetes – an overview, In: A. Kozłowski and J. Wiszniewska (eds), *Granitoids in Poland*, AM (Archivum



- Mineralogiae) Monograph No. 1, Komitet Nauk Mineralogicznych PAN & Wydział Geologii Uniwersytetu Warszawskiego, 59-87, 2007.
- Mazur, S., Aleksandrowski, P., Turniak, K., Krzemiński, L., Mastalerz, K., Górecka-Nowak, A., Kurowski, L., Krzywiec, P.,  
550 Żelaźniewicz, A. and Fanning, M.C.: Uplift and late orogenic deformation of the Central European Variscan belt as revealed  
by sediment provenance and structural record in the Carboniferous foreland basin of western Poland, *International Journal  
of Earth Sciences*, 99, 47-64, 2010.
- Mazur, S., Aleksandrowski, P., Gaęła, Ł., Krzywiec P., Żaba J., Gaidzik K. and Sikora R.: Late Palaeozoic strike-slip  
tectonics versus oroclinal bending at the SW outskirts of Baltica: case of the Variscan belt's eastern end in Poland,  
555 *International Journal of Earth Sciences*, 109, 1133-1160, 2020.
- McCann, T., Kiersnowski, H., Krainer, K., Vozárová, A., Peryt, T.M., Oplustil, S., Stollhofen, H., Schneider, J., Wetzel, A.,  
Boulvain, F., Dugar, M., Török, Á., Haas, J., Tait, J. and Körner, F.: Permian. In McCann T. (ed.): *The Geology of Central  
Europe*, The Geological Society, London, 531-597, 2008a.
- Mazur S., Scheck-Wenderoth M. and Krzywiec P.: Different modes of the Late Cretaceous-Early Tertiary inversion in the  
560 North German and Polish basins. *International Journal of Earth Sciences (Geologische Rundschau)*, 94, 782-798, 2005.
- McCann, T., Skompski, S., Poty, E., Dugar, M., Vozárová, A., Schneider, J., Wetzel, A., Krainer, K., Kornpohl, K., Schäfer,  
A., Krings, M., Oplustil, S. and Tait, J.: Carboniferous. In McCann T. (ed.): *The Geology of Central Europe*, The Geological  
Society, London, 411-529, 2008b.
- McClay, K.R.: Glossary of thrust tectonics terms in: McClay, K.R. (Ed), *Thrust tectonics*, Chapman & Hall, London, 419-  
565 433, 1992.
- Migoń, P. and Danišik, M.: Erosional history of the Karkonosze Granite Massif – constraints from adjacent sedimentary  
basins and thermochronology. *Geological Quarterly*, 56, 3, 440–454, doi: 10.7306/gq.1032, 2012.
- Milewicz, J.: The geological structure of the North-Sudetic Depression, *Biuletyn Instytutu Geologicznego*, 227, 5–27, 1968.
- Milewicz, J.: Górna kreda depresji północnosudeckiej, *Acta Universitatis Wratislaviensis, Prace Geologiczno-  
570 Mineralogiczne*, 61, 1-58, 1997.
- Navabpour, P., Malz, A., Kley, J., Sieburg, M., Kasch, N. and Ustaszewski, K.: Intraplate brittle deformation and states of  
paleostress constrained by fault kinematics in the central German platform, *Tectonophysics*, 694, 146–163,  
10.1016/j.tecto.2016.11.033, 2017.
- Nádaskay, R., Žák, J., Sláma, J., Sidorinová, T., and Valečka, J.: Deciphering the Late Paleozoic to Mesozoic  
575 tectonosedimentary evolution of the northern Bohemian Massif from detrital zircon geochronology and heavy mineral  
provenance, *International Journal of Earth Sciences*, 108, 2653–2681, 10.1007/s00531-019-01781-z, 2019.
- Nemec, W., Porębski, S. J. and Teisseyre, A. K.: Explanatory notes to the lithotectonic molasse profile of the Intra-Sudetic  
Basin, Polish part (Sudety Mts, Carboniferous-Permian), *Veröffentlichungen Zentralinstituts für Physik der Erde, Akad.  
der Wissenschaften der DDR*, 66, 267–278, 1982.



- 580 Nováková, L.: Evolution of paleostress fields and brittle deformation in Hronov-Poříčí Fault Zone, Bohemian Massif, *Stud. Geophys. Geodet.*, 58, 269–288, 2014.
- Oberc, J.: Sudety i obszary przyległe, [In:] *Budowa geologiczna Polski*. T. 4, Tektonika, cz. 2. Wydawnictwa Geologiczne, Warszawa, 1972.
- Oberc, J.: The Early Alpine epoch in South-West Poland: The North Sudetic Synclinorium, In: Pożaryski W. (Ed), *Geology of Poland*, Vol.4. Tectonics, Wydawnictwa Geologiczne, Warszawa, 419-424, 1977.
- 585 Oberc, J. and Salski, W.: Fałdy i spękania w skałach dolnocechsztyńskich na obszarze szybu wschodniego kopalni Lubin (Fold and joints in lower Zechstein rocks around the eastern shaft of the Lubin mine), *Kwartalnik Geologiczny*, 3, 519-536, 1968.
- Petrascheck, W.: Der böhmische Anteil der Mittelsudeten und sein Vorland. *Mitteilungen der Geol. Gesellschaft in Wien*, 590 24, 1933.
- Pożaryski, W.: *Mapa geologiczna Polski i krajów ościennych bez utworów kenozoicznych 1 : 1 000 000* (Geological map of Poland and adjacent countries without Cenozoic deposits), Instytut Geologiczny, Warszawa (in Polish), 1979.
- Prouza, V., Coubal, M. and Adamovič, J.: Specifika architektury hronovsko-poříčského zlomu (Specific architecture fo the Hronov –Poříčí Fault), *Zprávy o geologických výzkumech v roce 2014/A – Regionální geologie a stratigrafie, Česká* 595 *geologická služba*, Praha, 13-18, 2014.
- Radwański, S.: Kreda Sudetów Środkowych w świetle wyników nowych otworów wiertniczych [Upper Cretaceous of the central part of the Sudetes in the light of new borehole materials], *Biuletyn Instytutu Geologicznego*, 187, 5–59, 1975.
- Salski, W.: Problemy małej tektoniki w rejonie Lubina (Problems of minor tectonics in vicinities of Lubin). *Rudy i Metale Nieżelazne*, 4, 485-489 (in Polish), 1965.
- 600 Salski, W.: Charakterystyka litologiczna i drobne struktury łupków miedzionośnych monokliny przedsudeckiej (Lithological characteristics and minor structures of copper-bearing shales in Fore-Sudetic Homocline), *Kwartalnik Geologiczny*, 12/4, 855-873 (in Polish), 1968.
- Schroder, B.: Inversion tectonics along the Western margin of the Bohemian Massif, *Tectonophysics*, 137, 93-100, 10.1016/0040-1951(87)90316-7, 1987.
- 605 Sobczyk, A, Danišik, M., Aleksandrowski, P. and Anczkiewicz, A.: Post-Variscan cooling history of the central Western Sudetes (NE Bohemian Massif, Poland) constrained by apatite fission-track and zircon (U-Th)/He thermochronology, *Tectonophysics* 649, 47–57, <http://dx.doi.org/10.1016/j.tecto.2015.02.021>, 2015.
- Sobczyk, A., Sobel, E. R. and Georgieva, V.: Meso–Cenozoic cooling and exhumation history of the Orlica–Śnieżnik Dome (Sudetes, NE Bohemian Massif, Central Europe): Insights from apatite fission–track thermochronometry, *Terra Nova*, 32, 610 122–133, 2020.
- Solecki, A.T.: *Tektonika dysjunktywna i jej wpływ na warunki występowania kopalni w synklinorium północnosudeckim*, PhD thesis, University of Wrocław, 1-152 (in Polish), 1986.



- Solecki, A.: Conjugate cataclastic zones in the sandstones of the North-Sudetic Synclinorium, *Przegląd Geologiczny*, 36, 577–581 [In Polish, with English summary.], 1988.
- 615 Solecki, A.: Tectonics of the North Sudetic Synclinorium, *Acta Universitatis Wratislaviensis No 1618, Prace Geologiczno-Mineralogiczne*, 45, 1-60, University of Wrocław, 1994.
- Solecki, A.: Joints and shears of the N-Sudetic Synclinorium. In: Rossmanith, H-P. (ed), *Mechanics of jointed and faulted rocks*. Balkema, 341-346, 1995.
- Solecki, A.: Rozwój strukturalny epiwaryscyjskiej pokrywy platformowej w obszarze synklinorium północnosudeckiego (Structural development of the epi-variscan cover in the North Sudetic Synclinorium area). In: Żelaźniewicz, A. et al. (eds), *Mezozoik i kenozoik Dolnego Śląska, Polskie Towarzystwo Geologiczne, Wrocław*, 19-36, 2011.
- 620 (Structural development of the epi-variscan cover in the North Sudetic Synclinorium area). In: Żelaźniewicz, A. et al. (eds), *Mezozoik i kenozoik Dolnego Śląska, Polskie Towarzystwo Geologiczne, Wrocław*, 19-36, 2011.
- Teisseyre, H.: Ważniejsze dyslokacje ramowe Sudetów (Major boundary faults of the Sudetes), In: Teisseyre H. (Ed.), *Regionalna geologia Polski, T. III: Sudety, Z.1: Utwory przedtrzciorzędowe*, Polskie Towarzystwo Geologiczne, Kraków, 25-29 (in Polish), 1957.
- 625 Uličný, D., Špičáková, L., Grygar, R., Svobodová, M., Čech, S. and Laurin, J.: Palaeodrainage systems at the basal unconformity of the Bohemian Cretaceous Basin: roles of inherited fault systems and basement lithology during the onset of basin filling, *Bulletin of Geosciences*, 84, 4, 577–610, 2009.
- Voigt, S., Wagreich, M., Surlyk, F., Walaszczyk, I., Uličný, D., Čech, S., Voigt, T., Wiese, F., Wilmsen, M., Niebuhr, B., Reich, M., Funk, H., Michalík, J., Jagt, J.W.M., Felder, P.J. and Schulp, A.S.: Cretaceous, In McCann T. (ed.): *The Geology of Central Europe*, The Geological Society, London, 923-997, 2008.
- 630 Voigt, T., Kley, J. and Voigt, S.: Dawn and dusk of Late Cretaceous basin inversion in central Europe, *Solid Earth*, 12, 1443–1471, <https://doi.org/10.5194/se-12-1443-2021>, 2021.
- Wilmsen, M., Uličný, D. and Košťák, M.: Cretaceous basins of Central Europe: deciphering effects of global and regional processes – a short introduction, *Zeitschrift der Deutschen, Gesellschaft für. Geowissenschaften*, 165, 4, 495–499, 2014.
- 635 Wojewoda, J.: Sukcesja litoralno-szelfowa górnej kredy na obszarze niecki śródsudeckiej i rowu górnej Nysy w Sudetach, In: Wojewoda, J. (Ed.) *Obszary źródłowe: Zapis w osadach*, I, 81–96, WIND, Wrocław (in Polish), 1997.
- Wojewoda, J.: Neotectonic aspect of the Intrasudetic Shear Zone. *Acta Geodynamica et Geomaterialia*, Vol. 4, No. 4 (148), 31-41, 2007.
- Wojewoda, J.: Žďárky-Pstrážna Dome: a strike-slip fault-related structure at the eastern termination of the Poříčí-Hronov Fault Zone (Sudetes), *Acta Geodynamica et Geomaterialia*, 6, 273–290, 2009.
- 640 Wojewoda, J., Koszela, S. and Aleksandrowski, P.: A kilometre-scale low-angle detachment related to strike-slip faulting in Late Cretaceous mudstones of the Table Mountains (Central Sudetes, SW Poland). In: *Central European Tectonic Studies Group (CETEG) 8th Meeting, 22-25 April 2010, Mąchocice Kapitulne, Conference Proceedings* (ed. by M. Ludwiniak, A. Konon & A. Żylińska), University of Warsaw and Polish Geological Institute - National Research Institute, 127-128, 2010.
- 645 Wojewoda, J. and Kowalski, A.: Rola południowosudeckiej strefy ścinania w ewolucji Sudetów (The role of the South-Sudetic Shear Zone in the evolution of the Sudetes)., In: Wojewoda, J. and Kowalski, A. (eds): *Wyzwania Polskiej Geologii*,




3. Polski Kongres Geologiczny, Przewodnik do wycieczek kongresowych, Wycieczka 2.2., Polskie Towarzystwo Geologiczne, Wrocław, 28-43, 2016.
- Ziegler, P.A.: Late Cretaceous and Cenozoic intra-plate compressional deformations in the Alpine foreland-a geodynamic  
650 model, *Tectonophysics*, 137, 389-420, 10.1016/0040-1951(87)90330-1, 1987.
- Ziegler, P. A., Cloetingh, S. and van Wees, J. D.: Dynamics of intra-plate compressional deformation: the Alpine foreland and other examples, *Tectonophysics*, 252, 7–59, 1995.
- Żelaźniewicz, A. and Aleksandrowski, P.: Regionalizacja tektoniczna Polski: Polska południowo-zachodnia (Tectonic subdivision of Poland: southwestern Poland), *Przegląd Geologiczny*, 56, 904-911 (in Polish, English abstract), 2008.
- 655 Żelaźniewicz, A., Aleksandrowski, P., Buła, Z., Karnkowski, P.H., Konon, A., Oszczypko, N., Ślącza, A., Żaba, J. and Żytko, K.: Regionalizacja geologiczna Polski, Komitet Nauk Geologicznych PAN, Wrocław, 60, 2011.
- Żelaźniewicz, A. and Markiewicz, A.: Struktury ekstensyjne w cechsztyńskich ewaporatach monokliny przedsudeckiej a strefa tektoniczna Odry, *Przegląd Geologiczny*, 10, 463-471, 1991.


660

665


670



675 **Figure 1: The main Late Cretaceous – Early Palaeogene tectonic structures in Central Europe. Modified after Kley and Voigt 2008.. Abbreviations: Lf - Lusatian Main Fault, Od - Odra Fault, Ha-Harz Block, F – Flechtingen High, TW - Thuringian Forest, FL – Franconian Line, LB - Lusatian Block, SB - Sudetic Block, FSB - Fore-Sudetic Block, NBCB - North Bohemian Cretaceous Basin, HCM - Holy Cross Mountains**




**Figure 2: Outline geology of the Bohemian Massif NE margin (SW Poland and northern Czechia). Simplified from Dadlez (2000). NKG – Nysa-Králiky Graben. Red boxes show location of Figs. 6, 11a, 12 and 16.**




680


**Figure 3:** Schematic regional cross sections across the Bohemian Massif NE margin (highly vertically exaggerated). Location in Figure 2.



685 **Figure 4: Two seismic profiles from the Fore-Sudetic Homocline (location in Figure 2) showing very gentle folds in Triassic and Permian (Rotliegend not marked). Note steep to low-angle mainly reverse faults. The anticlines are structural traps for hydrocarbons, with gas fields known in Rotliegend strata (e.g. Grochowice gas field in Figure 4a). Please note the different scales of the two profiles.**



690 **Figure 5: Examples of tectonic structures observed on galleries' walls in copper and silver mines on the Fore-Sudetic Homocline at the base Zechstein ore-bearing formation (location in Figure 2). (a) Inverted steep fault, originally normal. The inversion caused fault-drag of Ca1 Zechstein Limestone. (b) Thrust fault in carbonate rock. Abbreviations: Bs- Weissliegend Sandstone, T1-Kupferschiefer Shale, Ca1-Zechstein Limestone.**



695 **Figure 6: Geological map of the North Sudetic Synclinorium (based on Balazińska and Bossowski, 1979; Krentz et al., 2001 and Cymerman, 2010). Locations of interpreted seismic profiles are shown. For full-length location of seismic profile of Figure 8a - see Figure 2. The map is based on Solecki (1994) and Cymerman (2010).**





Figure 7: Kurt Beyer's (1939) interpretation of the "young Saxonian" tectonics in the North Sudetic Synclinorium (Basin).

(a) Main tectonic deformation stages in the North Sudetic Basin. (b) Block-diagram showing simple open structure of the western part of the North Sudetic Synclinorium and complex pattern of blocks and synclinal grabens in its eastern part.

700




**Figure 8:** Folds in Muschelkalk in a quarry at Raciborowice (location given in Figure 6). (a) photo courtesy of Andrzej Solecki.



710 **Figure 9 a-c: Interpreted seismic profiles from the North-Sudetic Synclinorium (location in Figure 6).**

715



720

Figure 10 a-d: Interpreted seismic profiles from the North-Sudetic Synclinorium (location in Figure 6).




Figure 11: Simplified geological map and cross-sections of the Wleń Graben according to Kowalski (2020)






Figure 12: Geological map of the Intra-Sudetic Synclinorium (based on Grocholski and Augustyniak, 1971, Bossowski and Ichnatowicz, 2006, and Cymerman, 2010). Location of the interpreted seismic profiles and described localities is shown. CZT – Czerwieńczyce Trough (inverted graben).





730 **Figure 13: Seismic profiles from Mioszów vicinities showing probable effects of Late Cretaceous – Early Palaeogene reactivation of a Carboniferous flower structure. The Permian volcanic member (red) continues to the NNE to outcrop massively in the Suche Mountains. Location is given in Figure 12.**




735

**Figure 14 a,b:** (a) Seismic profile from vicinities of Broumov and its interpretation (made on printed paper version – Brada et al., 1982). Location in Fig. 12. The section shows that the Broumov well was drilled in the hanging-wall of a thrust fault and, hence, the Carboniferous in the neighbourhood may occur deeper than found in the well. The thrust continues to the NE parallel to the bedding – most probably along the Permian Anthracosia Shale member.

740





745




Figure 14 c-f: Seismic profiles from vicinities of Radków. Location in Fig. 12.





Figure 15: Geological cross-section of the Czerwieńczyce Reverse Graben after Oberc (1972). Location in Fig. 12.



750

Figure 16: Perspective view towards the north onto the base Cretaceous relief (known from boreholes) in the Upper Nysa-Králiky Graben (from Badura and Rauch, 2014). The oblique-striking blocks reflect NW-SE trending Late-Cretaceous – Early Palaeogene inverted blocks on the Cretaceous graben floor and folds in its fill (removed from the picture), once continuing laterally, before the graben formed in Miocene times. Location given in Fig. 2.




Figure 17: Schematic section across the Zieloniec Thrust (based on data from Cymerman, 1990, and Kozdrój, 2014). Location in Fig. 12.




Figure 18: Examples of outcrop-scale brittle structures related to major faults at the SW boundary of the Intra-Sudetic Synclinorium.

(a) The Pořiči-Hronov Fault, interpreted as high-angle reverse one, near the village of Malé Svatoňovice: tectonic contact between nearly vertically dipping Upper Cretaceous sandstones in the SW and the Permian sandstone in the NE.

(b) Shallow NW-dipping detachment along the bedding of Middle Turonian mudstones and calcareous claystones in a road cut near the village of Lewin Klodzki, probably related to the NW continuation of the Zieloniec Thrust. Location given in Fig. 12.





Figure 19: Schematic sections across the Lusatian Thrust in three localities, showing along-strike structure changes (based on Coubal et al., 2014). Location in Fig. 2.



770 **Figure 20:** Comparison of stereograms and rose diagrams of the dominant steep joint sets coming from several local studies (by Jerzykiewicz et al., 1974, Aleksandrowski, 1976, Solecki, 2011, Selerowicz et al., 2014) made at various locations in the Polish Sudetes. Background: the geological map of Figure 2.